

AAB MOLECULAR DIAGNOSTICS PRACTICE TEST (SAMPLE)

STUDY GUIDE



**SAMPLE STUDY GUIDE
WITH LIMITED QUESTIONS**

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://aabmoleculardiagnosics.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. What is heterogeneous nuclear RNA primarily composed of?**
 - A. Introns and exons
 - B. Only exons
 - C. Only introns
 - D. Messenger RNA only
- 2. What is the primary function of heterochromatins in relation to protein access?**
 - A. To enhance transcription speed
 - B. To provide a more compact structure for protein access
 - C. To promote gene expression
 - D. To facilitate DNA repair
- 3. What is the primary testing platform for monitoring response to antiviral therapy for HCV?**
 - A. ELISA
 - B. Qualitative PCR
 - C. Quantitative PCR
 - D. Western Blot
- 4. What type of control monitors specific steps of an amplification method?**
 - A. Verification control
 - B. Internal control
 - C. External control
 - D. Positive control
- 5. What is the role of linker histone H1 in the nucleosome structure?**
 - A. To degrade DNA
 - B. To bind the DNA between nucleosomes
 - C. To replicate DNA
 - D. To promote transcription

- 6. What is an example of a missense mutation?**
- A. AAA to AAG
 - B. UGC to UCC
 - C. CAA to CUA
 - D. GUA to GUC
- 7. What characterizes euchromatin in terms of its structure?**
- A. Highly compact and inactive
 - B. Less compact and accessible for transcription
 - C. Only found in non-dividing cells
 - D. Only present in inactive genes
- 8. Which of the following represents an example of an X-linked dominant disorder?**
- A. Wiskott-Aldrich syndrome
 - B. Prader-Willi syndrome
 - C. Becker muscular dystrophy
 - D. Fragile-X syndrome
- 9. Which of the following sequences is not characteristic of a eukaryotic polymerase II promoter?**
- A. TATAAA
 - B. TTGACA
 - C. CAAT box
 - D. GC-rich box
- 10. What is the condition resulting from Hemochromatosis?**
- A. Deficient iron absorption
 - B. Excess iron regulation
 - C. Increased calcium absorption
 - D. Chromosome misallocation

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. B
6. B
7. B
8. D
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is heterogeneous nuclear RNA primarily composed of?

- A. Introns and exons**
- B. Only exons
- C. Only introns
- D. Messenger RNA only

Heterogeneous nuclear RNA (hnRNA) is primarily composed of both introns and exons. This form of RNA is found in the nucleus of eukaryotic cells and represents the initial transcripts synthesized from DNA before they undergo processing. During this processing, introns—which are non-coding sequences—are removed, and exons—the coding sequences—are spliced together to form mature messenger RNA (mRNA). The presence of introns and exons in hnRNA is critical because it reflects the early stage of gene expression where the full RNA product is being prepared for translation. Exons will ultimately be retained in the final mRNA, while introns are excised. Therefore, understanding this composition is essential in molecular biology to comprehend the mechanisms of gene regulation and expression. In contrast, the other options either suggest a composition of only one type of sequence (either exons or introns) or imply that hnRNA consists solely of mature mRNA, which disregards the introns present in the primary transcript prior to its maturation.

2. What is the primary function of heterochromatins in relation to protein access?

- A. To enhance transcription speed
- B. To provide a more compact structure for protein access**
- C. To promote gene expression
- D. To facilitate DNA repair

The primary function of heterochromatins in relation to protein access is indeed best captured by the idea of providing a more compact structure for protein access. Heterochromatin represents regions of DNA that are tightly packed and often transcriptionally inactive. This compact structure serves to regulate access to the underlying genetic material, impacting how easily proteins, including transcription factors and other regulatory molecules, can interact with the DNA. Heterochromatin plays a crucial role in maintaining genome stability and regulating gene expression by controlling which genes are accessible for transcription. The dense compaction of these regions can effectively limit protein access, preventing the transcription machinery from binding and initiating gene expression. This function is essential for processes like maintaining cell identity and making sure that only the necessary genes are expressed in a particular cell type. While options related to transcription speed, gene expression promotion, and DNA repair touch on important cellular processes, they do not accurately describe the specific role of heterochromatin. Heterochromatin is known for its role in gene silencing and structural maintenance rather than enhancing transcription or directly facilitating DNA repair processes.

3. What is the primary testing platform for monitoring response to antiviral therapy for HCV?

- A. ELISA
- B. Qualitative PCR
- C. Quantitative PCR**
- D. Western Blot

The primary testing platform for monitoring response to antiviral therapy for Hepatitis C Virus (HCV) is quantitative PCR. This method quantifies the amount of viral RNA present in a patient's blood, providing critical information about the viral load at various points during treatment. Monitoring the change in viral load over time is essential in determining the effectiveness of the antiviral therapy. A significant decrease in viral load suggests that the treatment is working effectively, while a stable or increasing viral load may indicate treatment failure. Quantitative PCR is particularly important because it can detect even low levels of viral RNA, allowing healthcare providers to assess not just the presence of the virus but also the degree to which it is being suppressed by the therapy. This information is crucial for making informed decisions regarding continuing or adjusting treatment plans. In contrast, while ELISA can be used for initial screening and qualitative PCR can confirm the presence of HCV RNA, they do not provide the detailed quantification needed to evaluate therapeutic response comprehensively. Western blotting is generally not used for monitoring antiviral therapy response; it is primarily a confirmatory test for viral antibodies rather than for assessing active infection or monitoring treatment effectiveness.

4. What type of control monitors specific steps of an amplification method?

- A. Verification control
- B. Internal control**
- C. External control
- D. Positive control

The correct answer is internal control. Internal controls play a critical role in ensuring the reliability of amplification methods, such as polymerase chain reaction (PCR). They are incorporated into the assay to monitor the entire process from sample extraction through amplification, helping to verify that every step of the procedure is functioning correctly. An internal control typically consists of a target sequence that is co-amplified with the specific target of interest. This allows the laboratory to confirm that the amplification process is effective and that the reagents and conditions are suitable. If the internal control fails to amplify, it indicates a potential issue with the sample preparation or the amplification process itself, prompting a re-evaluation of the procedure. While other types of controls exist, such as positive controls which help confirm that the assay can detect the target under optimal conditions, and external controls which may provide information about the performance of the method across different runs, they do not specifically monitor the internal steps of the amplification method like internal controls do. Therefore, the emphasis on monitoring specific steps of the amplification process distinctly aligns with the function of internal controls.

5. What is the role of linker histone H1 in the nucleosome structure?

- A. To degrade DNA
- B. To bind the DNA between nucleosomes**
- C. To replicate DNA
- D. To promote transcription

The role of linker histone H1 in the nucleosome structure is primarily to bind the DNA between nucleosomes. H1 binds to the DNA that runs between adjacent nucleosomes, stabilizing the overall structure of the chromatin. By doing so, it helps to maintain the higher-order organization of DNA, facilitating the formation of the 30-nanometer fiber structure that plays a critical role in packing DNA into a compact form. This binding is crucial for the structural integrity of chromatin, effectively promoting the organization of genetic material in a way that balances accessibility for transcription and replication with the need for a compact storage form within the nucleus. H1's presence helps reinforce the interactions between nucleosomes, making it less accessible for transcription machinery when necessary, but also allowing for dynamic changes in chromatin structure that influence gene expression and DNA replication processes. Linker histone H1 does not degrade DNA, replicate DNA, or directly promote transcription. These functions are associated with other proteins and enzymes in the cell. For instance, certain nucleases are responsible for DNA degradation, while DNA polymerases carry out replication, and transcription factors mediate transcription. Thus, H1's specific role as a stabilizer between nucleosomes is what distinguishes its function within the nucleosome

6. What is an example of a missense mutation?

- A. AAA to AAG
- B. UGC to UCC**
- C. CAA to CUA
- D. GUA to GUC

A missense mutation occurs when a single nucleotide change results in the coding of a different amino acid. This type of mutation can potentially alter the function of a protein, depending on the role of the changed amino acid in the protein's structure and activity. In the given scenario, transforming UGC to UCC leads to a change from cysteine (Cys) to serine (Ser), representing a missense mutation. This change affects the amino acid sequence and, thus, can have functional implications for the resulting protein. Other options involve changes in nucleotide sequences, but they either lead to no change in the amino acid sequence (synonymous mutations) or to premature stop codons (nonsense mutations) rather than coding for a different amino acid. Hence, the correct choice exemplifies the concept of a missense mutation effectively, highlighting the significance of single nucleotide alterations in protein coding sequences.

7. What characterizes euchromatin in terms of its structure?

- A. Highly compact and inactive
- B. Less compact and accessible for transcription**
- C. Only found in non-dividing cells
- D. Only present in inactive genes

Euchromatin is characterized by being less compact and more accessible for transcription. This structural property allows for the active expression of genes, as the less dense packing of DNA facilitates the binding of transcription factors and RNA polymerase to the DNA. The open conformation of euchromatin is crucial for transcriptional activity, enabling genes located in these regions to be readily transcribed into RNA. In contrast, highly compacted forms of chromatin, such as heterochromatin, are generally associated with inactive or silent genes, as they restrict access to the DNA for transcription machinery. Furthermore, euchromatin is not limited to any specific cell type; it can be found in both dividing and non-dividing cells. It is also not exclusive to active genes, as regulatory regions of inactive genes can still reside in euchromatin, exemplifying its diverse role in gene regulation and accessibility. Thus, the major distinction of euchromatin lies in its functional and structural properties that promote gene expression.

8. Which of the following represents an example of an X-linked dominant disorder?

- A. Wiskott-Aldrich syndrome
- B. Prader-Willi syndrome
- C. Becker muscular dystrophy
- D. Fragile-X syndrome**

Fragile-X syndrome is an example of an X-linked dominant disorder, which means that the gene responsible for the disorder is located on the X chromosome and that only one copy of the mutated gene is sufficient to present with the disorder in both males and females. Since females have two X chromosomes, they can be affected even if only one of their X chromosomes carries the mutation. This is in contrast to X-linked recessive disorders, where typically males are more frequently affected because they have only one X chromosome. Fragile-X syndrome specifically results from a mutation in the FMR1 gene, which leads to developmental issues, intellectual disabilities, and various behavioral challenges. The inheritance pattern seen in Fragile-X syndrome is indicative of X-linked dominant disorders as affected individuals can have an affected parent. In this context, it's crucial to recognize that the disorder can be passed from an affected mother to her children regardless of their gender, resulting in various levels of expression in both males and females. Other options to consider involve different inheritance patterns. Wiskott-Aldrich syndrome is X-linked recessive, Prader-Willi syndrome is associated with imprinting on chromosome 15 and not linked to sex chromosomes, and Becker muscular dystrophy is also X-linked recessive.

9. Which of the following sequences is not characteristic of a eukaryotic polymerase II promoter?

- A. TATAAA
- B. TTGACA**
- C. CAAT box
- D. GC-rich box

A eukaryotic polymerase II promoter typically comprises several key elements that are important for the initiation of transcription. Among the options provided, the sequence that is not characteristic of this type of promoter is indeed the TTGACA sequence. The TATAAA box, commonly referred to as the TATA box, is a well-known core promoter element found in many eukaryotic promoters that helps position the RNA polymerase II enzyme for transcription initiation. It is usually located about 25 to 30 base pairs upstream of the transcription start site. The CAAT box is another important sequence element that can be found in eukaryotic promoters. It is generally located further upstream and plays a role in enhancing the transcriptional activity of that promoter. Similarly, a GC-rich box is associated with many eukaryotic promoters, particularly those that are constitutively active. This element contains a high frequency of guanine (G) and cytosine (C) residues, which can also contribute to the regulation of gene expression. In contrast, the TTGACA sequence is a characteristic sequence found in prokaryotic promoters, specifically as part of the -35 region recognized by sigma factors in bacterial RNA polymerase. This sequence does not typically appear

10. What is the condition resulting from Hemochromatosis?

- A. Deficient iron absorption
- B. Excess iron regulation**
- C. Increased calcium absorption
- D. Chromosome misallocation

Hemochromatosis is a condition characterized by the body's excessive absorption of iron from the diet, leading to iron overload. This excessive iron accumulation occurs because the regulatory mechanisms that normally control iron homeostasis are impaired. In a healthy individual, iron is absorbed according to the body's needs, and any excess is typically managed by hepcidin, a hormone that regulates iron metabolism. In individuals with hemochromatosis, mutations in genes responsible for iron regulation (most commonly the HFE gene) lead to an increased amount of iron being absorbed from food, even when the body's iron stores are already sufficient or elevated. This accumulation can cause serious damage to organs, including the liver, heart, and pancreas, leading to complications such as liver cirrhosis, heart disease, and diabetes. Given this understanding, the choice that indicates "excess iron regulation" accurately describes the pathological process in hemochromatosis, where there is not just excess absorption, but also an inability of the body to regulate iron levels effectively. This results in systemic iron overload rather than a deficiency or misallocation of other elements like calcium.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://aabmoleculardiagnosics.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE