AAB MOLECULAR DIAGNOSTICS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://aabmoleculardiagnostics.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What best describes a paracentric inversion?

A. It includes sequences involving both arms of a chromosome

B. It includes the centromere in the inverted region

C. It involves sequences within one arm of the chromosome

D. It changes the overall number of chromosomes

2. Which technique is used for engraftment testing following bone marrow transplantation?

A. SNP analysis

B. STR identification by PCR

C. Western blotting

D. Gel electrophoresis

3. How do small RNA molecules, like siRNA and saRNA, function in gene regulation?

A. By destroying nucleic acids

B. By promoting gene silencing exclusively

C. By inducing gene expression or silencing

D. By unwinding DNA for transcription

4. What is the key characteristic of DNA fragmentation in Maxam-Gilbert Sequencing?

A. It is labeled at the 3′ end

B. Chemical treatment generates breaks at all four nucleotide bases

C. Only one nucleotide base is treated in all reactions

D. The DNA fragment to be sequenced is radioactively labeled at the 5′ end

5. Which type of reagents are used for research purposes only?

A. ASR

B. RUO

C. GMP

D. IUO

SAMPLE

6. What is the primary screening method used to confirm a diagnosis of cystic fibrosis (CF)?

A. Genetic testing

B. Sweat test

C. Blood test

D. Chest X-ray

7. What is the primary function of the polyA tail in eukaryotic mRNA?

A. Stabilization of the mRNA

B. Translation initiation

C. Function is unknown

D. Facilitation of splicing

8. High stringency in nucleic acid hybridization is characterized by which of the following conditions?

A. Low temperature, high salt

B. High temperature, low salt, high formamide

C. Room temperature, medium salt

D. Low temperature, no formamide

9. What is the primary function of heterochromatins in relation to protein access?

A. To enhance transcription speed

B. To provide a more compact structure for protein access

C. To promote gene expression

D. To facilitate DNA repair

10. Which type of genetic mutation is primarily discussed in relation to hereditary hemochromatosis?

A. Point mutation

B. Frameshift mutation

C. Chromosomal translocation

D. All of the above

SAMPLE

Answers

SAMPLE

1. C
2. B
3. C
4. D
5. B
6. B
7. A
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What best describes a paracentric inversion?

A. It includes sequences involving both arms of a chromosome

B. It includes the centromere in the inverted region

C. It involves sequences within one arm of the chromosome

D. It changes the overall number of chromosomes

A paracentric inversion is a specific type of chromosomal rearrangement where a segment of a chromosome is inverted, and this inversion occurs while excluding the centromere. This means that the inverted segment is located entirely within one arm of the chromosome, either the short arm (p) or the long arm (q), but not both. By maintaining this confinement within a single arm and not incorporating the centromere, the structure and function of the chromosome are altered without impacting the centromere's positioning. The other options do not accurately describe a paracentric inversion. Sequences involving both arms would suggest a pericentric inversion, which includes the centromere. An implication of changing the overall number of chromosomes relates more closely to chromosomal duplications or deletions rather than to inversions, which do not change the chromosome count but can affect genetic expression and stability. The reference to including the centromere directly contradicts the definition of a paracentric inversion, as it specifically pertains to inversions that exclude it. Thus, the defining characteristic of a paracentric inversion is that it definitively involves sequences within one arm of the chromosome.

2. Which technique is used for engraftment testing following bone marrow transplantation?

A. SNP analysis

B. STR identification by PCR

C. Western blotting

D. Gel electrophoresis

The technique used for engraftment testing following bone marrow transplantation is STR (Short Tandem Repeat) identification by PCR (Polymerase Chain Reaction). This method is particularly effective in determining the presence and quantity of donor cells in the recipient's bone marrow post-transplantation. ENGRAFTMENT TESTING involves analyzing genetic markers that differ between the donor and recipient. STRs are highly polymorphic regions of DNA that vary in length between individuals, making them excellent targets for distinguishing between donor and recipient cells. By amplifying these STR regions through PCR, clinicians can assess the proportion of donor versus recipient DNA, thereby determining the success of the transplant and whether the donor marrow has engrafted effectively. The other techniques mentioned do not serve the same purpose in this context. SNP (Single Nucleotide Polymorphism) analysis also assesses genetic variations but is less commonly used for engraftment testing in practical settings due to its complexity and the fact that the variations may not be as numerous as STRs for distinguishing between the two genetic profiles. Western blotting is a technique used to detect specific proteins and is not applicable for analyzing DNA or monitoring engraftment. Gel electrophoresis is a method used to separate DNA or protein fragments but is typically used as a supportive technique

SAMPLE

3. How do small RNA molecules, like siRNA and saRNA, function in gene regulation?

A. By destroying nucleic acids

B. By promoting gene silencing exclusively

C. By inducing gene expression or silencing

D. By unwinding DNA for transcription

Small RNA molecules, such as small interfering RNA (siRNA) and small activating RNA (saRNA), play a crucial role in gene regulation through their ability to modulate gene expression. siRNAs primarily function by targeting specific messenger RNA (mRNA) sequences for degradation, which effectively silences gene expression. They achieve this through a process known as RNA interference (RNAi), where the siRNA guides the RNA-induced silencing complex (RISC) to the complementary mRNA, leading to its destruction. On the other hand, saRNAs are involved in promoting gene expression rather than silencing it. They can enhance the transcription of specific genes by binding to their promoters, facilitating recruitment of the transcriptional machinery and thus increasing mRNA production. Therefore, the correct choice reflects the dual role that small RNAs can play—they can induce gene expression in some contexts (as with saRNA) while also silencing genes in other contexts (as with siRNA). This versatility in function underlines the complexity of gene regulation mechanisms and showcases the significance of small RNA molecules in cellular functions.

4. What is the key characteristic of DNA fragmentation in Maxam-Gilbert Sequencing?

A. It is labeled at the 3' end

B. Chemical treatment generates breaks at all four nucleotide bases

C. Only one nucleotide base is treated in all reactions

D. The DNA fragment to be sequenced is radioactively labeled at the 5' end

The key characteristic of DNA fragmentation in Maxam-Gilbert Sequencing is that the DNA fragment to be sequenced is radioactively labeled at the 5' end. In this method, the DNA is first labeled to allow for detection of the fragments that are generated through specific chemical treatments. By tagging the 5' end, the resultant fragments produced through cleavage can be easily identified during the separation and analysis steps that follow. This labeling is crucial because it provides a way to visualize the fragments in later steps, typically using gel electrophoresis. The unique aspect of labeling at the 5' end in Maxam-Gilbert Sequencing is contrasted with other sequencing methods where 3' end labeling might be employed. This characteristic ensures that the specific fragments generated during the sequencing reactions can be accurately tracked and resolved. In contrast, the other options do not correctly define the primary characteristic of this sequencing methodology. For instance, while chemical treatment does indeed generate cuts at nucleotide bases, the specificity and method of labeling are particularly notable in Maxam-Gilbert Sequencing. Thus, the radioactively labeled 5' end is critically relevant in the context of this sequencing technique.

SAMPLE

5. Which type of reagents are used for research purposes only?

A. ASR

B. RUO

C. GMP

D. IUO

The correct answer is "RUO," which stands for "Research Use Only." These reagents are specifically intended for use in research settings and are not approved for diagnostic or therapeutic purposes in humans. They are often utilized in laboratories to explore new technologies, validate new assays, or conduct preliminary studies that can provide insights leading to further developments in molecular diagnostics. RUO reagents are important because they allow researchers to experiment and gather data without the stringent regulatory requirements placed on diagnostic products. This flexibility can accelerate the pace of innovation in the field of molecular diagnostics. In contrast, ASR (Analyte Specific Reagent) is designed for the laboratory use of specific analytes and is subject to quality controls set by regulatory bodies. GMP (Good Manufacturing Practice) denotes products that are manufactured under stringent regulations for clinical use, ensuring safety and efficacy. IUO (Investigational Use Only) refers to products under investigation that may be used in clinical trials but are not yet approved for general clinical use. These categories emphasize the varying levels of regulatory compliance and intended use, highlighting the research-specific focus of RUO reagents.

6. What is the primary screening method used to confirm a diagnosis of cystic fibrosis (CF)?

A. Genetic testing

B. Sweat test

C. Blood test

D. Chest X-ray

The primary screening method used to confirm a diagnosis of cystic fibrosis (CF) is the sweat test. This test measures the concentration of chloride in sweat. Individuals with cystic fibrosis have defects in the CFTR gene, which affects the transport of chloride and sodium ions across epithelial cell membranes. As a result, people with CF typically have elevated levels of chloride in their sweat. The sweat test is considered the gold standard for diagnosing cystic fibrosis due to its specificity and reliability. A chloride concentration of greater than 60 millimoles per liter is typically indicative of the disease. This direct method helps clinicians assess the functional impact of CFTR mutations. In contrast, while genetic testing can identify specific mutations in the CFTR gene, it is usually utilized to confirm a diagnosis rather than serve as the primary screening method. Blood tests may help rule out other conditions but are not specific for cystic fibrosis. Similarly, chest X-rays can indicate lung issues associated with CF, but they do not confirm the presence of the disease. Thus, the sweat test remains the most effective and widely accepted method for diagnosing cystic fibrosis.

SAMPLE

7. What is the primary function of the polyA tail in eukaryotic mRNA?

A. Stabilization of the mRNA

B. Translation initiation

C. Function is unknown

D. Facilitation of splicing

The polyA tail primarily stabilizes mRNA. By adding a string of adenine nucleotides at the 3' end, the transcript becomes more resistant to exonucleases, which slows its degradation in the cytoplasm and lengthens how long the message can be used to make protein. This increased stability often translates to higher protein output because the mRNA remains available longer for translation. The tail also helps with efficient translation by binding poly(A)-binding protein, which fosters a close loop between the 3' end and the 5' cap through interactions with translation factors. This looping promotes translation efficiency and ribosome recycling, but the fundamental, primary effect remains the extended lifetime of the mRNA. Splicing occurs earlier in processing and is not driven by the polyA tail; the tail is added after transcription termination. The idea that its function is unknown is incorrect.

8. High stringency in nucleic acid hybridization is characterized by which of the following conditions?

A. Low temperature, high salt

B. High temperature, low salt, high formamide

C. Room temperature, medium salt

D. Low temperature, no formamide

High stringency in nucleic acid hybridization refers to conditions that promote more specific binding between complementary nucleic acid sequences while minimizing non-specific interactions. This is typically achieved through the combination of elevated temperatures, reduced salt concentrations, and possibly the addition of components like formamide, which destabilize mismatched duplexes. Under high temperature conditions, the stability of the hybridized DNA or RNA is affected, leading to the denaturation of any non-perfectly matched strands. A lower salt concentration contributes to the weakening of ionic interactions that stabilize hybrid duplexes, further ensuring that only perfectly matched sequences remain hybridized. The inclusion of a substance like formamide can disrupt hydrogen bonding to some extent, thus helping to enhance stringency. For these reasons, high temperature and low salt concentration, along with the presence of formamide, effectively support specific hybridization conditions, making this option the correct choice for high stringency.

SAMPLE

9. What is the primary function of heterochromatins in relation to protein access?

A. To enhance transcription speed

B. To provide a more compact structure for protein access

C. To promote gene expression

D. To facilitate DNA repair

The primary function of heterochromatins in relation to protein access is indeed best captured by the idea of providing a more compact structure for protein access. Heterochromatin represents regions of DNA that are tightly packed and often transcriptionally inactive. This compact structure serves to regulate access to the underlying genetic material, impacting how easily proteins, including transcription factors and other regulatory molecules, can interact with the DNA. Heterochromatin plays a crucial role in maintaining genome stability and regulating gene expression by controlling which genes are accessible for transcription. The dense compaction of these regions can effectively limit protein access, preventing the transcription machinery from binding and initiating gene expression. This function is essential for processes like maintaining cell identity and making sure that only the necessary genes are expressed in a particular cell type. While options related to transcription speed, gene expression promotion, and DNA repair touch on important cellular processes, they do not accurately describe the specific role of heterochromatin. Heterochromatin is known for its role in gene silencing and structural maintenance rather than enhancing transcription or directly facilitating DNA repair processes.

10. Which type of genetic mutation is primarily discussed in relation to hereditary hemochromatosis?

A. Point mutation

B. Frameshift mutation

C. Chromosomal translocation

D. All of the above

Hereditary hemochromatosis is predominantly associated with point mutations, particularly in the HFE gene. A point mutation involves a change in a single nucleotide base pair in the DNA sequence. In the case of hereditary hemochromatosis, the most common mutation is a single base change (C to G) at position 282 of the HFE gene, leading to an amino acid substitution that disrupts iron regulation in the body. This single alteration can have significant downstream effects on iron metabolism, thereby resulting in the disease. While frameshift mutations and chromosomal translocations can cause various genetic disorders, they are not primarily linked to hereditary hemochromatosis in the same way that point mutations are. Frameshift mutations change the reading frame of the genetic code, often resulting in entirely different protein products, which is not the mechanism responsible for hereditary hemochromatosis. Chromosomal translocations involve segments of chromosomes being rearranged, which also does not contribute to the mutations specific to hereditary hemochromatosis. Thus, the focus on point mutations accurately reflects the underlying genetic cause of this inherited condition.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://aabmoleculardiagnostics.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE