AANP ADULT GERONTOLOGY PRIMARY CARE NURSE PRACTIOTIONER PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://aanp.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is a key feature of osteoporosis?

A. Increased bone density

B. Decreased bone density

C. Joint inflammation

D. Increased muscle mass

2. What technique is used to assess the psoas/iliopsoas muscle?

A. Flex the hip

B. Measure leg length

C. Draw the knee to the chest

D. Push against a wall

3. What does a peak expiratory flow rate (PEF) of 80-100% indicate for an asthma patient?

A. Severe asthma exacerbation

B. Green Zone, maintain treatment

C. Yellow Zone, increase maintenance therapy

D. Red Zone, call emergency services

4. Which factor does NOT precipitate gout attacks?

A. Seafood consumption

B. Excessive fluid intake

C. Alcohol consumption

D. Organ meats intake

5. What is the primary purpose of a fall risk assessment?

A. To evaluate medication adherence

B. To identify patients at risk for falls and implement preventive measures

C. To assess nutritional intake

D. To determine mobility issues

6. What is a contraindication for the use of potassium-sparing diuretics?

A. Severe hyperkalemia

B. Hypotension

C. Chronic kidney disease

D. Asthma

SAMPLE

7. What does the acronym "Frail" stand for in geriatric assessment?

A. Fatigue, response, ambulation, illness, and loss of weight

B. Fatigue, resistance, ambulation, illness, and loss of weight

C. Focus, resilience, activity, illness, and lack of weight

D. Fluid, resting, ambulation, injury, and loneliness

8. What should a patient report when taking Synthroid?

A. Decreased appetite

B. Excessive thirst

C. Palpitations

D. Skin rashes

9. What is the main concern with long-term steroid therapy?

A. Hypoglycemia

B. Hyperglycemia

C. Increased muscle strength

D. Improved wound healing

10. What laboratory parameter is most indicative of liver function?

A. Alkaline phosphatase

B. Alanine aminotransferase (ALT)

C. Creatinine

D. Bilirubin

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. B
5. B
6. A
7. B
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is a key feature of osteoporosis?

A. Increased bone density

B. Decreased bone density

C. Joint inflammation

D. Increased muscle mass

A key feature of osteoporosis is decreased bone density. This condition is characterized by a loss of bone mass and deterioration of bone tissue, leading to an increased risk of fractures. Osteoporosis typically occurs when the balance between bone resorption (breakdown) and bone formation shifts in favor of resorption, causing bones to become weaker and more porous. As bone density decreases, the structural integrity of bones is compromised, making them more fragile. This is particularly crucial for older adults, as they are at a higher risk for osteoporosis due to age-related changes in bone metabolism and hormonal shifts, particularly in postmenopausal women with decreased estrogen levels. In contrast, increased bone density, joint inflammation, and increased muscle mass are not characteristic of osteoporosis. Increased bone density suggests stronger bones, joint inflammation is more associated with conditions like arthritis, and increased muscle mass relates to overall fitness and strength rather than bone health.

2. What technique is used to assess the psoas/iliopsoas muscle?

A. Flex the hip

B. Measure leg length

C. Draw the knee to the chest

D. Push against a wall

The technique used to assess the psoas/iliopsoas muscle involves flexing the hip. This muscle is primarily responsible for hip flexion and is an important component in movements such as walking, running, and climbing stairs. When assessing the psoas muscle, the practitioner will often ask the patient to flex their hip against resistance. This action engages the psoas/iliopsoas muscle, allowing the practitioner to evaluate its strength and any potential tenderness or pain that may indicate an underlying condition, such as psoas muscle strain or irritation. Flexing the hip allows for the evaluation of the functional capacity of the psoas muscle, making it a direct assessment technique. Other movements, like measuring leg length or drawing the knee to the chest, do not specifically isolate the psoas/iliopsoas muscle in the same way. Pushing against a wall involves different muscle groups and assesses stability and strength rather than focusing on the hip flexors. Therefore, flexing the hip is the appropriate technique for assessing the condition and function of the psoas/iliopsoas muscle.

3. What does a peak expiratory flow rate (PEF) of 80-100% indicate for an asthma patient?

A. Severe asthma exacerbation

B. Green Zone, maintain treatment

C. Yellow Zone, increase maintenance therapy

D. Red Zone, call emergency services

A peak expiratory flow rate (PEF) of 80-100% indicates that the patient's lung function is within the expected range and suggests controlled asthma. This measurement falls within the "Green Zone," which is characterized by good control of asthma symptoms. In this zone, patients are encouraged to maintain their current treatment regimen, as they are effectively managing their condition. Regular monitoring of PEF can help patients recognize when they are stable and when changes in their condition may necessitate adjustments to their management plan. In contrast, PEF readings that fall below this range would typically indicate a worsening of asthma, warranting an increase in therapy or a need for emergency action, depending on severity. Thus, understanding PEF readings helps in the ongoing management of asthma and in fostering patient engagement in their own health care.

SAMPLE

4. Which factor does NOT precipitate gout attacks?

A. Seafood consumption

B. Excessive fluid intake

C. Alcohol consumption

D. Organ meats intake

Excessive fluid intake does not precipitate gout attacks, making it the correct choice in this context. Gout is a type of inflammatory arthritis caused by the accumulation of uric acid crystals in the joints. Factors that typically trigger gout attacks include dietary choices high in purines, as well as alcohol consumption, which can increase uric acid production and decrease its excretion. Seafood and organ meats are known to be high in purines, leading to elevated uric acid levels and potential gout flares. Alcohol, particularly beer and spirits, can also contribute to gout attacks for similar reasons. In contrast, adequate hydration and excessive fluid intake can actually help reduce uric acid levels in the body, promoting better kidney function and assisting in the excretion of uric acid. Thus, maintaining a high fluid intake is generally advised for individuals at risk for gout rather than being a precipitating factor.

5. What is the primary purpose of a fall risk assessment?

A. To evaluate medication adherence

B. To identify patients at risk for falls and implement preventive measures

C. To assess nutritional intake

D. To determine mobility issues

The primary purpose of a fall risk assessment is to identify patients at risk for falls and implement preventive measures. Falls are a significant health concern, particularly in older adults, leading to serious injuries such as fractures and head trauma. A comprehensive fall risk assessment involves evaluating various risk factors, including medical history, physical function, medication use, and environmental hazards. By identifying individuals at higher risk, healthcare providers can tailor interventions such as strength training, balance exercises, medication reviews, home safety evaluations, and education to mitigate these risks effectively. This proactive approach ultimately aims to enhance patient safety and prevent fall-related injuries. While evaluating medication adherence, assessing nutritional intake, and determining mobility issues are important components of overall patient health, they do not specifically focus on the multifaceted nature of fall risk. These other factors are often part of a broader assessment but are not the primary goal of a fall risk assessment.

6. What is a contraindication for the use of potassium-sparing diuretics?

A. Severe hyperkalemia

B. Hypotension

C. Chronic kidney disease

D. Asthma

Severe hyperkalemia is a significant contraindication for the use of potassium-sparing diuretics because these medications work by inhibiting sodium reabsorption in the renal tubules while conserving potassium. This action can exacerbate elevated potassium levels in the blood, leading to potentially life-threatening conditions such as cardiac arrhythmias. In patients who already have high potassium levels, the addition of a potassium-sparing diuretic could result in dangerously high levels of potassium, making it critical to avoid their use in this scenario. In contrast to severe hyperkalemia, the presence of hypotension, chronic kidney disease, or asthma typically does not outright contraindicate the use of potassium-sparing diuretics but may require careful monitoring and adjustment of medications based on the individual's overall health status and specific conditions. Therefore, the risk associated with using potassium-sparing diuretics in the context of severe hyperkalemia is markedly higher, highlighting the necessity of avoiding them in such cases.

SAMPLE

7. What does the acronym "Frail" stand for in geriatric assessment?

A. Fatigue, response, ambulation, illness, and loss of weight

B. Fatigue, resistance, ambulation, illness, and loss of weight

C. Focus, resilience, activity, illness, and lack of weight

D. Fluid, resting, ambulation, injury, and loneliness

The acronym "Frail" in the context of geriatric assessment stands for Fatigue, Resistance, Ambulation, Illness, and Loss of weight. Each of these components provides critical insights into the health status and functional capacity of older adults. Fatigue is an important aspect as it can indicate underlying problems and affects the overall quality of life. Resistance refers to both physical resistance in terms of strength and resilience against illness and challenges, which is essential for maintaining independence. Ambulation is about mobility, and difficulties in this area can lead to increased risk of falls and reduced ability to perform daily activities. Illness encompasses the various chronic conditions that older adults may have, influencing their overall health and functional status. Finally, Loss of weight is a significant indicator of malnutrition or underlying disease and is a common concern in the geriatric population. This framework helps health care providers identify frailty, assess risks for adverse outcomes, and implement appropriate interventions. Understanding this acronym assists in delivering holistic and tailored care to elderly patients, improving their quality of life and health outcomes.

8. What should a patient report when taking Synthroid?

A. Decreased appetite

B. Excessive thirst

C. Palpitations

D. Skin rashes

When a patient is taking Synthroid, which is a brand name for levothyroxine, it is important for them to report symptoms that may indicate they are experiencing side effects related to thyroid hormone replacement therapy. Palpitations are a significant side effect that patients may experience if they are on an inappropriate dosage or if their body is reacting to the medication. Palpitations can manifest as feelings of a racing heart, irregular heartbeat, or increased heart rate. These symptoms may indicate that the patient is receiving too high of a dose, leading to hyperthyroid-like symptoms, which can include cardiovascular effects such as increased heart rate and palpitations. Therefore, reporting these symptoms allows healthcare providers to assess the patient's thyroid hormone levels and adjust the medication as needed to ensure safe and effective treatment. In contrast, while decreased appetite, excessive thirst, and skin rashes might be concerning symptoms for other medications or conditions, they are not commonly associated with the use of Synthroid. Hence, they do not hold the same level of significance in the context of monitoring patients on this particular medication.

SAMPLE

9. What is the main concern with long-term steroid therapy?

A. Hypoglycemia

B. Hyperglycemia

C. Increased muscle strength

D. Improved wound healing

Long-term steroid therapy is primarily associated with the risk of hyperglycemia. Corticosteroids, such as prednisone, can increase blood sugar levels by stimulating gluconeogenesis in the liver and decreasing insulin sensitivity in muscle and adipose tissues. This effect can lead to elevated glucose levels in the blood, which can be particularly concerning for individuals who may already have conditions such as diabetes or those at risk of developing diabetes. In terms of the context for the other options, hypoglycemia is less commonly associated with steroid use, as steroids typically raise blood sugar levels rather than lower them. Improved muscle strength and enhanced wound healing are often seen as benefits of short-term steroid use in specific conditions; however, with chronic use, the opposite may occur due to steroid-induced muscle wasting and slowed tissue repair. Therefore, the main concern with long-term use of steroids lies significantly in the management of hyperglycemia and its related complications.

10. What laboratory parameter is most indicative of liver function?

A. Alkaline phosphatase

B. Alanine aminotransferase (ALT)

C. Creatinine

D. Bilirubin

Alanine aminotransferase (ALT) is considered one of the most sensitive indicators of liver function. This enzyme is primarily found in the liver, and its levels in the blood often increase in response to liver cell damage. When the liver is inflamed or injured, ALT is released into the bloodstream, making its measurement a valuable tool in assessing liver health. In clinical practice, elevated ALT levels can signify various liver conditions, such as hepatitis or fatty liver disease. Monitoring this enzyme helps healthcare providers evaluate the degree of liver injury and assess ongoing liver function. While alkaline phosphatase, creatinine, and bilirubin are important laboratory markers, they do not provide as clear a picture of liver function specifically. Alkaline phosphatase can indicate bile duct obstruction or certain bone disorders, creatinine is primarily a marker for kidney function, and bilirubin levels are more reflective of the liver's ability to conjugate and excrete bilirubin rather than overall liver function. Thus, ALT remains a key marker in the assessment of liver function.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://aanp.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE