ABC ORTHOTIC & PROSTHETIC PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://abcorthoticprosthetic.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which phase occurs just after initial contact in the gait cycle?

A. Mid stance

B. Terminal stance

C. Loading response

D. Preswing

2. How many secondary diagnoses are needed, in addition to diabetes, for Medicare coverage of shoes and inserts?

A. Four

B. Five

C. Six

D. Three

3. What characterizes a boutonniere deformity?

A. PIP hyperextension and DIP flexion

B. PIP flexion and DIP hyperextension

C. Thumb flexion and finger extension

D. Complete extension of all joints

4. Which organization oversees the rules relating to the safe use of potentially hazardous materials in orthotic fabrication?

A. Center for Disease Control (CDC)

B. Occupational Safety and Health Administration (OSHA)

C. Food and Drug Administration (FDA)

D. National Institutes of Health (NIH)

5. What forces act on the knee during single limb stance?

A. Compressive forces from hip extensor muscles

B. Valgus thrust on knee secondary to lateral shift

C. Tensile forces from knee flexors

D. Normal contact forces from the ground

SAMPLE

6. In a hand orthosis, what function does the C-Bar serve?

A. Wrist support

B. Thumb adduction stop

C. Finger extension aid

D. Palmar support

7. What is the primary orthotic goal for patients with Dens fractures?

A. Flexion stabilization

B. Hyperextension

C. External rotation assistance

D. Neutral alignment support

8. Which muscles assist with wrist extension?

A. Flexor digitorum superficialis and flexor carpi ulnaris

B. Extensor indicis and extensor carpi radialis longus

C. Palmaris longus and abductor pollicis brevis

D. Flexor pollicis longus and flexor carpi radialis

9. Which nerve innervates the wrist extensors?

A. Median Nerve

B. Ulnar Nerve

C. Radial Nerve

D. Musculocutaneous Nerve

10. For a patient with Fair (2/5) plantar flexion strength and good (4/5) dorsiflexion strength, which double adjustable configuration is recommended?

A. Anterior Pins and Posterior Springs

B. Posterior Pins and Anterior Springs

C. Fixed Ankle Joint

D. Single Adjustable AFO

SAMPLE

Answers

SAMPLE

1. C
2. C
3. B
4. B
5. B
6. B
7. B
8. B
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. Which phase occurs just after initial contact in the gait cycle?

A. Mid stance

B. Terminal stance

C. Loading response

D. Preswing

In the gait cycle, the phase that occurs just after initial contact is known as the loading response. This phase begins when the foot initially makes contact with the ground and continues until the opposite foot leaves the ground. During this period, the body begins to absorb the impact of the weight as it transitions from the initial contact phase to the next stages of gait. During the loading response, critical changes occur, including the knee flexing to help absorb shock and the foot adapting to the surface it is contacting. This phase is vital for establishing stability and distributing weight properly across the limb while preparing for the next phases of walking. Understanding this phase emphasizes its importance in maintaining balance and preventing falls, as it sets the foundation for smooth transitions into subsequent phases of gait.

2. How many secondary diagnoses are needed, in addition to diabetes, for Medicare coverage of shoes and inserts?

A. Four

B. Five

C. Six

D. Three

For Medicare coverage of therapeutic shoes and inserts for patients with diabetes, the requirement is that the patient has diabetes as the primary diagnosis and, in addition, at least one of a specified list of secondary diagnoses is necessary. The patient must have at least one qualifying secondary diagnosis, and in this context, a total of six specific conditions are recognized as necessary for coverage. This means that in addition to the primary diagnosis of diabetes, the documentation must include the presence of certain other medical conditions, which can support the need for therapeutic footwear. Therefore, having six secondary diagnoses in total fulfills the requirement for Medicare coverage alongside diabetes. This ensures that the patient not only has diabetes but also additional complications or risk factors that necessitate the use of therapeutic shoes.

3. What characterizes a boutonniere deformity?

A. PIP hyperextension and DIP flexion

B. PIP flexion and DIP hyperextension

C. Thumb flexion and finger extension

D. Complete extension of all joints

A boutonniere deformity is characterized by the specific positioning of the joints in the finger, primarily impacting the proximal interphalangeal (PIP) joint and the distal interphalangeal (DIP) joint. In this condition, there is flexion at the PIP joint while the DIP joint is in extension, which results from a rupture or disruption of the central slip of the extensor tendon. This particular presentation leads to a characteristic "buttonhole" appearance, hence the name "boutonniere." The flexion of the PIP joint occurs due to the imbalance of the tendon forces, while the DIP joint appears to be hyperextended due to the pull from the intact flexor tendons. Understanding this characteristic positioning is essential for proper diagnosis and treatment, as management may involve splinting, therapy, or surgery to restore function and alignment to the affected digits.

SAMPLE

4. Which organization oversees the rules relating to the safe use of potentially hazardous materials in orthotic fabrication?

A. Center for Disease Control (CDC)

B. Occupational Safety and Health Administration (OSHA)

C. Food and Drug Administration (FDA)

D. National Institutes of Health (NIH)

The Occupational Safety and Health Administration (OSHA) is the organization responsible for overseeing regulations concerning the safe use of potentially hazardous materials in various work environments, including orthotic fabrication. OSHA's mandate is to ensure safe and healthful working conditions by setting and enforcing standards and providing training, outreach, education, and assistance. In the context of orthotic fabrication, this means that materials used in the production process must comply with OSHA regulations regarding safety, handling, and exposure limits to ensure that workers are protected from occupational hazards associated with these materials. This may include proper ventilation, the use of personal protective equipment, and adherence to safety protocols when handling materials that may pose a risk to health. Understanding the role of OSHA helps professionals in the field of orthotics and prosthetics maintain a safe working environment, promoting the health and safety of both employees and patients.

5. What forces act on the knee during single limb stance?

A. Compressive forces from hip extensor muscles

B. Valgus thrust on knee secondary to lateral shift

C. Tensile forces from knee flexors

D. Normal contact forces from the ground

During single limb stance, the knee experiences various forces that impact its stability and function. The correct answer emphasizes the valgus thrust on the knee, which can occur due to a lateral shift of the body's center of mass. This lateral shift often happens when a person is standing on one limb, leading to an increased lateral load on the knee joint. In this scenario, the forces from other parts of the body, particularly the hip and torso, may create a mechanical advantage that leads to a valgus moment at the knee. This necessitates effective management by the knee's soft tissue structures and ligaments to maintain proper alignment and prevent injury. The other choices pertain to different types of forces that may not be as directly relevant in this specific context. Compressive forces from the hip extensor muscles are generally felt during activities involving hip extension rather than standing alone, while tensile forces from knee flexors are mainly engaged during knee flexion, not in a static stance. Normal contact forces from the ground do exist but do not specifically highlight the unique mechanics involved in a single limb stance as does the valgus thrust. Thus, the emphasis on the valgus thrust is key in understanding the forces acting on the knee in this particular position.

SAMPLE

6. In a hand orthosis, what function does the C-Bar serve?

A. Wrist support

B. Thumb adduction stop

C. Finger extension aid

D. Palmar support

The C-Bar in a hand orthosis primarily serves as a thumb adduction stop, which is crucial for maintaining the functional position of the thumb. This component of the orthosis helps to separate the thumb from the index finger, preventing unwanted movement that could lead to thumb adduction during grasping or pinching activities. By restricting adduction, the C-Bar ensures that the thumb is positioned properly to assist in gripping and functional hand movements, thereby enhancing the stability and effectiveness of the orthosis. In comparison, wrist support is mainly concerned with stabilizing the wrist joint and does not specifically address the position of the thumb. Similarly, finger extension aids focus on assisting the fingers in achieving extension, which is unrelated to the specific function of the C-Bar. Palmar support, while important for hand stability and comfort, also does not perform the same role as the C-Bar in preventing thumb adduction. Thus, the primary function of the C-Bar is integral to promoting optimal hand function by maintaining the thumb's necessary position.

7. What is the primary orthotic goal for patients with Dens fractures?

A. Flexion stabilization

B. Hyperextension

C. External rotation assistance

D. Neutral alignment support

The primary orthotic goal for patients with Dens fractures, also known as odontoid fractures, involves the provision of hyperextension. This is because these types of injuries often occur at the cervical spine, specifically affecting the second cervical vertebra (C2) where the Dens, or odontoid process, is located. Providing hyperextension through the use of orthoses helps to stabilize and realign the cervical spine, reducing the risk of further injury. Hyperextension is essential in this context because it assists in decompressing the spinal structures and can help prevent excessive movement that may aggravate the fracture site. Ensuring that the neck remains in a state of hyperextension allows for optimal healing by maintaining proper alignment of the C1 and C2 vertebrae, which is critical for restoring stability and function. Regarding the other options, while they may have relevance in different contexts or types of spinal injuries, they do not align with the primary goal of managing Dens fractures specifically. The focus for these patients is on maintaining hyperextension to promote healing and prevent complications.

SAMPLE

8. Which muscles assist with wrist extension?

A. Flexor digitorum superficialis and flexor carpi ulnaris

B. Extensor indicis and extensor carpi radialis longus

C. Palmaris longus and abductor pollicis brevis

D. Flexor pollicis longus and flexor carpi radialis

Wrist extension is primarily facilitated by muscles that are located on the dorsal or posterior aspect of the forearm. The extensor indicis and extensor carpi radialis longus are two such muscles that play a significant role in this action. The extensor carpi radialis longus originates from the lateral supracondylar ridge of the humerus and inserts into the base of the second metacarpal. Through its action, it extends and abducts the wrist. The extensor indicis, on the other hand, originates from the distal ulna and assists in extending the wrist while also extending the index finger. Together, these two muscles contribute effectively to wrist extension, making this answer correct. In contrast, the other options contain muscles that do not primarily contribute to wrist extension. For instance, the flexor digitorum superficialis and flexor carpi ulnaris are involved in wrist flexion rather than extension. Palmaris longus and abductor pollicis brevis also focus on wrist flexion and thumb abduction, respectively. Finally, the flexor pollicis longus and flexor carpi radialis also do not assist in wrist extension; they primarily facilitate wrist flexion or thumb flex

9. Which nerve innervates the wrist extensors?

A. Median Nerve

B. Ulnar Nerve

C. Radial Nerve

D. Musculocutaneous Nerve

The correct choice is the radial nerve, as it is primarily responsible for innervating the wrist extensors. The wrist extensors, which include muscles such as the extensor carpi radialis longus and brevis, as well as the extensor carpi ulnaris, are involved in the extension of the wrist joint. These muscles receive their motor supply from branches of the radial nerve that travel into the posterior compartment of the forearm. The role of the radial nerve in this context is crucial, as it not only supplies the muscles responsible for wrist extension but also has a significant function in finger extension. Damage to this nerve can lead to wrist drop, a condition where the individual is unable to extend the wrist due to loss of function in the extensors, further highlighting the importance of the radial nerve in upper limb function. Other nerves mentioned, such as the median and ulnar nerves, primarily innervate different muscle groups, including the wrist flexors and intrinsic hand muscles, which do not contribute to the extension of the wrist. The musculocutaneous nerve is primarily associated with the innervation of the biceps brachii and other flexors of the elbow, which is distinct from the role of wrist extensors.

SAMPLE

10. For a patient with Fair (2/5) plantar flexion strength and good (4/5) dorsiflexion strength, which double adjustable configuration is recommended?

A. Anterior Pins and Posterior Springs

B. Posterior Pins and Anterior Springs

C. Fixed Ankle Joint

D. Single Adjustable AFO

In the case of a patient exhibiting Fair (2/5) plantar flexion strength and Good (4/5) dorsiflexion strength, selecting a double adjustable configuration that effectively facilitates mobility while accommodating the patient's strength limitations is crucial. Choosing the anterior pins and posterior springs configuration allows for controlled plantar flexion, which is particularly beneficial for supporting the reduced strength in that movement. The posterior springs offer a dynamic response, enabling the foot to move into dorsiflexion more easily, thus enhancing stability during ambulation. This configuration actively assists the plantar flexion, providing support without entirely restricting movement, which is essential for this patient profile. Combining the strength the patient has in dorsiflexion with adjustable settings helps in achieving a more natural gait pattern. It allows for adaptations during walking, compensating for the weaker plantar flexion. This choice is particularly important since it allows the user to benefit from both active involvement in movement and necessary support for the weaker muscle groups, promoting a more functional and adaptive AFO.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://abcorthoticprosthetic.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE