

ABRET ELECTROENCEPHALOGRAPHY (EEG) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE

<https://abreteeg.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Which cranial nerve primarily controls ocular movements?**
 - A. II
 - B. III
 - C. IV
 - D. VI

- 2. What is the most likely EEG finding in a patient with a cerebral abscess?**
 - A. Normal activity
 - B. Localized high voltage delta
 - C. Synchronized periodic spikes
 - D. Extensive theta activity

- 3. Which legislation emphasizes the minimum necessary use of protected health information?**
 - A. Health Insurance Accountability Act
 - B. Health Information Security Act
 - C. Health Insurance Portability and Accountability Act
 - D. Patient Privacy Protection Act

- 4. Which of the following is an effect of a large lesion near the optic chiasm?**
 - A. Double vision
 - B. Complete visual loss
 - C. Tunnel vision
 - D. Color vision deficiency

- 5. What combination of settings enhances a low voltage slow wave that is masked by muscle activity?**
 - A. Sensitivity of 3, Low Frequency 0.1, High Frequency 20
 - B. Sensitivity of 5, Low Frequency 0.3, High Frequency 35
 - C. Sensitivity of 7, Low Frequency 0.5, High Frequency 30
 - D. Sensitivity of 10, Low Frequency 0.1, High Frequency 25

- 6. What settings could help reduce sweat artifact in a patient with asymmetrical low voltage fast activity?**
- A. S=5 μ V/mm, LF=1 Hz, HF=30 Hz
 - B. S=5 μ V/mm, LF=5 Hz, HF=70 Hz
 - C. S=10 μ V/mm, LF=10 Hz, HF=100 Hz
 - D. S=2 μ V/mm, LF=2 Hz, HF=20 Hz
- 7. What causes a spike or sharp wave in an EEG?**
- A. Synchronous activation of multiple neurons
 - B. Defective neurotransmitter release
 - C. Increased metabolic activity in the brain
 - D. Disruption of cerebral circulation
- 8. Which type of tumor is least likely to cause epileptiform activity in EEG?**
- A. Meningioma
 - B. Pituitary adenoma
 - C. Glioblastoma
 - D. Astrocytoma
- 9. Which medication can cause triphasic waves and periodic discharges on an EEG?**
- A. Carbamazepine
 - B. Phenobarbital
 - C. Lithium
 - D. Valproate
- 10. What does the EEG of a very tense and anxious patient typically show?**
- A. High amplitude theta activity
 - B. Widespread low voltage activity
 - C. Periodic lateralized epileptiform discharges
 - D. Normal alpha rhythm

Answers

SAMPLE

1. B
2. B
3. C
4. B
5. B
6. B
7. A
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. Which cranial nerve primarily controls ocular movements?

- A. II
- B. III**
- C. IV
- D. VI

The cranial nerve that primarily controls ocular movements is cranial nerve III, known as the oculomotor nerve. This nerve is responsible for the majority of eye movements, including the movement of the eyelid and the constriction of the pupil. It innervates four of the six extraocular muscles, which allow for precise movements of the eyes in various directions. Cranial nerve II, or the optic nerve, is primarily responsible for vision but does not control any eye movements. Cranial nerve IV, known as the trochlear nerve, controls only the superior oblique muscle, which is responsible for downward and lateral eye movement. Cranial nerve VI, the abducens nerve, controls the lateral rectus muscle, which is involved in moving the eye laterally but does not assist with the comprehensive range of movements that cranial nerve III facilitates. Thus, the oculomotor nerve stands out as the primary nerve controlling ocular movements due to its extensive role in innervating multiple eye muscles.

2. What is the most likely EEG finding in a patient with a cerebral abscess?

- A. Normal activity
- B. Localized high voltage delta**
- C. Synchronized periodic spikes
- D. Extensive theta activity

In the context of a cerebral abscess, the most likely EEG finding is localized high voltage delta activity. This is indicative of the disruption of normal brain function in the area surrounding the abscess. The presence of high voltage delta waves suggests that there is localized impairment or irritation in brain activity, reflecting the abnormal physiological changes associated with the abscess. Cerebral abscesses can cause increased intracranial pressure and localized edema, which may lead to a slowing of cortical activity, evident in the form of delta waves. The high voltage characteristic indicates that the abnormality is significant and focused within a particular region of the brain. Normal activity, while it may appear in some cases, is less likely because a cerebral abscess typically alters the cortical dynamics in the affected area. Synchronized periodic spikes are more often associated with epileptogenic activity rather than a primary structural lesion like an abscess. Extensive theta activity may indicate diffuse cerebral dysfunction, which is less specific to the localized nature of a cerebral abscess and does not adequately represent the expected findings. Thus, localized high voltage delta activity is a more accurate representation of the EEG changes associated with this condition.

3. Which legislation emphasizes the minimum necessary use of protected health information?

- A. Health Insurance Accountability Act
- B. Health Information Security Act
- C. Health Insurance Portability and Accountability Act**
- D. Patient Privacy Protection Act

The emphasis on the minimum necessary use of protected health information is a key component of the Health Insurance Portability and Accountability Act (HIPAA). This legislation was enacted to enhance the privacy and security of individuals' health information. Under HIPAA, covered entities, such as healthcare providers and insurers, are required to take reasonable steps to limit the use and disclosure of protected health information to only what is necessary to achieve the intended purpose. This principle is crucial in minimizing the risk of unauthorized access or disclosure of sensitive health data, ensuring that individuals' privacy is respected while still allowing necessary information flow for treatment, payment, and healthcare operations. By mandating the minimum necessary standard, HIPAA balances the need for healthcare professionals to access patient information with the obligation to protect patient privacy. Other options do not capture this essential aspect of health information management and privacy that HIPAA specifically addresses. The Health Insurance Accountability Act and the Patient Privacy Protection Act do not exist as specific pieces of legislation recognized in the context of healthcare information privacy. The Health Information Security Act, while relevant to security standards, does not focus explicitly on the minimum necessary standard inherent to HIPAA.

4. Which of the following is an effect of a large lesion near the optic chiasm?

- A. Double vision
- B. Complete visual loss**
- C. Tunnel vision
- D. Color vision deficiency

A large lesion near the optic chiasm is particularly impactful due to the organization of visual pathways in the brain. The optic chiasm is the point where the optic nerves from each eye meet and partially cross over. Lesions in this area can disrupt the transmission of visual signals from both eyes to the brain. When such a lesion occurs, it can lead to complete visual loss depending on the extent and location of the damage, as it may affect both the nasal retina fibers from both eyes that carry crucial visual information from the peripheral visual field. This can result in a significant loss of vision, especially if the lesion is extensive enough to impact both pathways. In contrast, double vision typically arises from problems with the muscles that control eye movement or issues with the cranial nerves, which is not directly related to optic chiasm lesions. Tunnel vision would indicate a loss of peripheral vision, often caused by other conditions rather than an optic chiasm lesion. Color vision deficiency is usually linked to issues within the photoreceptors in the retina or the visual processing areas of the brain, rather than lesions affecting the optic chiasm specifically. Thus, the effect of a large lesion near the optic chiasm leading to complete visual loss is a direct outcome

5. What combination of settings enhances a low voltage slow wave that is masked by muscle activity?

- A. Sensitivity of 3, Low Frequency 0.1, High Frequency 20
- B. Sensitivity of 5, Low Frequency 0.3, High Frequency 35**
- C. Sensitivity of 7, Low Frequency 0.5, High Frequency 30
- D. Sensitivity of 10, Low Frequency 0.1, High Frequency 25

The correct answer enhances a low voltage slow wave that might be obscured by muscle activity through optimal settings of sensitivity and frequency ranges. A higher sensitivity setting, like 5 microvolts per millimeter, allows quieter or lower amplitude brain waves to be more easily detected and analyzed. This is critical when dealing with low-voltage slow waves because it increases their visibility against the background noise, such as muscle artifacts. The low frequency setting of 0.3 Hz is particularly effective in capturing slow waves, as it allows for the adequate filtering of faster frequencies that may introduce noise from muscle activity. This is essential for isolating the desired EEG signal without interference. On the other hand, the high-frequency setting of 35 Hz ensures that any muscle artifacts or high-frequency electrical noise can be effectively filtered out, focusing the analysis on the slower brain activity of interest. Thus, the combination of sensitivity, low frequency, and high frequency in this answer maximizes the ability to visualize low voltage slow waves while minimizing the impact of muscle activity, making it the most suitable choice for the stated requirement.

6. What settings could help reduce sweat artifact in a patient with asymmetrical low voltage fast activity?

- A. S=5 μ V/mm, LF=1 Hz, HF=30 Hz
- B. S=5 μ V/mm, LF=5 Hz, HF=70 Hz**
- C. S=10 μ V/mm, LF=10 Hz, HF=100 Hz
- D. S=2 μ V/mm, LF=2 Hz, HF=20 Hz

To reduce sweat artifacts in EEG recordings, it is essential to optimize the sensitivity and frequency settings of the EEG equipment. The correct choice emphasizes a balanced approach to the low-frequency (LF) and high-frequency (HF) settings. Selecting the setting with a low LF of 5 Hz allows the system to be more sensitive to the slower components of brain activity without being overly responsive to artifacts like sweat, which may have lower frequency components. The high frequency set at 70 Hz is also appropriate, as it helps eliminate higher-frequency noise, allowing for clearer brain signal recording. This combination of settings provides a compromise that emphasizes brain wave activity while filtering out rapid fluctuations caused by sweat or other muscle activity. The sensitivity setting of 5 μ V/mm ensures that the system captures sufficient detail without being overly sensitive to electrical noise. Other choices have either higher LF or HF settings that may not effectively filter out the rapid sweeps and spikes caused by sweat, or they have sensitivity settings which may either not capture the signal adequately or may amplify artifacts. Thus, option B effectively highlights the configuration that can mitigate sweat artifact while still capturing the necessary brain waves.

7. What causes a spike or sharp wave in an EEG?

A. Synchronous activation of multiple neurons

B. Defective neurotransmitter release

C. Increased metabolic activity in the brain

D. Disruption of cerebral circulation

The occurrence of a spike or sharp wave in an EEG is primarily due to the synchronous activation of multiple neurons. When a large group of neurons fires simultaneously, their electrical activity can generate an intense discharge that is manifest as a spike or sharp wave on the EEG tracing. This synchronized neuronal firing often reflects a pathological process, such as in epilepsy, where abnormal bursts of electrical activity lead to the characteristic waveforms observed on the EEG. In the context of the other options, while factors like neurotransmitter release and cerebral circulation can impact overall brain function and potentially lead to abnormal EEG patterns, they do not directly account for the specific formation of spikes or sharp waves. Increased metabolic activity in the brain can also influence EEG patterns; however, it is not specifically linked to the generation of spikes or sharp waves in the same way that synchronous neural activation is. Thus, understanding the mechanism of synchronous neuronal activation is key to recognizing the significance of spike and sharp wave patterns in EEG interpretations.

8. Which type of tumor is least likely to cause epileptiform activity in EEG?

A. Meningioma

B. Pituitary adenoma

C. Glioblastoma

D. Astrocytoma

Pituitary adenomas are typically considered the least likely to cause epileptiform activity in EEG recordings. The reason for this stems from their location and the nature of these tumors. Pituitary adenomas are located in the sella turcica at the base of the skull and primarily affect the pituitary gland, which is not directly associated with the cortex or the structures that generate cortical electrical activity. Because they often don't invade adjacent brain tissue or produce physical irritation or disruption to surrounding neuronal pathways, the likelihood of them causing seizures or epileptiform discharges on EEG is lower compared to other tumor types. In contrast, glioblastomas and astrocytomas are types of brain tumors that arise from glial cells and are more invasive, infiltrating nearby brain tissue. This invasive nature leads to cortical irritation, which can manifest as epileptiform activity on an EEG. Meningiomas, while often extra-axial and arising from the meninges, can also lead to seizures due to their location and potential to exert mass effects on the brain tissue, even if they do not directly invade it. Therefore, while various factors can influence seizure activity in relation to tumors, the pituitary adenoma stands out as the least likely to produce epilept

9. Which medication can cause triphasic waves and periodic discharges on an EEG?

- A. Carbamazepine
- B. Phenobarbital
- C. Lithium**
- D. Valproate

The presence of triphasic waves and periodic discharges on an EEG is most commonly associated with metabolic disturbances and certain medications that affect the central nervous system. Lithium is particularly noted for its potential to cause these specific EEG changes, especially in the context of lithium toxicity or when there are underlying metabolic issues, like renal impairment. Lithium affects neurotransmission and can lead to various neurological side effects. The triphasic waves seen on the EEG in patients taking lithium are indicative of a diffuse cerebral disturbance, potentially linked to the medication's effects on ion channels and excitatory neurotransmitters. Thus, the correlation between lithium administration and the appearance of triphasic waves and periodic discharges makes it a key medication to consider in this context. Other medications listed, while they can affect EEG readings in different ways, are not as directly associated with this specific pattern of abnormality. For instance, carbamazepine and phenobarbital are more commonly linked to generalized slowing, while valproate tends to cause specific types of spikes or slowing rather than the distinct patterns created by lithium. Therefore, understanding lithium's unique impact on the EEG is critical in recognizing its role in producing these findings.

10. What does the EEG of a very tense and anxious patient typically show?

- A. High amplitude theta activity
- B. Widespread low voltage activity**
- C. Periodic lateralized epileptiform discharges
- D. Normal alpha rhythm

The EEG of a very tense and anxious patient typically shows widespread low voltage activity. This pattern can be attributed to the heightened arousal and increased sympathetic nervous system activity that often accompany anxiety and tension. When a person is anxious, their brain can exhibit a state of heightened alertness, leading to increased frequency of low-amplitude waveforms, which are often associated with an active and alert mental state. In a tense state, the brain may not settle into the more organized rhythms such as alpha waves, which indicate relaxed alertness, or theta waves, which are typically seen in drowsy states or during certain cognitive tasks. Instead, the EEG may demonstrate a more chaotic pattern with low amplitude, representing diffuse cortical activation. This pattern reflects the underlying physiological state of the patient, as anxiety can disrupt normal brainwave patterns. It's important to note that conditions like periodic lateralized epileptiform discharges are indicative of specific types of seizure activity and typically would not be seen solely as a result of anxiety. Similarly, normal alpha rhythm would indicate a relaxed but alert state, which does not align with the presentation of a tense and anxious individual. High amplitude theta activity, while present in some altered states of consciousness, is not characteristic of anxiety in a straightforward context.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://abreteeg.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE