
ACAAI BOARD PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://acaaiboard.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which CD markers are alpha/beta double-negative T cells (DNTs) lacking?

A. CD4 and CD8

B. CD3 and CD19

C. CD25 and CD56

D. CD14 and CD16

2. What is a potential indicator of severe allergic reaction post insect sting that requires blood testing?

A. Elevated blood sugar

B. High tryptase level

C. Low hemoglobin level

D. Increased platelet count

3. What is the last sinus to develop in children?

A. Ethmoid sinus

B. Sphenoid sinus

C. Frontal sinus

D. Maxillary sinus

4. Which type of T cells express CD4?

A. Naïve T cells

B. Helper T cells

C. Cytotoxic T cells

D. Regulatory T cells

5. Which HLA types are critical for histocompatibility matching in transplants?

A. HLA A, B, C, DRB1, and DQB1

B. HLA DR, DQ, A2, B5, and C3

C. HLA A1, B8, DR7, and DQB2

D. HLA B, DR, and C only

SAMPLE

6. What are considered the most significant indoor molds?

A. Aspergillus, Alternaria, and Penicillium

B. Aspergillus, Cladosporium, and Penicillium

C. Cladosporium, Penicillium, and Fusarium

D. Penicillium, Trichoderma, and Alternaria

7. What is the most common cause of rhinitis during pregnancy?

A. Newly developed allergies

B. Preexisting allergic rhinitis

C. Physiological changes related to pregnancy

D. Respiratory infections during pregnancy

8. Which conditions can potentially cause a false positive AHR?

A. Asthma and diabetes

B. Allergic rhinitis and smoking

C. Hypertension and obesity

D. Chronic renal failure and hyperthyroidism

9. A woman with progressive hearing loss and recurrent episodes of fever and urticaria may be diagnosed with:

A. Muckle-Wells syndrome

B. Still's disease

C. Behçet's disease

D. Systemic lupus erythematosus

10. What are the two primary coreceptors utilized by HIV?

A. CD4 and CXCR4

B. CCR5 and CXCR4

C. CXCR5 and CD26

D. CCR1 and CCR3

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. A
6. B
7. B
8. B
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. Which CD markers are alpha/beta double-negative T cells (DNTs) lacking?

A. CD4 and CD8

B. CD3 and CD19

C. CD25 and CD56

D. CD14 and CD16

Alpha/beta double-negative T cells (DNTs) are a distinct subset of T cells characterized by their lack of both CD4 and CD8 markers. These markers are typically associated with conventional T cells, where CD4+ T cells are often helper T cells and CD8+ T cells are cytotoxic T cells. The absence of CD4 and CD8 helps to define the DNTs, indicating that they do not fit the standard profiles of helper or cytotoxic T cells. DNTs can still express other T cell-associated markers, which helps them perform specific immune functions despite not having these core surface markers. Hence, the identification of DNTs relies strongly on their expression of surface proteins, where the loss of CD4 and CD8 is the defining feature of this T cell population. The other options involve markers that do not directly define the double-negative phenotype in T cells, as they are specific cell surface proteins associated with different cell types or other aspects of T cell functionality. For example, CD3 is a T cell receptor complex component, CD19 is a B cell marker, and CD25 and CD56 are associated with different immune cell functionalities, further demonstrating that the most relevant defining markers in this context are indeed CD4

2. What is a potential indicator of severe allergic reaction post insect sting that requires blood testing?

A. Elevated blood sugar

B. High tryptase level

C. Low hemoglobin level

D. Increased platelet count

A high tryptase level is a significant indicator of a severe allergic reaction, especially in the context of an insect sting. Tryptase is an enzyme released from mast cells during an anaphylactic reaction. When an individual experiences such a severe allergic response, the levels of tryptase in the blood can rise markedly within a few hours, reflecting the degranulation of mast cells and the immune response to the allergen. In cases of anaphylaxis, measuring the tryptase level can help confirm the diagnosis, as it is often elevated in response to significant allergic reactions. Blood tests for tryptase are typically performed within a few hours after the event to assess the severity of the reaction. The other options do not specifically correlate with severe allergic reactions post-insect sting. Elevated blood sugar may indicate stress or an underlying metabolic condition but is not specific to an allergic reaction. Low hemoglobin levels are more suggestive of anemia rather than an immediate allergic response, while an increased platelet count can be related to various conditions but does not signify an anaphylactic reaction. Thus, elevated tryptase levels are the primary indicator guiding clinicians toward the diagnosis of a severe allergic reaction.

SAMPLE

3. What is the last sinus to develop in children?

A. Ethmoid sinus

B. Sphenoid sinus

C. Frontal sinus

D. Maxillary sinus

The frontal sinus is indeed the last sinus to develop in children. It typically begins to develop around the age of 2 years but does not reach full adult size until adolescence, usually around the ages of 12 to 14. This delayed development can be attributed to the growth patterns of the skull, where the frontal sinus expands as the face matures and the bones around the sinus grow. In contrast, the ethmoid and maxillary sinuses are present at birth and grow significantly during the early years of life. The sphenoid sinus usually develops between the ages of 2 to 3 years, which is earlier than the frontal sinus. Understanding the sequence and timing of sinus development is important in recognizing and diagnosing sinus-related issues in pediatric patients.

4. Which type of T cells express CD4?

A. Naïve T cells

B. Helper T cells

C. Cytotoxic T cells

D. Regulatory T cells

Helper T cells are a specific subset of T cells that play a critical role in the immune response. They are identified by the expression of the CD4 glycoprotein on their surface. This differentiation is key because CD4+ T cells assist other immune cells, such as B cells and cytotoxic T cells, by releasing cytokines that enhance the immune response. In contrast, while naïve T cells are precursor cells that can develop into different types of T cells upon activation, they do not specifically express CD4 until they mature into helper T cells. Cytotoxic T cells, which are responsible for directly killing infected or malignant cells, express CD8 instead of CD4. Regulatory T cells, which help maintain immune tolerance and prevent excessive immune responses, can also express CD4 but are more specifically characterized by the presence of the transcription factor FoxP3 rather than their CD4 expression alone. The distinction of helper T cells as CD4+ is essential in understanding their function in orchestrating the immune response, making this option the correct choice.

5. Which HLA types are critical for histocompatibility matching in transplants?

A. HLA A, B, C, DRB1, and DQB1

B. HLA DR, DQ, A2, B5, and C3

C. HLA A1, B8, DR7, and DQB2

D. HLA B, DR, and C only

The correct answer identifies a set of human leukocyte antigen (HLA) types that are essential for histocompatibility matching in transplantation. HLA molecules are critical in the immune response and play a significant role in determining compatibility between donors and recipients. The HLA A, B, C, DRB1, and DQB1 types mentioned in the correct answer encompass both class I and class II HLA molecules. Class I molecules (HLA A, B, and C) are present on nearly all nucleated cells and are involved in presenting endogenous antigens to CD8+ cytotoxic T cells. Class II molecules (HLA DRB1 and DQB1) are primarily found on professional antigen-presenting cells and interact with CD4+ T helper cells. In transplantation, the matching of these HLA types is crucial because mismatches can lead to acute and chronic rejection of transplanted organs. A robust match in these five specific HLA types increases the likelihood of transplant acceptance and improves patient outcomes. The other options do not accurately represent the critical markers needed for effective histocompatibility matching. For instance, options that list inappropriate or incomplete sets of HLA types may overlook important specificities necessary for successful transplantation. Recognizing the

SAMPLE

6. What are considered the most significant indoor molds?

A. Aspergillus, Alternaria, and Penicillium

B. Aspergillus, Cladosporium, and Penicillium

C. Cladosporium, Penicillium, and Fusarium

D. Penicillium, Trichoderma, and Alternaria

Aspergillus, Cladosporium, and Penicillium are indeed considered some of the most significant indoor molds due to their prevalence and potential health impacts. Aspergillus is particularly notable because it comprises several species, some of which can produce mycotoxins that may lead to various respiratory issues and allergic reactions. Cladosporium is another common mold found in homes that can trigger asthma and other allergic responses, while Penicillium is well-known for its role in both allergies and the production of mycotoxins. The presence of these molds in indoor environments raises concerns regarding indoor air quality and associated health risks, making them critical for consideration in both clinical and environmental contexts. This understanding reflects the importance of identifying and managing indoor molds effectively to mitigate health impacts.

7. What is the most common cause of rhinitis during pregnancy?

A. Newly developed allergies

B. Preexisting allergic rhinitis

C. Physiological changes related to pregnancy

D. Respiratory infections during pregnancy

The most common cause of rhinitis during pregnancy is often attributed to preexisting allergic rhinitis. Many women who already have allergies may notice an exacerbation of their symptoms during pregnancy due to hormonal changes and increased vascular permeability, which can enhance nasal congestion and other allergic responses. The physiological changes associated with pregnancy, including elevated estrogen and progesterone levels, can lead to increased mucosal edema and sensitivity, which in turn can amplify allergic reactions. While it is true that physiological changes related to pregnancy also contribute significantly to nasal symptoms, they primarily affect those with preexisting allergic conditions rather than being the most common cause in isolation. Newly developed allergies can occur, but they are less common compared to the exacerbation of existing allergic rhinitis. Respiratory infections may lead to nasal symptoms as well but typically do not represent a chronic issue compared to the reiterated challenges faced by those with preexisting allergic conditions. Hence, preexisting allergic rhinitis stands out as the predominant cause of rhinitis in pregnant individuals.

SAMPLE

8. Which conditions can potentially cause a false positive AHR?

A. Asthma and diabetes

B. Allergic rhinitis and smoking

C. Hypertension and obesity

D. Chronic renal failure and hyperthyroidism

A false positive in airway hyperresponsiveness (AHR) testing can occur due to various factors, and allergic rhinitis is particularly relevant in this context. Individuals with allergic rhinitis can experience increased airway inflammation and bronchial hyperreactivity, which can lead to a false positive result when testing for AHR. This condition enhances the sensitivity of the airways to various stimuli, potentially skewing test results. Smoking also plays a significant role in the respiratory response and may influence AHR results by contributing to airway inflammation, increased bronchial responsiveness, and altered airflow dynamics. Smokers may display airway changes that mimic AHR, leading to difficulty in interpreting results accurately. Asthma is a condition typically associated with true positive AHR responses rather than false positives, as it inherently involves bronchial hyperreactivity. Diabetes, hypertension, and obesity do not directly influence airway reactivity in the same manner as allergic rhinitis and smoking. Chronic renal failure and hyperthyroidism also do not have a well-established connection to airway hyperresponsiveness. Therefore, the combination of allergic rhinitis and smoking is particularly significant in understanding potential false positives in AHR testing.

9. A woman with progressive hearing loss and recurrent episodes of fever and urticaria may be diagnosed with:

A. Muckle-Wells syndrome

B. Still's disease

C. Behçet's disease

D. Systemic lupus erythematosus

The correct answer is Muckle-Wells syndrome, which is a genetic autoinflammatory condition associated with sensorineural hearing loss, intermittent fever, and skin manifestations such as urticaria. The episodes of urticaria and fever are characteristic of the periodic inflammatory attacks that occur in this syndrome. Additionally, the progressive hearing loss observed in the patient aligns with the known auditory complications associated with Muckle-Wells syndrome. Muckle-Wells syndrome is caused by mutations in the NLRP3 gene, which plays a crucial role in the inflammatory response. The combination of symptoms in this context makes Muckle-Wells the most fitting diagnosis since it specifically encompasses hearing loss and recurrent fever with urticarial-type rash. Other conditions listed may feature some overlapping symptoms but do not present with this specific combination to the same extent or context. For instance, Still's disease primarily involves arthritis, fever, and rash but is not characterized by progressive hearing loss. Behçet's disease involves systemic vasculitis and can cause oral and genital ulcers and eye inflammation, but it does not usually lead to hearing impairment. Systemic lupus erythematosus also presents with a wide variety of systemic symptoms, including skin rashes and fever, but tends to have different

SAMPLE

10. What are the two primary coreceptors utilized by HIV?

A. CD4 and CXCR4

B. CCR5 and CXCR4

C. CXCR5 and CD26

D. CCR1 and CCR3

The two primary coreceptors utilized by HIV for entry into host cells are CCR5 and CXCR4. HIV primarily targets CD4 positive T cells, which serve as the main cell type for viral entry. However, the entry process also requires these coreceptors. CCR5 is important for the initial stages of infection, particularly in the early phases of HIV transmission, as many strains of HIV use this coreceptor to infect mucosal tissues. Later in the disease, the virus may switch to using CXCR4, which is often associated with widespread disease progression and is found on activated T cells. The ability of HIV to utilize both coreceptors—understanding that CCR5 is primarily used in the acute phase and CXCR4 in the chronic phase—illustrates a crucial aspect of HIV pathogenesis and treatment considerations. In this context, the other options do not reflect the coreceptor interactions relevant to HIV. CD4 is indeed required for the virus to attach to the host cell, but it is not classified as a coreceptor. Likewise, CXCR5 and CD26 are not utilized by HIV for entry, nor are CCR1 and CCR3, which are involved in other immune responses but do not serve the function required for HIV entry into cells. This

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://acaaiboard.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE