
ACVIM GENERAL BOARDS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://acvimgenboards.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the primary physiological process affected by myasthenia gravis?

A. Muscle relaxation

B. Acetylcholine receptor function

C. Nerve impulse transmission

D. Calcium reabsorption

2. Which diuretic class does spironolactone belong to?

A. Loop diuretics

B. Thiazide diuretics

C. Potasium-sparing diuretics

D. Osmotic diuretics

3. How is plasma osmolality calculated?

A. Posm = 2Na + glucose/18 + BUN/2.8

B. Posm = Na + K + Cl

C. Posm = 2Na + BUN

D. Posm = glucose/18 + 2BUN

4. What is the primary mechanism of action of aspirin?

A. Inhibiting reabsorption of sodium and chloride

B. Loop diuretic effect

C. Irreversibly blocking cyclooxygenase

D. Carbonic anhydrase inhibition

5. What is the effect of mannitol in the lumen of the renal tubules?

A. Promotes sodium reabsorption

B. Inhibits reabsorption of bicarbonate

C. Retains water, promoting diuresis

D. Blocks sodium channels

6. Which hormones are released from the anterior pituitary?

A. Cortisol and adrenaline

B. Prolactin, GH, ACTH, TSH, LH, FSH

C. Insulin and glucagon

D. Oxytocin and ADH

SAMPLE

7. How do CD8 cells execute their cytotoxic function?

A. By producing antibodies

B. Through perforin, granzyme, granulysin, and Fas ligand

C. Via phagocytosis

D. By releasing cytokines

8. Which heartworm testing method is more sensitive in cats compared to dogs?

A. Antigen testing

B. Antibody testing

C. Direct blood smear

D. Radiographic assessment

9. What chemotherapy drug is commonly used to treat polycythemia vera?

A. Cyclophosphamide

B. Hydroxyurea

C. Vincristine

D. Ruxolitinib

10. Which type of V/Q mismatch would not benefit from oxygen supplementation?

A. Physiologic dead space

B. Physiologic shunt

C. Intrapulmonary shunting

D. Airway obstruction

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. C
5. C
6. B
7. B
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is the primary physiological process affected by myasthenia gravis?

A. Muscle relaxation

B. Acetylcholine receptor function

C. Nerve impulse transmission

D. Calcium reabsorption

The primary physiological process affected by myasthenia gravis is acetylcholine receptor function. Myasthenia gravis is an autoimmune neuromuscular disorder characterized by weakness and rapid fatigue of voluntary muscles. In this condition, antibodies are produced that target and block or destroy nicotinic acetylcholine receptors at the neuromuscular junction. This impedes the ability of acetylcholine, a neurotransmitter, to bind to its receptors effectively, thereby disrupting the transmission of nerve signals that stimulate muscle contraction. In essence, acetylcholine normally binds to these receptors to facilitate muscle contraction, and when their function is compromised, it leads to the hallmark symptoms of muscle weakness and fatigue. Thus, understanding the involvement of acetylcholine receptor function provides crucial insights into the pathophysiology of myasthenia gravis and its clinical manifestations.

2. Which diuretic class does spironolactone belong to?

A. Loop diuretics

B. Thiazide diuretics

C. Potasium-sparing diuretics

D. Osmotic diuretics

Spironolactone is classified as a potassium-sparing diuretic. This class of diuretics works by antagonizing the effects of aldosterone, a hormone that promotes sodium retention and potassium excretion in the kidneys. By blocking aldosterone, spironolactone allows for the retention of potassium while promoting the excretion of sodium and water, thus helping to reduce blood volume and lower blood pressure. Potassium-sparing diuretics are particularly useful in situations where avoiding potassium loss is critical, such as in patients at risk for hypokalemia or those on concurrent medications that can further deplete potassium levels. These diuretics are commonly used in the management of conditions like heart failure, hypertension, and conditions characterized by excessive aldosterone production, such as primary hyperaldosteronism. In contrast, loop diuretics and thiazide diuretics primarily promote the excretion of both sodium and potassium, while osmotic diuretics mainly increase osmolarity in the renal tubules to draw water out of the body. Using spironolactone appropriately can enhance the diuretic effect when used in combination with other classes while minimizing the risk of hypokalemia.

SAMPLE

3. How is plasma osmolality calculated?

A. Posm = 2Na + glucose/18 + BUN/2.8

B. Posm = Na + K + Cl

C. Posm = 2Na + BUN

D. Posm = glucose/18 + 2BUN

The calculation of plasma osmolality is based on the concentration of solutes in the plasma that contribute significantly to its osmotic pressure. The formula for calculating plasma osmolality includes the major contributors to plasma osmolality, which are sodium, glucose, and urea. The correct formula, which is used clinically to estimate plasma osmolality, is expressed as: Posm = 2(Na) + (glucose/18) + (BUN/2.8) In this equation: - The term 2(Na) accounts for the effect of sodium ions, which are predominant in determining osmotic balance. - Glucose is included and divided by 18 because glucose contributes to osmolality, and the division converts the concentration from mg/dL to mmol/L. - BUN (blood urea nitrogen) is also included, divided by 2.8 to adjust for its contribution to osmotic pressure since this conversion similarly transforms the measurement into mmoles/L. This comprehensive formula accurately reflects the primary solutes contributing to plasma osmolality, providing a reliable estimation that can be critical in various clinical settings, especially when assessing fluid and electrolyte balance or evaluating conditions such as hypernatremia or hyponatremia

4. What is the primary mechanism of action of aspirin?

A. Inhibiting reabsorption of sodium and chloride

B. Loop diuretic effect

C. Irreversibly blocking cyclooxygenase

D. Carbonic anhydrase inhibition

The primary mechanism of action of aspirin is the irreversible blocking of cyclooxygenase (COX). Cyclooxygenase is an enzyme critical in the conversion of arachidonic acid to prostaglandins, which are compounds that play various roles in inflammation, pain signaling, and platelet aggregation. By inhibiting COX, aspirin effectively decreases the production of these prostaglandins, which in turn reduces inflammation and pain. This action also leads to antiplatelet effects, making aspirin useful in preventing thromboembolic events such as heart attacks and strokes. The irreversible nature of aspirin's inhibition means that its effects on the COX enzyme persist until new COX enzymes are synthesized by the body, which can take a period of time, particularly in platelets that lack nuclei and cannot regenerate the enzyme quickly. This unique property underpins many of the therapeutic applications of aspirin, distinguishing it from other anti-inflammatory and analgesic medications that may work through reversible mechanisms. Understanding this mechanism is crucial for leveraging aspirin's benefits in clinical practice, emphasizing its role not only in pain relief but also in cardiovascular protection.

SAMPLE

5. What is the effect of mannitol in the lumen of the renal tubules?

A. Promotes sodium reabsorption

B. Inhibits reabsorption of bicarbonate

C. Retains water, promoting diuresis

D. Blocks sodium channels

Mannitol is an osmotic diuretic that works primarily in the renal tubules, particularly in the proximal convoluted tubule and the loop of Henle. Its mechanism is based on its ability to remain in the tubular lumen and create an osmotic pressure that prevents the reabsorption of water. When mannitol is administered, it is filtered by the kidneys and remains in the renal tubules, where it draws water into the tubular fluid due to the osmotic gradient it establishes. As a result, the presence of mannitol in the tubular lumen leads to increased excretion of water, thus promoting diuresis. This is beneficial in conditions requiring the reduction of fluid overload, such as in cases of acute renal failure or cerebral edema, where rapid diuresis can alleviate pressure and improve renal function. The other options suggest different mechanisms that do not apply to mannitol's action in the renal tubules. For example, mannitol does not promote sodium reabsorption or inhibit bicarbonate reabsorption. It also does not block sodium channels but rather acts osmotically to increase the volume of urine by preventing water reabsorption. Consequently, option C accurately captures the pivotal role of mannitol in promoting

6. Which hormones are released from the anterior pituitary?

A. Cortisol and adrenaline

B. Prolactin, GH, ACTH, TSH, LH, FSH

C. Insulin and glucagon

D. Oxytocin and ADH

The anterior pituitary, also known as the adenohypophysis, plays a crucial role in regulating various physiological processes through the secretion of several key hormones. The hormones released from the anterior pituitary include prolactin, growth hormone (GH), adrenocorticotropic hormone (ACTH), thyroid-stimulating hormone (TSH), luteinizing hormone (LH), and follicle-stimulating hormone (FSH). Prolactin is primarily involved in milk production and lactation, GH plays a vital role in growth and metabolism, ACTH stimulates the adrenal cortex to produce cortisol, TSH regulates thyroid hormone production, LH and FSH are critical for reproductive functions such as ovulation and spermatogenesis. In contrast, cortisol and adrenaline are produced by the adrenal glands, insulin and glucagon are hormones produced by the pancreas, and oxytocin and antidiuretic hormone (ADH) are synthesized in the hypothalamus and released from the posterior pituitary, not the anterior pituitary. Thus, the hormones listed in the correct answer represent the primary functions and outputs of the anterior pituitary gland.

SAMPLE

7. How do CD8 cells execute their cytotoxic function?

A. By producing antibodies

B. Through perforin, granzyme, granulysin, and Fas ligand

C. Via phagocytosis

D. By releasing cytokines

CD8 T cells, also known as cytotoxic T lymphocytes, are specifically designed to eliminate infected or cancerous cells. Their cytotoxic function is primarily executed through several key effector molecules, including perforin, granzyme, granulysin, and Fas ligand. When CD8 T cells recognize a target cell that presents the appropriate antigen via MHC class I molecules, they engage in direct cytotoxicity. Perforin is released by the CD8 T cells and forms pores in the target cell membrane. This pore formation allows granzyme, a serine protease, to enter the target cell, where it initiates apoptosis (programmed cell death) by activating intracellular apoptotic pathways. Granulysin also has cytotoxic properties; it can enter the target cell and induce apoptosis as well, while also having antimicrobial effects against some pathogens. The Fas ligand interaction with the Fas receptor on the target cell is another crucial pathway that signals apoptosis, allowing for targeted destruction of cells that are infected or dysfunctional. In summary, the cytotoxic activity of CD8 T cells is a sophisticated and coordinated mechanism involving the release of specific cytolytic proteins and factors that induce cell death, ensuring the removal of harmful cells from the organism.

8. Which heartworm testing method is more sensitive in cats compared to dogs?

A. Antigen testing

B. Antibody testing

C. Direct blood smear

D. Radiographic assessment

Antibody testing is particularly significant in the context of diagnosing heartworm disease in cats compared to dogs due to differences in the host's immune response to the parasite. In cats, the immune response to infection with Dirofilaria immitis often leads to the production of antibodies, which can be detected through this testing method even when the actual adult heartworms are present in smaller numbers or when they are immature. Cats tend to have a lower worm burden when infected, and antigen tests, which rely on detecting specific proteins produced by adult female heartworms, may not be as effective given the fewer adult females present. Furthermore, many infected cats can have a transient or incomplete infection wherein they do not harbor enough adult female worms to trigger a positive antigen test, making antibody testing a more sensitive method in this species. The direct blood smear method and radiographic assessments are generally less reliable for detecting heartworm infection as they do not provide the same level of specificity or sensitivity regarding the immune response. Thus, antibody testing stands out as the preferred method in cats, enhancing the likelihood of early and accurate diagnosis of heartworm disease.

SAMPLE

9. What chemotherapy drug is commonly used to treat polycythemia vera?

A. Cyclophosphamide

B. Hydroxyurea

C. Vincristine

D. Ruxolitinib

Hydroxyurea is commonly used in the treatment of polycythemia vera because it helps to reduce the high number of red blood cells by inhibiting the production of DNA in the bone marrow. This ultimately reduces the overall cell proliferation, which is a key characteristic of polycythemia vera. By lowering the red blood cell count, hydroxyurea helps to alleviate symptoms and decrease the risk of thrombotic events associated with the condition. While other medications like Ruxolitinib may also play a role in managing polycythemia vera—particularly in cases that are resistant to conventional therapies—hydroxyurea remains a standard and widely used treatment primarily due to its effectiveness in controlling the hyperproliferative aspects of the disease. Cyclophosphamide and Vincristine are not typically utilized for polycythemia vera, making hydroxyurea the most established choice in this context.

10. Which type of V/Q mismatch would not benefit from oxygen supplementation?

A. Physiologic dead space

B. Physiologic shunt

C. Intrapulmonary shunting

D. Airway obstruction

Intrapulmonary shunting refers to a situation where blood passes from the right side of the heart to the left side without becoming oxygenated in the lungs due to conditions such as pneumonia, atelectasis, or lung consolidation. In this scenario, blood is effectively "shunted" past the areas of the lung that are functioning normally for gas exchange, meaning that the blood does not pick up adequate oxygen. When oxygen is administered to a patient with intrapulmonary shunting, the oxygen does not reach the areas of the lung where it can be effectively exchanged into the bloodstream because these areas are poorly ventilated or not ventilated at all. As a result, it does not improve the oxygenation of the blood, leading to hypoxemia. This is a critical distinction because supplemental oxygen is typically effective in cases where there is impaired ventilation in normally perfused areas (like in some types of V/Q mismatch). However, in the case of intrapulmonary shunting, despite the presence of supplemental oxygen, the underlying problem of inadequate ventilation persists, so it does not resolve the mismatch and can be considered ineffective. In summary, intrapulmonary shunting does not benefit from oxygen supplementation because the condition involves blood flow bypassing ventilated

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://acvimgenboards.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE