ADEX DIAGNOSTIC SKILLS EXAMINATION (DSE) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adexdse.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the appropriate action for a radiolucency next to the PM?

A. Observe for changes

B. Refer to general dentistry

C. Refer to oral surgery

D. Perform a biopsy

2. When replacing an anterior crown, how much incisive reduction is typically required?

A. 1 mm

B. 2 mm

C. 3 mm

D. 4 mm

3. What aesthetic issue can mesiodens lead to?

A. Increased tooth whiteness

B. Poor aesthetics

C. Enhanced smile symmetry

D. Improved facial structure

4. What is the characteristic feature of dentinogenesis imperfecta?

A. Thin enamel

B. Discolored dentin

C. Missing teeth

D. Gingival inflammation

5. What action is recommended if a tooth still has radiolucency two years post-root canal treatment?

A. Continue monitoring

B. Re-treat the tooth

C. Refer to a specialist

D. Perform a surgical extraction

SAMPLE

6. Where is it advisable to store scrap amalgam?

A. In a dry container

B. Under sulfide

C. In a glass jar

D. In an open bin

7. How is a Candida infection typically treated?

A. Clindamycin

B. Fluconazole

C. Nystatin

D. Amphotericin B

8. What causes mydriasis in patients?

A. Anti-cholinergics

B. Beta-blockers

C. Calcium channel blockers

D. SSRIs

9. What is the likely diagnosis for a purplish lesion of the lip?

A. Fibroma

B. Hemangioma

C. Squamous cell carcinoma

D. Psoriasis

10. What dental condition can result from inflammation of the tissue surrounding a tooth?

A. Gingivitis

B. Periodontitis

C. Abscess

D. Canker sore

SAMPLE

Answers

SAMPLE

1. C
2. B
3. B
4. B
5. B
6. B
7. B
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is the appropriate action for a radiolucency next to the PM?

A. Observe for changes

B. Refer to general dentistry

C. Refer to oral surgery

D. Perform a biopsy

In the context of a radiolucency located adjacent to a premolar, referring the patient to oral surgery is often the most appropriate action due to several reasons. Radiolucencies can indicate various conditions such as dental cysts, tumors, or periodontal disease, which may require surgical intervention for diagnosis and treatment. Oral surgeons are specially trained to handle these types of lesions, especially if there is a possibility of malignancy or if the lesion needs to be excised for further evaluation. Referring to general dentistry may not address the complexity of the situation, as general dentists typically manage more routine cases and may not be equipped to deal with the intricate assessment or surgical requirements that a radiolucency could entail. Observing for changes could lead to a delay in treatment, which is particularly concerning if the radiolucency is associated with a serious condition that warrants prompt attention. Performing a biopsy is indeed a necessary action for definitive diagnosis in certain situations, but it is usually carried out by specialists who have the necessary expertise and facilities, thus emphasizing the importance of the referral to oral surgery. In summary, the referral to oral surgery is the most appropriate response for addressing findings of a radiolucency next to a premolar, prioritizing accurate diagnosis and

2. When replacing an anterior crown, how much incisive reduction is typically required?

A. 1 mm

B. 2 mm

C. 3 mm

D. 4 mm

When replacing an anterior crown, typically 2 mm of incisive reduction is required. This amount of reduction is important to ensure there is adequate space for the materials used in the crown, which may vary in thickness depending on the type of crown being fabricated (e.g., porcelain, metal-ceramic). Achieving the right amount of reduction helps to avoid issues with crown retention and aesthetics, and ensures that the occlusion is correctly aligned. If less than 2 mm is removed, the crown may not have sufficient material to create a durable restoration or it may appear overly bulky, compromising the natural appearance. Conversely, excessive reduction could weaken the tooth structure and lead to sensitivity or the need for further restorative procedures. Thus, the 2 mm reduction strikes a balance that promotes long-term success in anterior crown restorations.

3. What aesthetic issue can mesiodens lead to?

A. Increased tooth whiteness

B. Poor aesthetics

C. Enhanced smile symmetry

D. Improved facial structure

The presence of mesiodens, which are extra teeth that typically appear in the midline of the dental arch, can result in poor aesthetics. These supernumerary teeth can lead to crowding or misalignment of adjacent teeth, which disrupts the symmetry and overall appearance of an individual's smile. When mesiodens are present, they can interfere with the proper eruption of normal teeth and may create space issues that affect how teeth align and position themselves in the mouth. This misalignment can detract from the aesthetic harmony that is often desired in a smile, making it less visually pleasing. In contrast, increased tooth whiteness, enhanced smile symmetry, and improved facial structure do not typically relate to the presence of a mesiodens. Instead, mesiodens are more commonly associated with complications that negatively impact dental aesthetics.

SAMPLE

4. What is the characteristic feature of dentinogenesis imperfecta?

A. Thin enamel

B. Discolored dentin

C. Missing teeth

D. Gingival inflammation

Dentinogenesis imperfecta is an inherited disorder that primarily affects the structure and appearance of dentin, a vital component of teeth. The hallmark of this condition is discolored dentin, which can give the teeth a yellow, brown, or even opalescent appearance. This discoloration occurs due to the defective formation of collagen, which impacts the mineralization process and the overall quality of dentin. While other features such as thin enamel or a potential for missing teeth can be observed in various dental conditions, they are not the defining characteristics of dentinogenesis imperfecta. Principal complications may include issues related to the strength and support provided by dentin, leading to further susceptibility to dental wear or fractures. However, the key feature that distinguishes dentinogenesis imperfecta is indeed the discoloration of dentin itself, affirming its primary role in the diagnosis and understanding of this dental anomaly.

5. What action is recommended if a tooth still has radiolucency two years post-root canal treatment?

A. Continue monitoring

B. Re-treat the tooth

C. Refer to a specialist

D. Perform a surgical extraction

If a tooth still shows radiolucency two years after root canal treatment, it often indicates the presence of persistent infection or failure of the initial treatment. In this context, re-treatment is typically recommended, as it allows for a thorough evaluation and intervention to address any lingering infection that may not have been completely resolved during the original procedure. Re-treating the tooth involves cleaning out the root canal system again, which may include the use of different techniques or materials to eliminate the infection and promote healing. This approach prioritizes preserving the tooth and alleviating any ongoing symptoms or risks associated with infection. Monitoring the tooth over time might be an initial approach, but with the evidence of ongoing radiolucency, it suggests that more active treatment is necessary. Referral to a specialist can be useful in some cases, particularly for complex scenarios, but it may not be the first line of action if re-treatment is viable and the dentist is equipped to handle it. Performing a surgical extraction should be considered only if all other options fail or if the tooth is deemed non-salvageable, which is not the case here.

6. Where is it advisable to store scrap amalgam?

A. In a dry container

B. Under sulfide

C. In a glass jar

D. In an open bin

Storing scrap amalgam under sulfide is advisable due to its ability to mitigate the potential release of toxic metals, primarily mercury, which can be present in amalgam waste. By storing it under sulfide, the mercury is chemically bound and prevented from escaping into the environment, which is critical for maintaining safety in dental practice and compliance with environmental regulations. This method of storage is also generally recognized to minimize health risks associated with exposure to mercury vapors. In contrast, storing scrap amalgam in a dry container or a glass jar might not adequately contain the mercury, allowing for possible leakage or evaporation into the atmosphere. An open bin would pose the highest risk, as it would leave the amalgam exposed to air and potential contamination or accidental exposure, which is not a safe practice in any clinical setting.

SAMPLE

7. How is a Candida infection typically treated?

A. Clindamycin

B. Fluconazole

C. Nystatin

D. Amphotericin B

The treatment of a Candida infection often involves the use of antifungal medications that specifically target the Candida species. Fluconazole is a commonly prescribed oral antifungal that belongs to the azole class, and it is effective against a range of Candida infections, including candidiasis of the mouth, throat, and esophagus. It works by inhibiting the synthesis of ergosterol, an essential component of fungal cell membranes, which leads to the disruption of cellular function and ultimately results in the death of the fungi. Fluconazole is particularly favored due to its efficacy, oral availability, and relatively low side-effect profile compared to other antifungals. This makes it a first-line treatment option for localized mucosal infections as well as some systemic forms of candidiasis. While other medications like Nystatin, Amphotericin B, and Clindamycin may be used in specific cases or for different infections, Fluconazole is generally the go-to choice for routine Candida infections, especially when a systemic treatment is indicated. For example, Nystatin is primarily used for topical or oral fungal infections and is not as commonly used systemically. Amphotericin B is typically reserved for serious systemic infections due to its toxicity, and Clindamycin is an

8. What causes mydriasis in patients?

A. Anti-cholinergics

B. Beta-blockers

C. Calcium channel blockers

D. SSRIs

Mydriasis, or dilation of the pupils, occurs when there is an increase in sympathetic nervous system activity or a decrease in parasympathetic activity affecting the iris sphincter muscle. Anti-cholinergics, specifically, block the action of acetylcholine on the muscarinic receptors in the eye. This inhibition reduces the tone of the sphincter muscle, leading to pupil dilation. The other options do not have a primary mechanism that leads to mydriasis. Beta-blockers primarily affect the cardiovascular system by blocking adrenergic receptors, and while they may have some effect on ocular pressure, they do not directly cause pupil dilation. Calcium channel blockers affect vascular smooth muscle and cardiac tissue but do not have a direct effect on pupil size. SSRIs primarily target serotonin reuptake in the brain and are associated mainly with mood regulation and do not cause pupil dilation as a primary effect. Thus, the action of anti-cholinergics is distinctly responsible for causing mydriasis.

SAMPLE

9. What is the likely diagnosis for a purplish lesion of the lip?

A. Fibroma

B. Hemangioma

C. Squamous cell carcinoma

D. Psoriasis

A purplish lesion on the lip is most indicative of a hemangioma. Hemangiomas are benign tumors made up of blood vessels and are characterized by their reddish to purplish color due to the presence of these vessels. They can appear anywhere on the body, but when located on the lip, their distinctive coloration and texture help differentiate them from other types of lesions. The color and properties of hemangiomas often make them quite noticeable, leading to their identification when performing examinations. The diagnosis is further supported by their common presentation in various age groups and the fact that they typically do not cause any significant symptoms unless they are large or in a location that creates functional issues. In contrast, other conditions listed have distinct characteristics that do not align with a purplish lesion. For example, fibromas are usually firm, skin-colored or slightly pigmented nodules that do not display the vascular characteristics seen in hemangiomas. Squamous cell carcinoma may present as an ulcerated or scaly lesion without the distinct purplish coloration, often associated with a history of sun exposure or other risk factors. Psoriasis is typically characterized by well-defined, scaly plaques that are usually silvery in appearance and do not present as vascular lesions. Thus

10. What dental condition can result from inflammation of the tissue surrounding a tooth?

A. Gingivitis

B. Periodontitis

C. Abscess

D. Canker sore

The condition that arises from inflammation of the tissue surrounding a tooth is periodontitis. This condition progresses from gingivitis, which is an initial form of gum disease characterized by inflammation of the gums without loss of supporting bone. As gingivitis is left untreated, it can advance to periodontitis, where the inflammation extends deeper into the bone and supporting structures around the teeth, leading to tissue and bone loss. Periodontitis serves as a significant dental concern because it can affect the stability of teeth and is associated with various systemic health issues. The inflammation in periodontitis is generally a response to the presence of plaque, bacteria, and the body's immune response, which can lead to further complications if not properly managed. In contrast, the other options refer to different conditions: gingivitis is the earlier stage of gum disease; an abscess is a localized collection of pus that can occur due to infection, while canker sores are small, painful ulcers in the mouth that are not directly related to periodontal tissue inflammation. Understanding the progression and distinctions between these conditions aids in effective diagnosis and treatment planning.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adexdse.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE