
ADULT ARC CPR/AED/ FIRST AID CERT PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adultarccpraedfirstaid.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the compression-to-breath ratio for adult CPR?

A. 15 compressions to 2 breaths

B. 30 compressions to 1 breath

C. 30 compressions to 2 breaths

D. 50 compressions to 1 breath

2. What is an important step in managing an asthma attack?

A. Using a spacer with the inhaler if needed

B. Providing oxygen through a mask immediately

C. Performing chest compressions

D. Giving a mixture of fluids

3. Which symptom indicates a possible sudden illness that may require urgent care?

A. Fatigue

B. Extreme thirst

C. Chest pain

D. Runny nose

4. Upon detecting a life-threatening sudden illness, what should you do after ensuring safety?

A. Start CPR immediately

B. Assist with medications

C. Call 9-1-1 and seek help

D. Ignore the symptoms

5. Which of the following is a sign of shock?

A. Increased energy levels

B. Pale or clammy skin

C. Slight fever

D. Excited behavior

SAMPLE

6. What is the first action you take in a choking adult who is unable to speak?

A. Place the person in a sitting position

B. Perform the Heimlich maneuver

C. Provide water to the person

D. Call emergency services

7. How can you identify if someone is having a heart attack?

A. They will have a faint pulse

B. They may experience chest pain, shortness of breath, or sweating

C. They will likely become unconscious immediately

D. They will have a sudden headache

8. What is a common sign of anaphylaxis?

A. Stomach pain

B. Severe headache

C. Difficulty breathing

D. Muscle soreness

9. What is the correct action to take before applying AED pads to an older adult wearing a sweatshirt?

A. Apply the pads over the sweatshirt

B. Leave the sweatshirt on

C. Remove the sweatshirt

D. Cut the sweatshirt for access

10. Which area is most helpful for assessing changes in skin color?

A. Face

B. Less pigmented areas

C. Arms

D. Legs

SAMPLE

Answers

SAMPLE

1. C
2. A
3. C
4. C
5. B
6. B
7. B
8. C
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What is the compression-to-breath ratio for adult CPR?

A. 15 compressions to 2 breaths

B. 30 compressions to 1 breath

C. 30 compressions to 2 breaths

D. 50 compressions to 1 breath

The compression-to-breath ratio for adult CPR is 30 compressions to 2 breaths. This guideline is grounded in evidence-based practices that emphasize a balance between chest compressions and rescue breaths to optimize blood flow and oxygen delivery during cardiac arrest situations. In adult CPR, high-quality chest compressions are paramount, as they maintain circulation and help preserve vital organ function until emergency medical services can take over. The 30:2 ratio was established to ensure that participants provide a sufficient number of compressions to circulate blood effectively while also delivering adequate breaths to oxygenate the blood. This approach allows for a continuous and effective cycle of compressions with minimal interruptions for breaths, thereby maximizing the potential for survival. Other ratios do not conform to the current guidelines for adult victims undergoing CPR, making it essential to stick to the established 30 compressions followed by 2 breaths during resuscitation efforts.

2. What is an important step in managing an asthma attack?

A. Using a spacer with the inhaler if needed

B. Providing oxygen through a mask immediately

C. Performing chest compressions

D. Giving a mixture of fluids

Using a spacer with the inhaler during an asthma attack is an important step because it ensures that the medication is delivered effectively to the lungs. A spacer is a device that holds the medication released from an inhaler, allowing the person to inhale the medicine more easily and deeply. This helps in reducing the amount of medication that gets deposited in the mouth and throat, enhancing the amount that reaches the lungs where it is needed most. Effective delivery of the medication can alleviate the symptoms of the asthma attack more rapidly. Other methods like providing oxygen or performing chest compressions are not standard immediate responses to an asthma attack, as the primary goal is to use bronchodilators (inhalers) to relieve airway constriction. Giving fluids can also be inappropriate in this scenario, as it does not directly address the issue of airway constriction or bronchospasm that occurs during an asthma attack.

3. Which symptom indicates a possible sudden illness that may require urgent care?

A. Fatigue

B. Extreme thirst

C. Chest pain

D. Runny nose

Chest pain is a significant symptom that often indicates a potential acute medical emergency, especially if it is severe or accompanied by other distressing signs such as shortness of breath, dizziness, or radiating pain. It could suggest serious conditions like a heart attack, angina, or other cardiovascular issues that require immediate medical evaluation and intervention. Recognizing this symptom is crucial in emergency situations as timely treatment can be life-saving. In contrast, while fatigue can indicate various health issues, it is not necessarily urgent by itself. Extreme thirst can be a symptom of conditions like dehydration or diabetes but does not always require immediate care. A runny nose is typically associated with minor ailments such as allergies or colds, which are not urgent. Thus, chest pain stands out as the most urgent symptom necessitating immediate attention.

EXAMPLE

4. Upon detecting a life-threatening sudden illness, what should you do after ensuring safety?

A. Start CPR immediately

B. Assist with medications

C. Call 9-1-1 and seek help

D. Ignore the symptoms

In the case of a life-threatening sudden illness, calling 9-1-1 and seeking professional help is essential as it ensures that advanced medical assistance is on the way. This is particularly important because life-threatening conditions, such as a heart attack, stroke, or severe allergic reactions, can deteriorate rapidly. Emergency medical personnel are equipped to provide treatments that are critical in these situations, including administering oxygen, medications, or advanced life support. Additionally, calling for emergency services allows you to focus on the individual in need without delay. While performing CPR or assisting with medications may be necessary in some circumstances, those actions should come after ensuring help is on the way, ensuring that you are not compromising the patient's safety during this critical time. Ignoring the symptoms is not an option, as this could lead to dire consequences and put the person's life at risk.

5. Which of the following is a sign of shock?

A. Increased energy levels

B. Pale or clammy skin

C. Slight fever

D. Excited behavior

A sign of shock is pale or clammy skin because shock results in reduced blood flow and oxygen to the skin and extremities. This physiological response often leads to the skin losing its warmth and moisture, appearing pale or cool and clammy to the touch. Recognizing this sign is critical for diagnosing shock, as it indicates that the body is not adequately circulating blood. The other options do not indicate shock; increased energy levels and excited behavior can reflect a heightened state of arousal rather than distress, while a slight fever may suggest an infection or other medical issue, rather than shock itself.

6. What is the first action you take in a choking adult who is unable to speak?

A. Place the person in a sitting position

B. Perform the Heimlich maneuver

C. Provide water to the person

D. Call emergency services

When an adult is choking and unable to speak, the first action you should take is to perform the Heimlich maneuver, also known as abdominal thrusts. This technique is specifically designed to create an artificial cough, which can help expel the object blocking the airway. Rapid intervention is crucial in these situations, as a person cannot breathe or speak if their airway is fully obstructed, which can lead to loss of consciousness or even death within minutes. While placing the person in a sitting position could make them more comfortable, it does not address the immediate threat of airway obstruction. Providing water is also not an appropriate response, as it could exacerbate the choking situation by introducing more fluid into the airway. Calling emergency services is important but serves as a secondary action; the priority must be to clear the obstruction right away while waiting for professional assistance to arrive. Thus, performing the Heimlich maneuver is the correct and most effective immediate action in this emergency scenario.

EXAMPLE

7. How can you identify if someone is having a heart attack?

A. They will have a faint pulse

B. They may experience chest pain, shortness of breath, or sweating

C. They will likely become unconscious immediately

D. They will have a sudden headache

Identifying whether someone is having a heart attack is crucial for prompt and effective response. The most common symptoms include chest pain, shortness of breath, and sweating, which can occur as the heart muscle faces reduced blood flow. Chest pain may feel like pressure, squeezing, or a fullness, and it can radiate to other areas such as the arms, back, neck, or jaw. Shortness of breath may occur with or without chest discomfort. Sweating, often described as cold sweats, is another significant indicator. While some may assume that a faint pulse or immediate loss of consciousness are definitive signs, these situations do not represent the most common presentations of a heart attack. Additionally, a sudden headache is not a typical symptom associated with heart attacks. Understanding the primary symptoms helps in recognizing the situation swiftly and can lead to quicker medical assistance, which is vital for better outcomes.

8. What is a common sign of anaphylaxis?

A. Stomach pain

B. Severe headache

C. Difficulty breathing

D. Muscle soreness

Difficulty breathing is a hallmark sign of anaphylaxis, which is a severe and potentially life-threatening allergic reaction. During anaphylaxis, the body's immune system overreacts to a substance it perceives as harmful, leading to the release of chemicals that can cause swelling of the airways, tightening of the throat, and difficulty in breathing. This tightening can also cause wheezing or stridor, which further illustrates the critical nature of this symptom. The other signs mentioned, such as stomach pain, severe headache, or muscle soreness, while they may occur in different conditions or reactions, are not signature indicators of anaphylaxis. Stomach pain may arise from food allergies but is not as critical as breathing difficulties. Similarly, headaches and muscle soreness can be associated with a variety of other health issues but do not directly indicate anaphylactic shock. Recognizing difficulty breathing as a consequence of anaphylaxis is crucial for timely intervention, such as administration of epinephrine and calling for emergency medical assistance.

9. What is the correct action to take before applying AED pads to an older adult wearing a sweatshirt?

A. Apply the pads over the sweatshirt

B. Leave the sweatshirt on

C. Remove the sweatshirt

D. Cut the sweatshirt for access

When using an AED, it is crucial to ensure that the pads have direct contact with the skin for effective defibrillation. Removing the sweatshirt allows for the proper placement of the pads on the bare chest, ensuring that there is no barrier that could interfere with the electrical shock intended to restore a normal heart rhythm. While leaving the sweatshirt on, placing pads over it, or simply cutting it could potentially create issues with pad adhesion or effectiveness, removing the garment entirely mitigates any variables that could compromise the AED's function. Adequate skin contact is vital, and it's essential to expose the area where the pads will be applied, which is why removing the sweatshirt is the most appropriate action in this scenario.

SAMPLE

10. Which area is most helpful for assessing changes in skin color?

A. Face

B. Less pigmented areas

C. Arms

D. Legs

The area that is most helpful for assessing changes in skin color is less pigmented areas. These regions, such as the palms of the hands, soles of the feet, and the inner surfaces of the eyelids, typically show changes in color more distinctly. This is particularly important in emergency situations or in assessing conditions like cyanosis or jaundice, where changes in blood oxygenation or liver function can manifest as color alterations in the skin. Less pigmented areas have less melanin, making any changes in blood flow or oxygenation more noticeable. For instance, if someone is experiencing a lack of oxygen, these areas may turn a bluish color, which is more evident than in darker pigmented areas where such changes can be harder to discern. Conversely, while the face, arms, and legs can show changes in color, they can also be influenced by factors like natural skin tone, bruising, or external conditions, which might obscure the assessment of critical changes in skin color. Therefore, focusing on less pigmented areas provides a clearer and more reliable indication of the person's health status during an emergency assessment.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adultarccpraedfirstaid.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE