ADULT CCRN PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adultccrn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which option correctly describes the purpose of ICP monitoring?

A. To assess and manage increased intracranial pressure, which can lead to brain damage.

B. To measure systemic arterial oxygen saturation

C. To monitor peripheral nerve conduction

D. To evaluate liver function

2. What is cardiac tamponade?

A. Obstruction of the coronary arteries

B. Enlargement of the heart due to dilated cardiomyopathy

C. Inflammation of the pericardium without effusion

D. Compression of the heart due to blood or fluid accumulation between the myocardium and pericardium

3. What is the definition of acute coronary syndrome?

A. A condition where blood flow to a part of the heart is blocked briefly but heart muscle remains intact

B. A chronic inflammatory condition of the arteries

C. A sudden decrease in blood pressure causing syncope

D. A condition where blood flow to a part of the heart is blocked long enough that part of the heart muscle is damaged or dies

4. What is the purpose of administering nitroglycerin in this context?

A. To dilate cerebral arteries

B. To dilate coronary arteries

C. To increase heart rate

D. To decrease systemic vascular resistance

5. In traumatic brain injury, CPP should be maintained to prevent secondary brain injury. Which option best reflects this goal?

A. To minimize CPP to reduce brain edema

B. To maintain adequate CPP and prevent secondary brain injury

C. To maximize CPP regardless of intracranial pressure

D. To ignore perfusion status

SAMPLE

6. What is the function of inotropic agents?

A. They increase heart rate

B. They decrease conduction velocity

C. They affect the force of muscle contraction

D. They change venous return

7. Autonomic dysreflexia management includes which action?

A. Elevate the head of the bed

B. Administer vasopressors

C. Increase physical activity

D. Eliminate the cause, such as relieving bladder distension or treating skin irritation

8. Which vasopressor provides both inotropic and vasopressor effects by stimulating alpha and beta-1 receptors?

A. Epinephrine

B. Dopamine

C. Norepinephrine (Levophed)

D. Phenylephrine

9. In diabetic ketoacidosis, what is the primary cause of intracellular dehydration?

A. Osmotic diuresis from hyperglycemia.

B. Decreased insulin production.

C. Elevated blood glucose level.

D. Decreased serum osmolality.

10. Which combination drug is used in angiotensin receptor-neprilysin inhibitors (ARNIs)?

A. Enalapril, lisinopril

B. Sacubitril/valsartan (Entresto)

C. Valsartan/losartan

D. Sacubitril/losartan

SAMPLE

Answers

SAMPLE

1. A
2. D
3. D
4. B
5. B
6. C
7. D
8. C
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. Which option correctly describes the purpose of ICP monitoring?

A. To assess and manage increased intracranial pressure, which can lead to brain damage.

B. To measure systemic arterial oxygen saturation

C. To monitor peripheral nerve conduction

D. To evaluate liver function

Intracranial pressure monitoring is designed to continuously assess the pressure inside the skull so clinicians can detect rising pressure early and guide interventions to prevent secondary brain injury. When ICP increases, cerebral perfusion pressure (CPP) can drop, reducing blood flow to brain tissue and increasing the risk of ischemia or herniation; monitoring helps maintain adequate CPP by guiding therapies that lower ICP or support blood pressure. Normal ICP is roughly 5–15 mm Hg, and interventions aim to keep ICP within the normal range while preserving sufficient CPP (often targeting CPP above about 60–70 mm Hg). The data from ICP monitoring informs actions such as adjusting drainage with a ventriculostomy, optimizing head position and ventilation, and using osmotic therapies or sedation to control pressure. The other options describe measures of oxygenation, nerve function, or liver function, not intracranial pressure, so they do not describe the purpose of ICP monitoring.

2. What is cardiac tamponade?

A. Obstruction of the coronary arteries

B. Enlargement of the heart due to dilated cardiomyopathy

C. Inflammation of the pericardium without effusion

D. Compression of the heart due to blood or fluid accumulation between the myocardium and pericardium

Tamponade happens when fluid or blood collects in the pericardial sac enough to squeeze the heart from the outside. That external pressure prevents the ventricles from filling properly during diastole, so the amount the heart can pump (stroke volume) drops and overall cardiac output decreases. Because the pericardial pressure is transmitted to all chambers, diastolic pressures tend to equalize and venous return is impaired, leading to symptoms like low blood pressure, raised neck venous pressure, and often muffled heart sounds (Beck's triad); a rapid rise in pressure can also cause a noticeable drop in systolic blood pressure with inspiration (pulsus paradoxus) and may show electrical changes on the monitor such as electrical alternans. Emergent relief comes from removing the fluid, typically with pericardiocentesis or surgical drainage, while supporting circulation. This phenomenon is distinct from blockage of a coronary artery, enlargement from dilated cardiomyopathy, or inflammation of the pericardium without fluid.

SAMPLE

3. What is the definition of acute coronary syndrome?

A. A condition where blood flow to a part of the heart is blocked briefly but heart muscle remains intact

B. A chronic inflammatory condition of the arteries

C. A sudden decrease in blood pressure causing syncope

D. A condition where blood flow to a part of the heart is blocked long enough that part of the heart muscle is damaged or dies

Acute coronary syndrome means a sudden reduction in blood flow to a portion of the heart that is severe enough to cause myocardial injury, with cell death if reperfusion isn't achieved promptly. The crucial idea is the duration and consequence of the blockage: when flow is blocked long enough to injure or kill heart muscle, you have an infarction, which is the hallmark of ACS. If the blockage is brief and the heart muscle remains intact, that scenario is more consistent with transient ischemia or angina, not ACS. ACS encompasses unstable angina, NSTEMI, and STEMI, with the distinguishing feature being whether myocardial necrosis has occurred (often indicated by biomarkers and ECG changes). The other statements describe a brief blockage without damage, a chronic arterial inflammatory process, or a systemic cause of syncope—none of which define ACS.

4. What is the purpose of administering nitroglycerin in this context?

A. To dilate cerebral arteries

B. To dilate coronary arteries

C. To increase heart rate

D. To decrease systemic vascular resistance

Nitroglycerin is given to improve blood flow to the ischemic heart by dilating the coronary arteries, increasing perfusion to already stressed myocardium. It releases nitric oxide, causing relaxation of vascular smooth muscle. In the context of chest pain or suspected ischemia, this coronary dilation helps relieve ischemia and angina. Nitroglycerin also lowers preload and afterload (through venodilation and some arterial dilation), which reduces the heart's oxygen demand, but the key therapeutic aim here is to enhance coronary perfusion to the ischemic tissue.

SAMPLE

5. In traumatic brain injury, CPP should be maintained to prevent secondary brain injury. Which option best reflects this goal?

A. To minimize CPP to reduce brain edema

B. To maintain adequate CPP and prevent secondary brain injury

C. To maximize CPP regardless of intracranial pressure

D. To ignore perfusion status

Maintaining cerebral perfusion pressure to ensure reliable cerebral blood flow and prevent secondary injury from ischemia is the key idea. CPP is the pressure that drives blood through the brain and is calculated as mean arterial pressure minus intracranial pressure. In traumatic brain injury, the goal isn't to maximize blood flow at all costs or to shrink perfusion entirely; it's to keep CPP in a safe range so the brain receives enough oxygen and nutrients without letting intracranial pressure rise unchecked. If CPP falls too low, brain tissue becomes ischemic, triggering secondary brain injury. If CPP is pushed too high while ICP is elevated, you can worsen edema and intracranial hypertension, also worsening injury. So the best approach is to maintain adequate CPP by supporting MAP and controlling ICP, thereby protecting brain tissue from secondary damage. In practice, that means preventing hypotension, targeting a CPP roughly around 60–70 mmHg (adjusted to the individual patient), and using ICP-limiting strategies such as head-of-bed elevation, appropriate sedation, osmotic therapy, CSF drainage if needed, and other neurocritical care measures. The other ideas—minimizing CPP to reduce edema, maximizing CPP regardless of ICP, or ignoring perfusion status—risk ischemia or worsening intracranial hypertension, making them unsuitable.

6. What is the function of inotropic agents?

A. They increase heart rate

B. They decrease conduction velocity

C. They affect the force of muscle contraction

D. They change venous return

The principal idea here is contractility—the strength of the heart's muscle contraction. Inotropic agents directly influence this force. Positive inotropes raise the contractile strength by increasing the availability of calcium to the cardiac myocytes or by making the contractile apparatus more responsive to calcium, leading to a stronger squeeze and higher stroke volume. Negative inotropes do the opposite, reducing contractility. This is why inotropes are used to boost cardiac output in heart failure or shock. They may have other effects—some can alter heart rate (chronotropy) or conduction or preload indirectly—but their defining function is to change the force of contraction. They do not primarily change venous return, which is more about preload and venous tone.

7. Autonomic dysreflexia management includes which action?

A. Elevate the head of the bed

B. Administer vasopressors

C. Increase physical activity

D. Eliminate the cause, such as relieving bladder distension or treating skin irritation

Autonomic dysreflexia is a life threatening reflex that occurs in people with high spinal cord injuries when a noxious stimulus below the level of injury triggers a sudden, excessive sympathetic response, causing a dangerous rise in blood pressure. The most important management step is to identify and eliminate the triggering stimulus. By relieving bladder distension, emptying the bowel, removing tight clothing, or treating skin irritation, you remove the source of the abnormal signaling that keeps driving the dangerous hypertension. Once the trigger is removed, you can then support blood pressure by positioning the patient upright to help lower pressure and monitor closely, addressing further needs as needed. Administering vasopressors would worsen the situation, and increasing physical activity would introduce more stimuli, so those are not appropriate primary actions. The emphasis is on stopping the trigger, which stops the reflex at its source.

SAMPLE

8. Which vasopressor provides both inotropic and vasopressor effects by stimulating alpha and beta-1 receptors?

A. Epinephrine

B. Dopamine

C. Norepinephrine (Levophed)

D. Phenylephrine

Activating both alpha-1 and beta-1 receptors provides both vasoconstriction and increased cardiac contractility. Alpha-1 stimulation raises systemic vascular resistance, which boosts blood pressure, while beta-1 stimulation increases the force of heart contractions (inotropy) and can modestly raise heart rate, helping improve cardiac output. The drug that most reliably combines strong alpha-1–driven vasoconstriction with beta-1–driven inotropy is norepinephrine. Its alpha-1 effects produce the needed vasoconstriction to raise blood pressure, and its beta-1 effects enhance contractility to support forward flow, with minimal beta-2–mediated vasodilation compared with epinephrine. Other options either lack substantial inotropic effect (pure alpha-1 vasoconstrictor), have dose-dependent or broader receptor actions (dopamine), or include beta-2–mediated vasodilation (epinephrine), making norepinephrine the best match for those receptor targets.

9. In diabetic ketoacidosis, what is the primary cause of intracellular dehydration?

A. Osmotic diuresis from hyperglycemia.

B. Decreased insulin production.

C. Elevated blood glucose level.

D. Decreased serum osmolality.

When blood glucose rises dramatically in diabetic ketoacidosis, plasma osmolality increases. This creates an osmotic gradient that pulls water out of cells into the extracellular space to balance the forces, leading to intracellular dehydration. The high glucose level is the direct driver of this osmotic shift, making it the best explanation for why cells become dehydrated inside. Osmotic diuresis from the hyperglycemia contributes to overall fluid loss and volume depletion, but the intracellular water loss stems from the osmotic effect of the elevated glucose itself. Decreased serum osmolality would not cause this dehydration, since osmolality is increased, not decreased.

10. Which combination drug is used in angiotensin receptor-neprilysin inhibitors (ARNIs)?

A. Enalapril, lisinopril

B. Sacubitril/valsartan (Entresto)

C. Valsartan/losartan

D. Sacubitril/losartan

The key idea is that an ARNI combines a neprilysin inhibitor with an angiotensin receptor blocker to both boost protective natriuretic peptide effects and suppress the harmful RAAS. The drug that does this exact pairing is sacubitril/valsartan (Entresto), where sacubitril inhibits neprilysin and valsartan blocks the angiotensin II type 1 receptor. This combination reduces preload and afterload, improves hemodynamics, and has shown mortality and hospitalization benefits in heart failure with reduced ejection fraction. Enalapril and lisinopril are ACE inhibitors, which lower angiotensin II formation but do not provide neprilysin inhibition. Valsartan/losartan is two ARBs with no neprilysin inhibition. Sacubitril/losartan pairs neprilysin inhibition with an ARB, but the approved ARNI is sacubitril/valsartan, making it the correct choice.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adultccrn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE