
ADULT CRITICAL CARE SPECIALTY (ACCS) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adultcriticalcarespecialty.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is the purpose of a urinary catheter in critically ill patients?**

   A. To administer medication directly

   B. To monitor urine output and assess renal function

   C. To prevent urinary tract infections

   D. To facilitate bladder training

2. **What is the most rapid method to temporarily lower a patient's intracranial pressure?**

   A. Administration of corticosteroids

   B. Intentional hyperventilation

   C. Insertion of an intracranial catheter

   D. Administration of loop diuretic

3. **In preparation for a difficult intubation with a video assist device, what should a specialist also obtain?**

   A. A preformed rigid stylet

   B. A fiberoptic bronchoscope

   C. A retrograde wire kit

   D. A gum elastic bougie

4. **A 18 y.o. male with a fractured femur develops confusion and dyspnea 36 hours later. What immediate intervention is advised?**

   A. Fluid restriction

   B. Inhaled flolan

   C. Heparin therapy

   D. Low tidal volume ventilation

5. **Which type of shock is characterized by severe vasodilation and pooling of blood in the extremities?**

   A. Hypovolemic shock

   B. Cardiogenic shock

   C. Obstructive shock

   D. Distributive shock

SAMPLE

6. The common chest X-ray findings of CHF include all except?

A. Bilateral lung opacity

B. Cardiomegaly

C. Increased vascular markings

D. Durable pleural effusion

7. After intubation, which method confirms endotracheal tube placement?

A. Chest movement

B. Breath sounds

C. PetCo2

D. Condensation in the tube

8. Which patient position is often used to improve oxygenation in patients with severe respiratory distress?

A. Supine position

B. Sitting upright

C. Prone position

D. Semi-Fowler's position

9. In which scenario would the patient most likely be diagnosed with congestive heart failure based on vital signs and available data?

A. Elevated PCWP, low MAP, and low Qt

B. Low CVP, high PAP, and high PVR

C. Low MAP, low CVP, and low PV

D. Elevated PAP, elevated SVR, and low PVR

10. Neuromuscular blocking agents primarily affect which types of muscles?

A. Intercostal muscles

B. Cardiac muscle

C. Diaphragm

D. Skeletal muscle

SAMPLE

Answers

SAMPLE

1. B
2. D
3. A
4. B
5. D
6. D
7. C
8. D
9. A
10. D

SAMPLE

Explanations

SAMPLE

1. What is the purpose of a urinary catheter in critically ill patients?

A. To administer medication directly

B. To monitor urine output and assess renal function

C. To prevent urinary tract infections

D. To facilitate bladder training

The purpose of a urinary catheter in critically ill patients primarily focuses on monitoring urine output and assessing renal function. In a critical care setting, it is essential to accurately measure urine output because it serves as an important indicator of kidney function and overall fluid balance. Monitoring urine output can help clinicians detect early signs of renal impairment or fluid overload, which is crucial when managing critically ill patients who may have various comorbidities or conditions affecting renal function. This data also aids in adjusting fluid resuscitation and the administration of medications that may impact renal perfusion. While the other options may present related concepts, they do not capture the primary role of urinary catheters in this specific context. For example, while urinary tract infections can occur in patients with catheters, the catheter's primary function is not to prevent these infections but rather to provide critical data regarding kidney health. Similarly, administering medication directly through a catheter is not standard practice, nor is facilitating bladder training relevant in the context of critically ill patients, who typically have urgent medical needs that take precedence over such interventions.

2. What is the most rapid method to temporarily lower a patient's intracranial pressure?

A. Administration of corticosteroids

B. Intentional hyperventilation

C. Insertion of an intracranial catheter

D. Administration of loop diuretic

The most rapid method to temporarily lower a patient's intracranial pressure is through the administration of a loop diuretic. Loop diuretics, such as furosemide, can effectively decrease intracranial pressure by promoting diuresis, which leads to a reduction in cerebral edema and a decrease in the volume of cerebrospinal fluid (CSF). This quick action is essential in critical situations when intracranial pressure must be managed rapidly to prevent further neurological damage. In contrast, corticosteroids can reduce inflammation and cerebral edema over time, but they do not provide an immediate effect. Intentional hyperventilation can lower intracranial pressure by inducing hypocapnia (reduced carbon dioxide levels), leading to vasoconstriction, but this method is generally slower and less effective in emergency conditions compared to the immediate effects seen with after administering loop diuretics. While the insertion of an intracranial catheter is essential for monitoring and managing intracranial pressure, it is an invasive procedure that does not immediately lower pressure and carries its own risks and complications. Therefore, administering a loop diuretic offers a rapid, effective measure to lower intracranial pressure in urgent clinical scenarios.

SAMPLE

3. In preparation for a difficult intubation with a video assist device, what should a specialist also obtain?

A. A preformed rigid stylet

B. A fiberoptic bronchoscope

C. A retrograde wire kit

D. A gum elastic bougie

Obtaining a preformed rigid stylet is key in preparation for a difficult intubation because it aids in providing a more stable and controlled means of guiding the endotracheal tube into the trachea. The rigid stylet maintains the shape of the tube, which is particularly advantageous in situations where anatomy might be distorted due to either physiological conditions or external factors. This tool allows for optimal positioning and alignment of the endotracheal tube with the trachea, significantly improving the chances of successful intubation in challenging scenarios, such as those presented by limited neck mobility, anatomical anomalies, or anterior laryngeal positioning. When approaching intubation procedures, having a variety of tools available is crucial because conditions can change rapidly. While devices like fiberoptic bronchoscope, retrograde wire kits, and gum elastic bougies have their respective roles in aiding intubation, the primary function of a preformed rigid stylet is to facilitate the passage of the endotracheal tube itself during a difficult intubation, making it an essential tool in this context.

4. A 18 y.o. male with a fractured femur develops confusion and dyspnea 36 hours later. What immediate intervention is advised?

A. Fluid restriction

B. Inhaled flolan

C. Heparin therapy

D. Low tidal volume ventilation

The scenario describes an 18-year-old male with a fractured femur who develops confusion and dyspnea 36 hours post-injury, which raises concerns about potential fat embolism, a well-known complication of long bone fractures. In cases of fat embolism syndrome, patients can experience respiratory distress and neurological symptoms, typically manifesting within 1 to 3 days after the injury. Inhaled Flolan, which is a prostacyclin (i.e., a vasodilator), helps improve oxygenation by dilating the pulmonary vasculature, thereby reducing the pulmonary artery pressure and improving blood flow in the lungs. It can be particularly beneficial for patients experiencing respiratory distress due to increased pulmonary vascular resistance or inflammation, which can occur with fat embolism. The use of inhaled Flolan in this scenario aims to address the immediate respiratory compromise and facilitate better oxygen delivery to the tissues. By improving ventilation-perfusion matching in the lungs, it can help mitigate the hypoxic effects associated with fat globules obstructing pulmonary capillaries. Thus, the choice of inhaled Flolan as the immediate intervention is justified given its role in managing the respiratory symptoms linked with potential fat embolism, ultimately improving patient outcomes in this acute setting.

SAMPLE

5. Which type of shock is characterized by severe vasodilation and pooling of blood in the extremities?

A. Hypovolemic shock

B. Cardiogenic shock

C. Obstructive shock

D. Distributive shock

Distributive shock is characterized by severe vasodilation and a relative hypovolemia due to the pooling of blood in the extremities. This type of shock occurs when there is a significant decrease in systemic vascular resistance, leading to impaired circulation and inadequate blood flow to vital organs. Conditions such as sepsis, anaphylaxis, and neurogenic factors can lead to this widespread vasodilation. In distributive shock, the heart may still be functioning normally, but the blood vessels are excessively dilated, which causes a mismatch between blood volume and the vascular space. As a result, blood may pool in the extremities, preventing adequate perfusion to the core organs, which mandates prompt identification and intervention to restore hemodynamic stability. Understanding the characteristics of different types of shock can help in rapidly identifying the appropriate management strategies, highlighting the importance of recognizing distributive shock's unique features, including the profound vasodilation and the consequent pooling of blood.

6. The common chest X-ray findings of CHF include all except?

A. Bilateral lung opacity

B. Cardiomegaly

C. Increased vascular markings

D. Durable pleural effusion

In the context of congestive heart failure (CHF), chest X-ray findings typically reveal distinct patterns due to fluid overload and heart enlargement. The key findings associated with CHF include bilateral lung opacity, indicative of pulmonary congestion or edema, cardiomegaly representing an enlarged heart due to volume overload, and increased vascular markings which signal elevated pressure in the pulmonary circulation. Durable pleural effusion, while it can occur in CHF patients due to fluid accumulation, is not a common finding that is routinely associated with CHF itself. It is more of a potential development that can arise secondarily, rather than a primary characteristic that one would expect to identify on a standard chest X-ray for CHF assessment. As a result, when evaluating chest X-rays for CHF, the presence of durable pleural effusion would be less definitive and specific to the direct effects of heart failure compared to the other listed findings.

SAMPLE

7. After intubation, which method confirms endotracheal tube placement?

A. Chest movement

B. Breath sounds

C. PetCo2

D. Condensation in the tube

Confirming endotracheal tube placement is crucial in ensuring adequate ventilation and avoiding potential complications. Evaluating the exhaled carbon dioxide levels using a device like a capnometer, commonly referred to as PetCO2, provides objective evidence of proper tube placement in the trachea. When the endotracheal tube is correctly positioned, CO2 from the patient's exhaled breath will be detected, indicating that the tube is in the airway rather than the esophagus. While chest movement, breath sounds, and condensation in the tube are useful tools in assessing tube placement, they are less definitive than the measurement of carbon dioxide. Chest movement might not show clear signs of effective ventilation if there are underlying respiratory issues. Breath sounds can be misleading, particularly in cases where there is bilateral lung disease or if the tube is partially obstructed. Condensation in the tube does not provide reliable confirmation of proper placement. Therefore, the presence of CO2 is the most specific and reliable indicator of successful endotracheal intubation, making it the best choice for confirming tube placement.

8. Which patient position is often used to improve oxygenation in patients with severe respiratory distress?

A. Supine position

B. Sitting upright

C. Prone position

D. Semi-Fowler's position

The semi-Fowler's position is often utilized to enhance oxygenation in patients experiencing severe respiratory distress. This position involves the patient being placed at a slight angle, typically between 30 to 45 degrees. Elevating the upper body can reduce pressure on the diaphragm, allowing for better lung expansion and ease of breathing. This facilitates increased chest wall compliance and reduces the work of breathing, which can be critical in managing respiratory distress. Additionally, the semi-Fowler's position can help improve venous return, decrease the risk of aspiration, and promote effective diaphragmatic movement. These factors collectively contribute to optimized gas exchange and better oxygenation, making this position beneficial for patients with respiratory challenges. In contrast, the supine position may restrict lung expansion and increase the work of breathing, especially in patients with respiratory issues. The sitting upright and prone positions can also be beneficial for certain conditions or specific patient needs but are not as universally applicable for all patients in distress as the semi-Fowler's position. The semi-Fowler's position is widely recommended in acute care settings for its balance between comfort and functional benefits in respiratory distress situations.

SAMPLE

9. In which scenario would the patient most likely be diagnosed with congestive heart failure based on vital signs and available data?

A. Elevated PCWP, low MAP, and low Qt

B. Low CVP, high PAP, and high PVR

C. Low MAP, low CVP, and low PV

D. Elevated PAP, elevated SVR, and low PVR

In the scenario where the patient presents with elevated pulmonary capillary wedge pressure (PCWP), low mean arterial pressure (MAP), and low cardiac output (Qt), the diagnosis of congestive heart failure becomes evident. Elevated PCWP is a critical marker indicating increased left atrial pressure, which often occurs due to left ventricular dysfunction. This elevation reflects the backup of blood flow due to the heart being unable to pump effectively, commonly seen in heart failure. The accompanying low MAP signifies that the patient's systemic blood pressure is inadequate, which is often a consequence of reduced cardiac output. Low cardiac output further indicates that the heart is not supplying sufficient blood to meet the body's demands, which is characteristic of congestive heart failure. In conjunction, these signs and symptoms collectively illustrate a poor hemodynamic status attributable to heart failure. The other scenarios described do not align with typical presentations of congestive heart failure. For example, low central venous pressure (CVP) or elevated pulmonary artery pressure (PAP) with high pulmonary vascular resistance (PVR) suggests other pathophysiological states rather than heart failure. Similarly, various combinations of low vascular parameters without corresponding elevations in pressures typically relate to conditions that do not signify congestive heart failure. Thus, the evidence from

10. Neuromuscular blocking agents primarily affect which types of muscles?

A. Intercostal muscles

B. Cardiac muscle

C. Diaphragm

D. Skeletal muscle

The correct choice is related to the nature of neuromuscular blocking agents and their primary effects on muscle types. Neuromuscular blocking agents specifically target skeletal muscles by interrupting the transmission of nerve impulses at the neuromuscular junction. This action leads to muscle paralysis, particularly affecting voluntary skeletal muscles that are responsible for movement. The intercostal muscles and diaphragm, although essential for respiration and considered as striated skeletal muscles, are indirectly affected since they are part of the skeletal muscle group. However, focusing specifically on the correct answer, it is essential to recognize that neuromuscular blockers are primarily utilized to induce relaxation of skeletal muscles during surgical procedures or mechanical ventilation. Cardiac muscle, on the other hand, functions differently; it is controlled by the autonomic nervous system and is not influenced by neuromuscular blockers, which specifically target the somatic nervous system affecting only skeletal muscles. In summary, the primary action of neuromuscular blocking agents is on skeletal muscles, including important muscles such as the diaphragm and intercostals, which are crucial for respiration, but are ultimately classified under the broader category of skeletal muscle dynamics.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adultcriticalcarespecialty.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE