
ADVANCED CARDIOVASCULAR LIFE SUPPORT (ACLS) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://acls.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. If an AED does not promptly analyze a rhythm, what is the next step to take?

A. Adjust the pads on the patient

B. Begin chest compressions

C. Check the patient's pulse

D. Switch to manual defibrillation

2. Which condition is characterized by irregular and rapid heart rhythms originating from the atria?

A. Ventricular fibrillation

B. Supraventricular tachycardia

C. Atrial fibrillation

D. Sinus tachycardia

3. What is the first treatment priority for a patient who achieves ROSC?

A. Administer medications

B. Optimizing ventilation and oxygenation

C. Begin cardiac rehabilitation

D. Transfer to ICU

4. What indicates the need for immediate defibrillation in an adult patient?

A. Presence of bradycardia

B. Presence of ventricular fibrillation or pulseless ventricular tachycardia

C. Presence of chest pain only

D. Presence of normal sinus rhythm

5. What is the recommended depth of chest compressions during adult CPR?

A. At least 1 inch (2.5 cm) deep

B. At least 2 inches (5 cm) deep

C. At least 2.5 inches (6.5 cm) deep

D. At least 3 inches (7.5 cm) deep

SAMPLE

6. Which of the following is a critical link in ensuring patient survival in adult cardiac emergencies?

A. Ventilation techniques

B. Early advanced life support

C. Rapid defibrillation

D. Post-cardiac arrest monitoring

7. A patient with pulseless ventricular tachycardia is defibrillated. What is the next action?

A. Start chest compressions at a rate of at least 100/min

B. Check for a pulse

C. Administer epinephrine

D. Monitor the ECG continuously

8. What does torsades refer to in the context of ventricular tachycardia?

A. Atrial fibrillation

B. Polymorphic ventricular tachycardia

C. Coarse ventricular fibrillation

D. Supraventricular tachycardia

9. Which of the following conditions constitutes symptomatic bradycardia?

A. Stable blood pressure

B. Presence of a clear airway

C. Hypotension

D. Regular heart rhythm

10. After verifying the absence of a pulse, you initiate CPR with adequate bag-mask ventilation. What is your next action?

A. Prepare for intubation

B. IV or IO access

C. Administer epinephrine

D. Check for a pulse again

SAMPLE

Answers

SAMPLE

1. B
2. C
3. B
4. B
5. B
6. C
7. A
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. If an AED does not promptly analyze a rhythm, what is the next step to take?

A. Adjust the pads on the patient

B. Begin chest compressions

C. Check the patient's pulse

D. Switch to manual defibrillation

When an Automated External Defibrillator (AED) does not promptly analyze a rhythm, it is essential to maintain the quality of chest compressions, as this is critical for preserving circulation and oxygenation to vital organs during a cardiac arrest scenario. Chest compressions should be initiated immediately to provide effective blood flow and increase the chances of a successful rhythm analysis once the AED is ready. In the context of an unresponsive patient in need of defibrillation, ensuring that chest compressions are ongoing is vital. The rhythm can then be analyzed again after a brief period of uninterrupted CPR to determine if a shockable rhythm is present. Initiating compressions also allows for the proper functioning of the AED if it is able to continue analyzing rhythms during the compressions, as it may not function optimally if the patient is not receiving adequate compressions. This response underscores the importance of following current resuscitation guidelines, which prioritize high-quality CPR, especially when an AED is not able to provide immediate analysis.

2. Which condition is characterized by irregular and rapid heart rhythms originating from the atria?

A. Ventricular fibrillation

B. Supraventricular tachycardia

C. Atrial fibrillation

D. Sinus tachycardia

Atrial fibrillation is a condition defined by irregular and often rapid heart rhythms that originate from the atria, the upper chambers of the heart. In this condition, the normal rhythmic contractions of the atria are disrupted, leading to chaotic electrical activity. This irregularity can result in an erratic heart rhythm that is typically identified on an electrocardiogram (ECG) by the absence of distinct P waves, which are normally seen in a healthy sinus rhythm. Instead, there may be indistinct, fibrillatory waves. The relationship between the atria and ventricles is also affected in atrial fibrillation, as the rapid impulses can cause the ventricles to respond with varying rates, further contributing to the irregularity of the heartbeat. This can lead to decreased cardiac efficiency and increased risk for thromboembolic events, such as stroke, due to the potential for blood clots forming in the atria. Other conditions mentioned involve different mechanisms of arrhythmias. While supraventricular tachycardia does involve rapid heart rhythms, it typically arises from areas above the ventricles and can include a regular rhythm. Ventricular fibrillation is a lethal arrhythmia that originates from the ventricles and is characterized by a loss of

SAMPLE

3. What is the first treatment priority for a patient who achieves ROSC?

A. Administer medications

B. Optimizing ventilation and oxygenation

C. Begin cardiac rehabilitation

D. Transfer to ICU

After a patient achieves Return of Spontaneous Circulation (ROSC), the immediate focus shifts to optimizing ventilation and oxygenation. This is crucial because, even after ROSC, the patient's body may still be in a state of oxygen debt due to the preceding cardiac arrest. Ensuring adequate ventilation and oxygenation helps to stabilize the patient, support brain recovery, and maintain organ function. In the post-ROSC phase, it's vital to monitor the patient's respiratory status closely. Providing supplemental oxygen, ensuring proper airway management, and monitoring the EtCO2 levels are essential components of this care. Optimizing these parameters can prevent hypoxia and contribute to better outcomes in recovery. While other options such as administering medications, cardiac rehab, or transferring to the ICU are important steps in the post-resuscitation care pathway, they are secondary to ensuring the patient is adequately ventilated and oxygenated first. Without proper ventilation and oxygenation, the effectiveness of subsequent treatments may be compromised.

4. What indicates the need for immediate defibrillation in an adult patient?

A. Presence of bradycardia

B. Presence of ventricular fibrillation or pulseless ventricular tachycardia

C. Presence of chest pain only

D. Presence of normal sinus rhythm

Immediate defibrillation is indicated specifically in the presence of ventricular fibrillation (VF) or pulseless ventricular tachycardia (VT). Both of these arrhythmias are life-threatening conditions that can lead to cardiac arrest and require urgent intervention to restore a normal rhythm. Ventricular fibrillation is characterized by chaotic electrical activity in the heart, preventing it from effectively pumping blood. Similarly, pulseless VT is a rapid heart rhythm that does not produce a measurable pulse, meaning the heart is also ineffective at maintaining circulation. In both cases, defibrillation delivers an electrical shock to the heart, which can interrupt the abnormal rhythm and allow the heart's natural pacemaker to regain control, potentially restoring a viable cardiac rhythm. Other conditions listed, such as bradycardia, chest pain, or normal sinus rhythm, do not warrant immediate defibrillation. Bradycardia refers to a slow heart rate, which may require other forms of intervention but not defibrillation. Chest pain may suggest myocardial ischemia or other cardiac problems but does not indicate the immediate need for defibrillation unless accompanied by the other arrhythmias mentioned. Normal sinus rhythm signifies a healthy heart rhythm, so there is no need for defibrillation. Thus,

SAMPLE

5. What is the recommended depth of chest compressions during adult CPR?

A. At least 1 inch (2.5 cm) deep

B. At least 2 inches (5 cm) deep

C. At least 2.5 inches (6.5 cm) deep

D. At least 3 inches (7.5 cm) deep

The recommended depth of chest compressions during adult CPR is at least 2 inches (5 cm) deep. This depth is essential to ensure effective circulation and to generate enough pressure to perfuse vital organs, particularly the heart and brain, during cardiac arrest. Compressing to this depth allows for adequate blood flow, which is critical during the unresponsive state of a person experiencing a cardiac emergency. Additionally, it's important to allow for complete chest recoil between compressions to maximize blood return to the heart. Achieving the correct depth can be challenging, and rescuers should aim to maintain a rate of 100 to 120 compressions per minute while ensuring that the compressions are both deep and consistent. Therefore, recognizing the significance of 2 inches as the minimum depth aligns with established guidelines to optimize the chances of survival and recovery after cardiac arrest.

6. Which of the following is a critical link in ensuring patient survival in adult cardiac emergencies?

A. Ventilation techniques

B. Early advanced life support

C. Rapid defibrillation

D. Post-cardiac arrest monitoring

Rapid defibrillation is a critical link in ensuring patient survival in adult cardiac emergencies, particularly in cases of shockable rhythms such as ventricular fibrillation (VF) and pulseless ventricular tachycardia (VT). Defibrillation delivers an electrical shock to the heart, aiming to restore a normal rhythm and effectively re-establish circulation. The sooner defibrillation is administered after the onset of a cardiac arrest, the higher the chances of survival; each minute that passes without defibrillation decreases the probability of successful resuscitation. In cardiac emergencies, the effectiveness of resuscitation efforts is significantly improved when defibrillation occurs within the critical time window, which is generally within the first few minutes post-cardiac arrest. The guidelines emphasize that achieving rapid defibrillation should be a priority in the chain of survival because data shows a direct correlation between the speed of defibrillation and patient outcomes. Other components of cardiac emergency response, such as ventilation techniques, early advanced life support, and post-cardiac arrest monitoring, are also important but come after the immediate action of defibrillation. While they contribute to overall patient care and can impact outcomes, they do not match the urgency and direct effect that rapid defibrillation has during the

SAMPLE

7. A patient with pulseless ventricular tachycardia is defibrillated. What is the next action?

A. Start chest compressions at a rate of at least 100/min

B. Check for a pulse

C. Administer epinephrine

D. Monitor the ECG continuously

Defibrillation is employed in the case of pulseless ventricular tachycardia to restore a normal cardiac rhythm. Following defibrillation, the immediate priority is to ensure effective circulation and oxygenation, which is achieved through chest compressions. Initiating high-quality chest compressions at a rate of at least 100 per minute is essential because it helps maintain blood flow to vital organs and supports survival until a normal rhythm can be established. This action is critical as it provides circulatory support during the crucial moments after defibrillation, especially given that the primary aim is to restore a perfusing rhythm. Effective chest compressions are proven to be invaluable in enhancing patient outcomes, especially in the early minutes after a cardiac event. While checking for a pulse might seem logical, it is generally recommended to continue compressions for a short period before reassessing, as pulse checks can interrupt circulation. Furthermore, administering epinephrine or continuous ECG monitoring may be important later in the resuscitation process, but they do not take precedence immediately after defibrillation. Hence, the focus should be on resuming chest compressions promptly.

8. What does torsades refer to in the context of ventricular tachycardia?

A. Atrial fibrillation

B. Polymorphic ventricular tachycardia

C. Coarse ventricular fibrillation

D. Supraventricular tachycardia

Torsades de pointes, often simply referred to as torsades, is a specific type of polymorphic ventricular tachycardia characterized by a distinctive undulating pattern of QRS complexes that change in amplitude and direction. This arrhythmia typically occurs in the context of a prolonged QT interval and can lead to more serious rhythms or even progress to ventricular fibrillation if not promptly treated. Understanding torsades is crucial for effective management, especially in emergency settings. It often requires interventions such as magnesium sulfate administration or addressing the underlying causes of the prolonged QT interval. Recognizing torsades as polymorphic ventricular tachycardia allows healthcare providers to differentiate it from other forms of tachycardia and implement appropriate treatment strategies swiftly.

9. Which of the following conditions constitutes symptomatic bradycardia?

A. Stable blood pressure

B. Presence of a clear airway

C. Hypotension

D. Regular heart rhythm

Symptomatic bradycardia is defined as a slower-than-normal heart rate that is accompanied by signs or symptoms indicating that the body is not receiving enough blood flow to meet its needs. Hypotension, or low blood pressure, is one of the key signs of decreased perfusion and inadequate circulation. When a patient presents with bradycardia and also has hypotension, this condition can lead to serious organ dysfunction or even shock, making it critical to address. Other conditions such as stable blood pressure or the presence of a clear airway may indicate a situation where the patient is stable, but they do not necessarily reflect the immediate threats posed by symptomatic bradycardia. In addition, a regular heart rhythm, while significant in assessment, does not imply that the heart rate is appropriate or that the patient is stable. Thus, hypotension is a defining factor indicating that the bradycardia is causing symptoms that require intervention.

SAMPLE

10. After verifying the absence of a pulse, you initiate CPR with adequate bag-mask ventilation. What is your next action?

A. Prepare for intubation

B. IV or IO access

C. Administer epinephrine

D. Check for a pulse again

After verifying the absence of a pulse and initiating CPR with adequate bag-mask ventilation, the critical next action is to establish IV (intravenous) or IO (intraosseous) access. This step is vital because it allows for the rapid administration of medications such as epinephrine and other necessary interventions, which are crucial for improving the patient's chances of survival during cardiac arrest. Establishing IV or IO access enables healthcare providers to deliver lifesaving drugs effectively and efficiently, especially in a resuscitation scenario where every second counts. The ACLS guidelines emphasize the importance of obtaining vascular access early in the resuscitation efforts as medications need to be administered to facilitate heart rhythm stabilization and enhance perfusion. While preparing for intubation is important in advanced airway management, it typically follows the establishment of IV access and the administration of medications. Administering epinephrine is a critical step as well, but it should be done after confirming that IV or IO access is in place. Checking for a pulse again is also essential, but in the immediate steps of resuscitation, establishing access to deliver medications takes precedence.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://acls.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE