
ADVANCED EMERGENCY MEDICAL TECHNICIAN (AEMT) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://
advancedemergencymedicaltechnician.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What should be monitored in a patient receiving nitroglycerin?

A. Blood glucose levels

B. Blood pressure

C. Heart rate

D. Respiratory rate

2. What substance keeps alveoli in the lungs open?

A. Tidal volume

B. Capillary sphincters

C. Expiratory pressure

D. Surfactant

3. Which condition could indicate a need for intubation during an allergic reaction?

A. Increased respiratory rate

B. Stridor and difficulty breathing

C. Swelling of the abdomen

D. Chest pain upon inhalation

4. What is the primary treatment for a patient who is unresponsive and not breathing?

A. Oxygen therapy

B. Chest compressions

C. Defibrillation

D. Endotracheal intubation

5. Which of the following is part of the Enhanced 911 system?

A. Dispatching law enforcement

B. Automatically dispatching an ambulance

C. Identifying the caller's location

D. Providing automatic call-back

SAMPLE

6. What is the typical dosage of epinephrine for a patient experiencing anaphylaxis?

A. 0.1 to 0.2 mg intramuscularly

B. 0.3 to 0.5 mg intramuscularly

C. 1.0 mg intravenously

D. 0.5 to 1.0 mg subcutaneously

7. The term dysrhythmia most accurately refers to:

A. An absence of heart activity

B. An irregular pulse rhythm

C. Inadequate cardiac output

D. Abnormality of electrical activity in the heart

8. A patient with severe dehydration may be treated with which fluid type?

A. Hypertonic saline

B. Isotonic crystalloid solution

C. Colloid solution

D. Hypotonic solution

9. All of the following are examples of solid medications except:

A. An acetaminophen capsule

B. Fine powdered form of albuterol in a metered-dose inhaler

C. Nitroglycerin paste

D. An aspirin tablet

10. Enteral routes of drug administration include?

A. Intramuscular injection

B. Oral

C. Intravenous bolus

D. Inhalation

SAMPLE

Answers

SAMPLE

1. B
2. D
3. B
4. B
5. C
6. B
7. D
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What should be monitored in a patient receiving nitroglycerin?

A. Blood glucose levels

B. Blood pressure

C. Heart rate

D. Respiratory rate

In a patient receiving nitroglycerin, monitoring blood pressure is crucial because nitroglycerin acts as a vasodilator, primarily causing dilation of blood vessels to reduce cardiac workload and relieve angina. As a result, the patient's blood pressure can significantly decrease following administration, potentially leading to hypotension. Regular monitoring allows the healthcare provider to detect any significant drops in blood pressure promptly, which can help avoid adverse effects such as dizziness, syncope, and other complications associated with excessive hypotension. While the other parameters like blood glucose levels, heart rate, and respiratory rate may provide useful information regarding the patient's overall condition, they do not have the same direct relationship with the administration of nitroglycerin and its immediate effects on the cardiovascular system. Therefore, maintaining vigilant blood pressure monitoring is essential for patient safety during nitroglycerin treatment.

2. What substance keeps alveoli in the lungs open?

A. Tidal volume

B. Capillary sphincters

C. Expiratory pressure

D. Surfactant

Surfactant is a critical substance produced by the cells lining the alveoli in the lungs. Its primary function is to reduce surface tension within the alveoli, which are the small air sacs where gas exchange occurs. By decreasing surface tension, surfactant prevents the alveoli from collapsing, particularly during exhalation when the lungs are at their smallest volume. This is essential for maintaining the stability of the alveoli and ensuring that they remain open and available for gas exchange with the bloodstream. Without sufficient surfactant, the alveoli would collapse (a condition known as atelectasis) and would not function properly, leading to impaired oxygenation of the blood. The other choices pertain to other aspects of the respiratory process but do not directly contribute to keeping the alveoli open. While tidal volume refers to the volume of air inhaled and exhaled during normal breathing, it does not influence the structural integrity of the alveoli. Capillary sphincters are involved in the regulation of blood flow in capillaries but do not have a role in maintaining alveolar structure. Expiratory pressure relates to the pressure during exhalation but likewise does not serve the specific function of preventing alveolar collapse.

SAMPLE

3. Which condition could indicate a need for intubation during an allergic reaction?

A. Increased respiratory rate

B. Stridor and difficulty breathing

C. Swelling of the abdomen

D. Chest pain upon inhalation

Intubation during an allergic reaction is typically indicated by the presence of stridor and difficulty breathing. Stridor is a high-pitched, wheezing sound resulting from turbulent airflow in the upper airway, often due to obstruction or swelling in that area. In the context of an allergic reaction, particularly anaphylaxis, edema can occur in the larynx and surrounding tissues, leading to this characteristic sound. Difficulty breathing suggests that the airway is compromised, and intubation may be necessary to ensure airway patency and adequate oxygenation. Increased respiratory rate can indicate distress or a compensatory mechanism in response to inadequate oxygenation, but it does not specifically point to a need for intubation on its own. Swelling of the abdomen may suggest gastrointestinal involvement of an allergic reaction but does not directly affect the airway. Chest pain upon inhalation could be indicative of a variety of respiratory issues but does not specifically signify the need for intubation in the context of an allergic reaction. Therefore, the presence of stridor and difficulty breathing is the most compelling reason for intubation in this scenario.

4. What is the primary treatment for a patient who is unresponsive and not breathing?

A. Oxygen therapy

B. Chest compressions

C. Defibrillation

D. Endotracheal intubation

The primary treatment for a patient who is unresponsive and not breathing is chest compressions. In such a situation, the absence of breathing indicates that the patient may be in cardiac arrest, which requires immediate initiation of cardiopulmonary resuscitation (CPR). Chest compressions are crucial because they help circulate blood to vital organs, particularly the brain and heart, until professional medical help arrives or the heart spontaneously resumes normal functioning. Oxygen therapy is not immediately effective in this scenario since the patient is not breathing, meaning that the delivery of oxygen would not be possible or useful without restoring adequate circulation first. Defibrillation is specifically indicated in cases of certain cardiac arrhythmias, such as ventricular fibrillation, but it is not the first step when a patient is unresponsive and not breathing. Endotracheal intubation might eventually be required to secure the airway, but it comes after initiating chest compressions and addressing basic life support measures. Therefore, starting chest compressions is the critical first action that significantly increases the patient's chances of survival.

SAMPLE

5. Which of the following is part of the Enhanced 911 system?

A. Dispatching law enforcement

B. Automatically dispatching an ambulance

C. Identifying the caller's location

D. Providing automatic call-back

The Enhanced 911 (E911) system is designed to improve emergency response by providing critical information to dispatchers. One of its key features is the ability to automatically identify and display the caller's location when a 911 call is made. This capability is crucial in emergency situations, as it allows responders to reach the scene as quickly as possible, even if the caller is unable to communicate their location due to various circumstances. In contrast, while dispatching law enforcement, automatically dispatching an ambulance, or providing automatic call-back may be part of overall emergency response protocols, these features are not specifically defined as elements of the Enhanced 911 system. The primary focus of E911 technology is the geographic identification of the caller, enhancing situational awareness and response accuracy for first responders.

6. What is the typical dosage of epinephrine for a patient experiencing anaphylaxis?

A. 0.1 to 0.2 mg intramuscularly

B. 0.3 to 0.5 mg intramuscularly

C. 1.0 mg intravenously

D. 0.5 to 1.0 mg subcutaneously

The typical dosage of epinephrine for a patient experiencing anaphylaxis is 0.3 to 0.5 mg administered intramuscularly. This dosage is consistent with guidelines from health organizations, which advocate for the use of intramuscular administration to ensure rapid absorption and a swift physiologic response in an emergency situation. Epinephrine acts as a vasoconstrictor and bronchodilator, making it effective in reversing the severe allergic reaction associated with anaphylaxis. The intramuscular route is preferred because it allows for better absorption into the bloodstream, particularly when the onset of anaphylactic symptoms can be rapid and life-threatening. In this context, while the other dosages may relate to different routes or situations involving epinephrine, they do not reflect the preferred and evidence-based approach for anaphylaxis management, which underscores the importance of administering 0.3 to 0.5 mg intramuscularly for effective treatment.

7. The term dysrhythmia most accurately refers to:

A. An absence of heart activity

B. An irregular pulse rhythm

C. Inadequate cardiac output

D. Abnormality of electrical activity in the heart

Dysrhythmia specifically refers to an abnormality of the heart's electrical activity. This term encompasses a variety of disturbances in the heart's rhythm, which can lead to problems such as irregular heartbeats, missed beats, or rapid heart rates. These disturbances can originate from various issues in the heart's conduction system, affecting how signals are transmitted through the heart muscle and leading to the aforementioned irregularities. Understanding dysrhythmia is crucial for identifying the underlying heart problems that can cause symptoms or complications. While an irregular pulse rhythm may indicate a dysrhythmia, the term itself is more comprehensive, as it directly relates to the electrical signaling within the heart rather than simply the external presentation of pulse. Additionally, conditions such as an absence of heart activity or inadequate cardiac output represent specific consequences or conditions resulting from dysrhythmias rather than defining the term itself.

SAMPLE

8. A patient with severe dehydration may be treated with which fluid type?

A. Hypertonic saline

B. Isotonic crystalloid solution

C. Colloid solution

D. Hypotonic solution

In cases of severe dehydration, the priority is to restore the body's fluid volume and electrolyte balance effectively. Isotonic crystalloid solutions, such as normal saline or lactated Ringer's solution, are commonly used for this purpose. These solutions have a similar osmolarity to blood plasma, which means they help to expand the intravascular volume without causing a significant shift of fluids between compartments in the body. The use of isotonic solutions is particularly important when dealing with dehydration because they can effectively rehydrate the patient while also maintaining a stable electrolyte environment. This is essential for normal cellular function and overall metabolic processes. Hypertonic saline, while it can be used in specific circumstances for rapid volume expansion or to manage cerebral edema, is not appropriate for general rehydration as it can draw fluid out of cells, potentially worsening dehydration. Colloid solutions contain larger molecules that can help to maintain intravascular volume but are typically not the first line for hydration in cases of dehydration. Hypotonic solutions could dilute the serum sodium and further complicate fluid and electrolyte balance, making them unsuitable for treating severe dehydration. Thus, isotonic crystalloid solutions are the best choice for effectively managing this condition.

9. All of the following are examples of solid medications except:

A. An acetaminophen capsule

B. Fine powdered form of albuterol in a metered-dose inhaler

C. Nitroglycerin paste

D. An aspirin tablet

Solid medications are typically characterized by their physical form, which includes tablets, capsules, and powders that can be compressed or shaped. In this context, nitroglycerin paste is not considered a solid medication; rather, it is a topical preparation that is applied to the skin for systemic absorption. This means it has a semisolid consistency, allowing for easy application and release of the medication over time. The other options are all forms of solid medications. An acetaminophen capsule contains solid medication in a gelatin shell, which dissolves in the stomach to release the drug. The fine powdered form of albuterol in a metered-dose inhaler, while it may appear as a liquid when dispensed, contains solid particles in suspension that deliver the medication to the lungs. An aspirin tablet is a classic example of a solid dosage form, where the drug is pressed into a specific shape for oral consumption.

10. Enteral routes of drug administration include?

A. Intramuscular injection

B. Oral

C. Intravenous bolus

D. Inhalation

The enteral route of drug administration involves delivering medications directly to the gastrointestinal tract. This is primarily achieved through oral consumption, where drugs are taken by mouth and absorbed through the digestive system. When a drug is administered orally, it passes through the stomach and intestines before entering the bloodstream, which allows for systematic distribution throughout the body. In the context of the given options, oral administration is the main method recognized as enteral. Other routes, such as intramuscular injection, intravenous bolus, and inhalation, do not involve the gastrointestinal tract and are therefore classified as parenteral or other routes of administration. Thus, the correct response highlights the significance of the oral route in the enteral administration category.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://advancedemergencymedicaltechnician.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE