
ADVANCED MEDICAL LIFE SUPPORT (AMLS) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://advancedmedicallifesupport.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What should be evaluated when assessing a pediatric patient according to AMLS guidelines?**

   A. Only vital signs

   B. Age and weight alone

   C. Developmental stage and body surface area

   D. Family medical history

2. **What key factor should be considered in transport decision-making for a patient requiring critical care?**

   A. Availability of medical staff at the transport destination

   B. Weather conditions during transport

   C. Transport time versus potential treatment benefits

   D. Type of transport vehicle available

3. **What condition is likely to be present in COPD patients with acute shortness of breath?**

   A. Pulmonary embolism

   B. Angina pectoris

   C. Angioedema

   D. Hypertensive crisis

4. **What does the SAMPLE acronym stand for in patient assessment?**

   A. Signs, Anticipated outcomes, Medications, Past history, Last check, Events

   B. Signs, Allergies, Medications, Past medical history, Last oral intake, Events leading to illness

   C. Sensations, Actions, Monitoring, Previous treatments, Lifestyle, Events

   D. Symptoms, Ailments, Medications, Previous treatments, Last inputs, Emergencies

5. **What is the primary importance of patient education in AMLS?**

   A. Understanding their condition and adherence to treatment plans

   B. Minimizing hospital admissions

   C. Encouraging physical activity

   D. Improving nutritional habits

SAMPLE

6. When facing a patient with a traumatic brain injury, what is a crucial intervention AMLS recommends?

A. Applying ice packs to the head

B. Keeping the patient flat on their back

C. Ensuring head elevation

D. Administering a pain relief medication

7. A patient with a family history of heart disease taking daily aspirin and has chest pain should be most concerned about:

A. Familial history of heart disease

B. Daily use of aspirin

C. Symptoms of nausea

D. Recent stent placement

8. How does AMLS recommend managing airway obstructions?

A. Using sedatives to relax the patient

B. Performing the Heimlich maneuver

C. Administering nasal steroids

D. Immediate intubation in all cases

9. A patient may present with wheezing and difficulty breathing due to which possible diagnosis?

A. Pneumonia

B. Anxiety attack

C. Asthma exacerbation

D. Heart failure

10. How does AMLS suggest managing a patient with suspected cardiac emergency?

A. Focusing exclusively on oxygen administration

B. Ensuring airway, breathing, and circulation; obtaining an ECG; and administering appropriate medication

C. Offering emotional support and reassurance

D. Prioritizing transport to the hospital over initial management

SAMPLE

Answers

SAMPLE

1. C
2. C
3. A
4. B
5. A
6. C
7. D
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What should be evaluated when assessing a pediatric patient according to AMLS guidelines?

A. Only vital signs

B. Age and weight alone

C. Developmental stage and body surface area

D. Family medical history

When assessing a pediatric patient according to AMLS guidelines, evaluating developmental stage and body surface area is critical because both factors significantly influence how a child presents with illness and how they should be treated. Developmental stage is essential as children of different ages have varying physiological responses and potential differences in their medical needs. For example, toddlers and adolescents have different levels of understanding, expectations of involvement in their care, and reactions to illness. Body surface area is particularly important in pediatrics for calculating medication dosages, fluid resuscitation, and other medical interventions, as children's bodies metabolize drugs and respond to treatments differently than adults. These measurements help healthcare providers make informed decisions in emergency situations where accurate dosing and appropriate interventions are crucial for patient safety and effectiveness of treatment. Evaluating only vital signs would provide incomplete information, as it would not capture the full clinical picture or the specific considerations required for effective pediatric care. Focusing exclusively on age and weight would also neglect developmental considerations that may impact the patient's management and treatment decisions. Lastly, while family medical history is useful in understanding the patient's background, it does not address the immediate needs and physiological assessment required in an acute situation like critical care or emergency medicine.

2. What key factor should be considered in transport decision-making for a patient requiring critical care?

A. Availability of medical staff at the transport destination

B. Weather conditions during transport

C. Transport time versus potential treatment benefits

D. Type of transport vehicle available

In critical care transport decision-making, evaluating the transport time in relation to potential treatment benefits is essential because it directly impacts a patient's survival and overall outcome. In emergency situations, rapid access to definitive care can make a significant difference, particularly for conditions requiring immediate intervention, such as traumatic injuries, cardiac events, or severe respiratory distress. Consideration of transport time emphasizes the urgency of the patient's condition and the necessity of balancing how quickly the patient can be moved against the potential advantages of advanced treatment available at the destination. If the time taken to transport exceeds the immediate benefit of reaching a facility with specialized care or resources, this could delay critical interventions that could stabilize the patient. While factors like the availability of medical staff, weather conditions, and the type of transport vehicle are certainly important in the context of safety and capability, they must be analyzed in light of how they influence the overall transport time and the urgency of care required. Thus, focusing on transport time versus treatment benefits aligns with the critical nature of patient care in emergency situations.

SAMPLE

3. What condition is likely to be present in COPD patients with acute shortness of breath?

A. Pulmonary embolism

B. Angina pectoris

C. Angioedema

D. Hypertensive crisis

In patients with Chronic Obstructive Pulmonary Disease (COPD) who experience acute shortness of breath, a pulmonary embolism is a condition that is particularly relevant. COPD is characterized by chronic airflow limitation, primarily due to emphysema or chronic bronchitis, which may predispose individuals to complications such as a pulmonary embolism. The presence of a pulmonary embolism can cause sudden worsening of respiratory symptoms because it effectively reduces the blood flow to parts of the lung, causing increased shortness of breath, hypoxemia, and in severe cases, can lead to respiratory failure. Additionally, the presence of a thrombus can exacerbate the already compromised pulmonary function of COPD patients, leading to a significant increase in difficulty breathing and often necessitating urgent medical intervention. The other options, while they may be serious conditions, are less directly linked to acute respiratory distress in the context of a pre-existing diagnosis of COPD. Pulmonary embolism is unique in its ability to acutely and significantly impact the respiratory status in these patients.

4. What does the SAMPLE acronym stand for in patient assessment?

A. Signs, Anticipated outcomes, Medications, Past history, Last check, Events

B. Signs, Allergies, Medications, Past medical history, Last oral intake, Events leading to illness

C. Sensations, Actions, Monitoring, Previous treatments, Lifestyle, Events

D. Symptoms, Ailments, Medications, Previous treatments, Last inputs, Emergencies

*The SAMPLE acronym is a critical tool used during patient assessments to gather important information about a patient's medical history and current situation. In this context, the elements represented by the acronym are as follows: - **Signs**: These are the observable indicators of a medical condition that can be detected during an assessment. - **Allergies**: Knowing a patient's allergies is vital, as it helps in avoiding medications or interventions that could cause an allergic reaction. - **Medications**: Documenting what medications a patient is taking helps to understand their ongoing treatment and any potential interactions or contraindications in emergency care. - **Past medical history**: A thorough understanding of a patient's past medical conditions provides context for their current health issues and can help tailor the immediate medical response. - **Last oral intake**: This information is particularly important when planning for potential surgical interventions or when assessing the risk of aspiration and other complications. - **Events leading to illness**: Identifying what happened prior to the current medical situation can offer crucial clues about the etiology of the condition and inform appropriate interventions. The inclusion of these specific dimensions within the SAMPLE acronym emphasizes a structured method of patient assessment that ensures comprehensive data collection, leading to better patient care and outcomes.*

SAMPLE

5. What is the primary importance of patient education in AMLS?

A. Understanding their condition and adherence to treatment plans

B. Minimizing hospital admissions

C. Encouraging physical activity

D. Improving nutritional habits

The primary importance of patient education in Advanced Medical Life Support (AMLS) lies in helping patients understand their condition and promoting adherence to treatment plans. This understanding is crucial because when patients are well-informed about their health status, they are more likely to comply with medical advice, recognize symptoms that require immediate attention, and make informed decisions about their care. Educated patients are empowered to manage their conditions more effectively, which can lead to better outcomes and may reduce the risk of emergencies that require advanced medical interventions. Effective communication and education foster a partnership between the healthcare team and the patient, enabling the latter to take an active role in their own health and treatment journey. This understanding also enhances the patient's ability to follow prescribed treatment regimens, which is essential for managing chronic conditions and preventing exacerbations that could lead to hospitalization or complications. While other factors such as minimizing hospital admissions, encouraging physical activity, and improving nutritional habits are certainly beneficial aspects of overall patient care, the cornerstone of effective management in AMLS is the patient's comprehension of their condition and commitment to their treatment plan.

6. When facing a patient with a traumatic brain injury, what is a crucial intervention AMLS recommends?

A. Applying ice packs to the head

B. Keeping the patient flat on their back

C. Ensuring head elevation

D. Administering a pain relief medication

In the management of a patient with a traumatic brain injury (TBI), ensuring head elevation is a crucial intervention. Elevating the head of the bed can help to decrease intracranial pressure (ICP), which is a significant concern in patients with TBI. Elevated positioning promotes venous outflow from the brain, thereby reducing pressure within the cranial cavity. This practice can also facilitate respiratory function and comfort, especially if the patient has other injuries or is at risk of airway compromise. Additionally, maintaining proper head positioning can help prevent additional complications such as secondary brain injury caused by increased intracranial pressure or impaired cerebral perfusion. The recommended angle of elevation is typically around 30 degrees, balancing the need for reducing ICP while ensuring adequate perfusion of the brain. Other interventions like applying ice packs to the head may provide some discomfort relief but do not directly address the critical issue of ICP. Keeping the patient flat on their back could potentially exacerbate ICP, while administering pain relief, although important for patient comfort, does not directly manage the physiological concerns associated with TBI. Therefore, head elevation is emphasized in AMLS guidelines as a fundamental step in improving outcomes for patients suffering from traumatic brain injuries.

SAMPLE

7. A patient with a family history of heart disease taking daily aspirin and has chest pain should be most concerned about:

A. Familial history of heart disease

B. Daily use of aspirin

C. Symptoms of nausea

D. Recent stent placement

When assessing the clinical scenario presented, the most significant concern for a patient with a family history of heart disease who is experiencing chest pain is the recent stent placement. This context is critical because the presence of chest pain in someone with a recent intervention like stenting could indicate potential complications, such as stent thrombosis or restenosis, which could lead to serious cardiac events. Stents are placed to keep coronary arteries open, and if complications arise, they can significantly increase the risk of myocardial infarction (heart attack). Therefore, the presence of chest pain in this specific context warrants immediate evaluation to rule out any acute issues related to the stent and ensure the proper vascular flow is maintained. While a family history of heart disease and the daily use of aspirin are important factors to consider in the overall risk stratification for heart disease, they do not present an immediate concern compared to the risks linked with recent stent placement. Symptoms of nausea can also occur during cardiac events but are not as directly related to the urgency posed by the recent stent placement. Hence, focusing on the potential complications arising from the stent is crucial in this case.

8. How does AMLS recommend managing airway obstructions?

A. Using sedatives to relax the patient

B. Performing the Heimlich maneuver

C. Administering nasal steroids

D. Immediate intubation in all cases

Airway obstruction is a critical situation that requires prompt recognition and intervention. The Heimlich maneuver, also known as abdominal thrusts, is specifically recommended for dealing with choking in patients who have a complete airway obstruction due to a foreign body. Performing this maneuver can effectively generate enough pressure to expel the object blocking the airway, allowing the patient to breathe again. In cases of partial obstruction or when the patient is still able to cough or speak, it is advised to monitor the patient closely, providing supportive care while encouraging them to continue coughing effectively. The Heimlich maneuver is not appropriate for patients who can still breathe adequately, nor is it useful in cases of airway obstruction caused by other factors, such as swelling or anaphylaxis, where different interventions would be necessary. Using sedatives is not a suitable approach for managing airway obstructions, as it could potentially exacerbate the situation by further relaxing the airway muscles or reducing the patient's ability to respond to the obstruction. Immediate intubation is also not recommended in all cases of airway obstruction, as this is an invasive procedure that may not be indicated if the obstruction can be resolved through less invasive means. Nasal steroids are typically used to reduce inflammation in chronic respiratory conditions and are not effective for immediate management of

SAMPLE

9. A patient may present with wheezing and difficulty breathing due to which possible diagnosis?

A. Pneumonia

B. Anxiety attack

C. Asthma exacerbation

D. Heart failure

Wheezing and difficulty breathing are hallmark signs of bronchial constriction and airway obstruction, commonly associated with asthma exacerbation. In asthma, the airways become inflamed, leading to narrowed passages which can trigger wheezing due to the turbulent airflow when a patient exhales. This condition can fluctuate greatly and may present acutely, making it a primary diagnosis when observing these respiratory symptoms. Other potential diagnoses, while they can also cause breathing difficulties, typically present with different primary symptoms or mechanisms. For example, pneumonia often leads to respiratory distress but is usually accompanied by productive cough, fever, and chest pain, rather than the wheezing characteristic of asthma. Anxiety attacks can cause shortness of breath, yet they typically do not produce wheezing; the breathlessness is often more related to hyperventilation. Lastly, heart failure can result in dyspnea due to fluid overload in the lungs, but it usually presents with crackles or rales on auscultation, rather than wheezing. Therefore, the combination of wheezing and difficulty breathing strongly points toward an asthma exacerbation as the most plausible diagnosis in this scenario.

10. How does AMLS suggest managing a patient with suspected cardiac emergency?

A. Focusing exclusively on oxygen administration

B. Ensuring airway, breathing, and circulation; obtaining an ECG; and administering appropriate medication

C. Offering emotional support and reassurance

D. Prioritizing transport to the hospital over initial management

The recommendation to ensure airway, breathing, and circulation; obtain an ECG; and administer appropriate medication is comprehensive and addresses the critical components of managing a patient with a suspected cardiac emergency. Early stabilization of these fundamental aspects is crucial to prevent further deterioration of the patient's condition. Airway management is the first step in patient assessment, as it is vital to ensure that the patient can breathe adequately. Adequate breathing is necessary to maintain oxygenation and prevent hypoxia. Circulation management involves assessing the patient's pulse and blood pressure and recognizing signs of shock or decreased perfusion, which could indicate a severe cardiac event. Obtaining an ECG is essential in cardiac emergencies, as it helps identify dysrhythmias, ischemia, or infarction that may guide subsequent treatment. Administering appropriate medication based on the findings can alleviate symptoms, stabilize the heart, and improve the overall condition of the patient. While other options may encompass good practices in general emergency care, focusing solely on one aspect such as oxygen administration or providing reassurance does not adequately address the multifaceted nature of cardiac emergencies. Similarly, prioritizing transport over initial management may delay critical interventions that can be life-saving. This holistic approach creates a greater chance for positive outcomes in patients experiencing cardiac issues.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://advancedmedicallifesupport.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE