AHA BLS FOR HEALTHCARE PROVIDERS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ahablsforhealthcareproviders.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What action should you take if a child is unresponsive and not breathing?

A. Begin first aid and wait for help

B. Monitor until they wake up

C. Begin CPR immediately and call for emergency help

D. Check for a pulse and then act

2. What is the initial step of the C-A-B sequence in CPR?

A. Airway

B. Breathing

C. Chest compression

D. Defibrillation

3. When should adult AED pads be used on a child or infant?

A. When there are no child pads available

B. When the child is unconscious

C. When the child is under 5 years old

D. When pediatric pads are too large

4. What is the recommended compression to ventilation ratio during CPR for adults?

A. 30:2

B. 15:2

C. 20:1

D. 10:1

5. Which of the following best describes effective chest compressions?

A. Soft and slow

B. Deep and fast

C. Rapid and rhythmic

D. Consistent and smooth

6. In what situation should you not attempt to use CPR on a victim?

A. If the victim has a strong pulse

B. If the victim is responsive

C. If the environment poses a danger

D. If you suspect a spinal injury

SAMPLE

7. How often should breaths be delivered to adults who are unresponsive and not breathing?

A. Every 3-5 seconds

B. Every 5-6 seconds

C. Every 10 seconds

D. Once per minute

8. What is one of the signs that a victim is experiencing a cardiac arrest?

A. The victim is breathing normally

B. The victim is alert and talking

C. The victim is unresponsive and not breathing

D. The victim is coughing continuously

9. When is it appropriate to stop CPR?

A. When the rescuer is too exhausted to continue

B. When emergency responders arrive, the patient shows signs of life, or you are too exhausted to continue

C. When the patient starts to cough

D. When 5 minutes have elapsed since CPR began

10. What should be the initial action when encountering an unresponsive adult?

A. Start CPR immediately

B. Call for emergency assistance

C. Provide rescue breaths

D. Check for a pulse

SAMPLE

Answers

SAMPLE

1. C
2. C
3. A
4. A
5. D
6. C
7. B
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What action should you take if a child is unresponsive and not breathing?

A. Begin first aid and wait for help

B. Monitor until they wake up

C. Begin CPR immediately and call for emergency help

D. Check for a pulse and then act

When a child is unresponsive and not breathing, the most critical action to take is to begin CPR immediately and call for emergency help. This response is essential because an unresponsive individual who is not breathing is likely experiencing a life-threatening emergency, such as cardiac arrest. Starting CPR provides critical blood flow to the brain and vital organs, increasing the chances of survival. Time is of the essence in such situations, and initiating CPR as quickly as possible is crucial for improving outcomes. The process involves giving rescue breaths and chest compressions, which together help maintain blood circulation and oxygenation until professional medical assistance arrives. Calling for emergency help at the same time ensures that additional medical resources are on the way, as CPR alone is not enough to resolve the underlying issue but is vital for sustaining life until further intervention can occur. This combined approach maximizes the child's chances of recovery. In this scenario, monitoring the child until they wake up or merely starting first aid while waiting for help would not effectively address the urgent need for immediate intervention. Checking for a pulse might delay the initiation of life-saving measures, which can be detrimental in a critical situation where every second counts. Therefore, the course of action that includes immediate CPR and notification for emergency services is the best

2. What is the initial step of the C-A-B sequence in CPR?

A. Airway

B. Breathing

C. Chest compression

D. Defibrillation

The correct initial step of the C-A-B sequence in CPR is chest compression. This step emphasizes the importance of providing effective and immediate chest compressions to initiate blood circulation in a cardiac arrest scenario. The focus on chest compressions first is based on the understanding that, in a situation where the heart has stopped pumping blood, the priority is to restore blood flow to vital organs, especially the brain. Chest compressions should be performed at a depth of approximately 2 to 2.4 inches for adults, at a rate of 100 to 120 compressions per minute. This method allows for the best chance of survival by maintaining blood flow to the heart and brain until advanced medical help can arrive or until the heart rhythm can be restored. Other steps in the C-A-B sequence, such as assessing airway and breathing, or defibrillation, come after ensuring that effective chest compressions are initiated. The change to C-A-B instead of the previous A-B-C sequence reflects updated guidelines that prioritize circulation first, improving outcomes in cases of cardiac arrest.

SAMPLE

3. When should adult AED pads be used on a child or infant?

A. When there are no child pads available

B. When the child is unconscious

C. When the child is under 5 years old

D. When pediatric pads are too large

Using adult AED pads on a child or infant is appropriate when there are no pediatric pads available. This guideline ensures that you can still deliver defibrillation if necessary. Adult pads are designed to deliver the appropriate level of energy for older children and adults but can also be used in an emergency situation when pediatric pads are not accessible. It's important to note that while adult pads can be used, they may not be ideal due to the larger size and energy dose they deliver. Therefore, if pediatric pads are available, they should be used, as they are specifically designed to cater to infants and children's smaller bodies. However, in emergencies, prioritizing the use of any available pads on a patient in need of defibrillation is crucial, hence the importance of this choice.

4. What is the recommended compression to ventilation ratio during CPR for adults?

A. 30:2

B. 15:2

C. 20:1

D. 10:1

The recommended compression to ventilation ratio during CPR for adults is 30:2, which means that for every 30 chest compressions given, 2 rescue breaths should be administered. This ratio is established based on research that indicates this balance maximizes the potential for successful resuscitation. The focus on delivering high-quality chest compressions is crucial, as they help maintain blood flow to vital organs during cardiac arrest. The ratio allows for a significant number of compressions in a timely manner, helping to optimize the chances of restoring a viable heartbeat and circulation while also providing adequate ventilation. In adult CPR, the primary goal is to provide effective compressions that maintain circulation, while the ventilations ensure oxygenation. The 30:2 ratio strikes a balance that improves outcomes in adult resuscitation scenarios, as it allows rescuers to maintain a continuous focus on compressions, which is critical to the overall effectiveness of CPR.

5. Which of the following best describes effective chest compressions?

A. Soft and slow

B. Deep and fast

C. Rapid and rhythmic

D. Consistent and smooth

Effective chest compressions during cardiopulmonary resuscitation (CPR) are characterized by being consistent and smooth. This means that compressions should be delivered at a steady rate and with a consistent depth, ensuring that the heart receives sufficient blood flow to maintain circulation. The goal is to create a reliable rhythm that allows for adequate perfusion of vital organs. Consistency in the depth of compressions also ensures that each compression is sufficient to generate the necessary pressures needed to circulate blood effectively. Smoothness helps maintain a rhythm that is easier for healthcare providers to sustain over time and makes it more effective in generating blood flow. In contrast, while being deep, fast, or rhythmic can have merits in specific contexts, focusing on consistency and smoothness encompasses the essential aspects of delivering high-quality CPR. Each of these components contributes to maintaining proper blood flow, but without the element of consistency, the effectiveness of each compression can be compromised.

SAMPLE

6. In what situation should you not attempt to use CPR on a victim?

A. If the victim has a strong pulse

B. If the victim is responsive

C. If the environment poses a danger

D. If you suspect a spinal injury

In a situation where the environment poses a danger, it is essential to prioritize your safety before attempting to perform CPR on a victim. If there are hazards such as fire, electrical wires, or any other life-threatening conditions, focusing on personal safety is crucial. Attempting to provide assistance in a hazardous scenario not only puts the rescuer at risk but could also exacerbate the situation for the victim if the rescuer becomes incapacitated. Therefore, ensuring that the environment is safe allows a rescuer to effectively help the victim once the area is secure. Other factors, such as a strong pulse or responsiveness, indicate that CPR may not be necessary at that moment. Moreover, a spinal injury requires careful handling but does not inherently prevent the performance of CPR if the situation allows for it. By prioritizing safety in dangerous environments, effective care can occur without undue risk to the rescuer.

7. How often should breaths be delivered to adults who are unresponsive and not breathing?

A. Every 3-5 seconds

B. Every 5-6 seconds

C. Every 10 seconds

D. Once per minute

Breaths should be delivered to adults who are unresponsive and not breathing every 5-6 seconds. This frequency aligns with the guidelines provided by the American Heart Association (AHA) for delivering rescue breaths in a situation where the individual is experiencing a respiratory arrest. Delivering breaths at this rate ensures that the victim receives adequate ventilation without overwhelming their lungs or causing complications such as hyperventilation. The recommended timing helps to maintain oxygenation while allowing enough time for both ventilation and the body's natural processes. In this context, the other frequencies do not align with AHA guidelines, making them less suitable in a real emergency scenario. For instance, delivering breaths every 3-5 seconds may be too frequent, risking over-ventilation, whereas the options of every 10 seconds or once per minute would not provide sufficient breaths to adequately oxygenate the individual's blood during critical moments when response time is essential.

8. What is one of the signs that a victim is experiencing a cardiac arrest?

A. The victim is breathing normally

B. The victim is alert and talking

C. The victim is unresponsive and not breathing

D. The victim is coughing continuously

A victim experiencing cardiac arrest will typically show signs such as being unresponsive and not breathing. During cardiac arrest, the heart stops pumping blood effectively, leading to a lack of oxygenated blood reaching the brain and other vital organs. This results in the person losing consciousness and the absence of normal breathing or any effective breathing patterns. In such situations, immediate action is required to assess and provide CPR, along with activating emergency medical services. The presence of normal breathing, alertness, or continuous coughing indicates that the person is likely not in cardiac arrest and may be experiencing different medical conditions. Recognizing the specific signs of cardiac arrest is crucial for initiating timely and appropriate life-saving interventions.

EXAMPLE

9. When is it appropriate to stop CPR?

A. When the rescuer is too exhausted to continue

B. When emergency responders arrive, the patient shows signs of life, or you are too exhausted to continue

C. When the patient starts to cough

D. When 5 minutes have elapsed since CPR began

Stopping CPR is appropriate in specific situations that indicate the need for further medical intervention or that the patient may be recovering. The correct answer encompasses multiple scenarios: when emergency responders arrive, when the patient shows signs of life, or when the rescuer becomes too exhausted to continue. Emergency responders arriving means that trained professionals can take over and provide advanced care, which is crucial for the patient's survival. Observing signs of life, such as effective breathing, movement, or consciousness, indicates that the person may have resumed normal physiological functions and does not require CPR. Finally, while personal exhaustion can prompt the cessation of CPR, it's vital for rescuers to ensure that someone else is ready to take over responsibilities to maintain the patient's care. This comprehensive approach reflects a balance between the continued need for CPR and the health of the responder, ensuring that care remains consistent and effective.

10. What should be the initial action when encountering an unresponsive adult?

A. Start CPR immediately

B. Call for emergency assistance

C. Provide rescue breaths

D. Check for a pulse

When encountering an unresponsive adult, the initial action should be to call for emergency assistance. This step is crucial because, in a medical emergency, timely intervention can significantly improve outcomes. By activating the emergency response system, you ensure that help is on the way while you assess the situation further. It also allows you to focus on providing care to the patient without delay. After ensuring emergency services have been contacted, you would then proceed to check for responsiveness and breathing. If the individual is not breathing or only gasping, you would begin cardiopulmonary resuscitation (CPR) as the next vital step. Starting CPR immediately without calling for help first can lead to delays in advanced care, which is why the initial action should prioritize alerting emergency services.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ahablsforhealthcareproviders.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE