AIR METHODS CRITICAL CARE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://airmethodscriticalcare.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which condition is an indication for FFP transfusion?

A. Acute Blood Loss

B. Burns

C. Chronic Pain

D. Asthma

2. Which of the following rhythms is characterized by QRS complexes that are the same shape, size, and direction?

A. Monomorphic ventricular tachycardia

B. Ventricular fibrillation

C. Junctional escape rhythm

D. Idioventricular rhythm

3. What is the typical range for magnesium levels in mg/dL?

A. 1.0-1.5

B. 1.5-2.0

C. 1.7-2.2

D. 2.2-2.6

4. Which lead in an EKG primarily looks at the apex of the heart?

A. Lead I

B. Lead II

C. Lead III

D. Lead V6

5. What is the normal platelet count range in thousands per microliter?

A. 100-300

B. 150-450

C. 200-500

D. 250-600

6. When is the RhoGAM shot administered to the mother?

A. Before delivery only

B. After the first delivery or miscarriage

C. Once a month during pregnancy

D. Only after a blood transfusion

SAMPLE

7. Which component of a basic metabolic panel measures kidney function?

A. Sodium

B. Chloride

C. Creatinine

D. Potassium

8. Calcium administration in pediatric patients should be considered only for which condition?

A. Hypercalcemia

B. Known/suspected hypocalcemia

C. Hyperkalemia

D. Severe dehydration

9. In elevated ICP management, what should be monitored while administering Mannitol?

A. Urine output only

B. Serum osmolality

C. Blood pressure

D. Electrolyte levels

10. What are the characteristics of tension pneumothorax?

A. Tracheal deviation toward affected side

B. Decreased breath sounds and hyperresonance

C. Increased central venous pressure

D. Bradycardia and hypotension

SAMPLE

Answers

SAMPLE

1. B
2. A
3. C
4. B
5. B
6. B
7. C
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which condition is an indication for FFP transfusion?

A. Acute Blood Loss

B. Burns

C. Chronic Pain

D. Asthma

Fresh frozen plasma (FFP) transfusion is indicated in conditions where there is a need for replacement of clotting factors, particularly in patients who are at risk of bleeding due to coagulopathy or when specific clotting factors are deficient. In the context of burns, especially extensive third-degree burns, there can be significant fluid and protein loss, including clotting factors. The administration of FFP can help restore these factors and improve clotting, thus managing the risk of bleeding and supporting overall hemostatic function in critically injured patients. Acute blood loss generally requires packed red blood cells (PRBCs) for volume and hemoglobin replacement, rather than FFP unless there is concurrent coagulopathy. Chronic pain and asthma are not conditions that typically involve deficiencies in clotting factors, making FFP transfusion unnecessary in those cases.

2. Which of the following rhythms is characterized by QRS complexes that are the same shape, size, and direction?

A. Monomorphic ventricular tachycardia

B. Ventricular fibrillation

C. Junctional escape rhythm

D. Idioventricular rhythm

Monomorphic ventricular tachycardia is defined by its consistent appearance of QRS complexes that are identical in shape, size, and direction. This uniformity helps to distinguish it from other types of arrhythmias, where the QRS complexes may vary in morphology. In monomorphic ventricular tachycardia, the origin of the electrical impulses is from a single ectopic focus within the ventricles, leading to these uniform complexes. This can result in rapid heart rates, typically over 100 beats per minute, but the regularity and uniformity give healthcare providers a clear picture of the underlying rhythm. Recognizing this rhythm is crucial in the clinical setting as it often necessitates immediate treatment, especially if the patient is experiencing hemodynamic instability. Understanding the distinct nature of monomorphic ventricular tachycardia helps differentiate it from other rhythms such as ventricular fibrillation, where there is chaotic electrical activity leading to varied QRS complexes without any discernible pattern. Other rhythms like junctional escape rhythms and idioventricular rhythms also exhibit different characteristics, such as differing QRS morphologies or slower rates respectively.

3. What is the typical range for magnesium levels in mg/dL?

A. 1.0-1.5

B. 1.5-2.0

C. 1.7-2.2

D. 2.2-2.6

The typical range for magnesium levels in the body is crucial for various physiological functions, including nerve conduction, muscle contraction, and enzyme activity. The accepted normal range for serum magnesium levels is generally 1.7 to 2.2 mg/dL. This range reflects the concentration of magnesium that is considered optimal for maintaining metabolic processes and overall health. Magnesium plays a vital role in many cellular functions and processes, and levels outside the normal range can lead to various clinical conditions. Levels lower than this range may indicate magnesium deficiency, which can lead to neuromuscular excitation, while levels above this range may suggest hypermagnesemia, often associated with renal insufficiency or excessive supplementation. Given this context, the range of 1.7 to 2.2 mg/dL not only aligns with established clinical guidelines but also highlights the importance of maintaining magnesium homeostasis for proper bodily function.

SAMPLE

4. Which lead in an EKG primarily looks at the apex of the heart?

A. Lead I

B. Lead II

C. Lead III

D. Lead V6

Lead II is particularly useful for examining the electrical activity of the heart as it primarily reflects the path of depolarization towards the apex. This lead is positioned between the right arm and the left leg, which provides a diagonal view of the heart's electrical activity. Because Lead II is oriented in a way that aligns with the heart's natural electrical conduction path, it effectively captures the overall activity of the heart as it moves towards the apex. This is significant in clinical practice, as observing the apex can help in identifying arrhythmias and other cardiac conditions. In contrast, Leads I and III offer different perspectives; Lead I looks at the heart from a horizontal plane (left to right), whereas Lead III views the heart from the right arm to the left leg in a slightly different angle. Lead V6, on the other hand, is a precordial lead that looks at the heart from the left lateral aspect, providing valuable information about the hearts' surface but is not as indicative of the apex as Lead II.

5. What is the normal platelet count range in thousands per microliter?

A. 100-300

B. 150-450

C. 200-500

D. 250-600

The normal platelet count range for an individual is typically between 150,000 and 450,000 platelets per microliter of blood, which is often expressed as 150-450 in thousands per microliter. This range is crucial in clinical practice because platelets play a vital role in coagulation and wound healing. A count below the lower limit could suggest thrombocytopenia, which can lead to increased bleeding risks, while a count above the upper limit could indicate thrombocytosis, which may increase the risk of thromboembolic events. Understanding this range is essential for the assessment and management of patients in critical care settings. The other ranges provided are either too low or too high, which would not align with established clinical guidelines.

6. When is the RhoGAM shot administered to the mother?

A. Before delivery only

B. After the first delivery or miscarriage

C. Once a month during pregnancy

D. Only after a blood transfusion

The RhoGAM shot is administered to a mother, particularly those who are Rh-negative, to prevent Rh immunization, which can occur if an Rh-negative mother is exposed to Rh-positive blood. The primary purpose of administering RhoGAM is to protect future pregnancies from complications that may arise due to Rh incompatibility. This shot is typically given after the first delivery if the newborn is Rh-positive, and it is also indicated after any miscarriage or abortion involving Rh-positive blood. Administering the shot in these scenarios helps ensure that the mother's immune system does not produce antibodies against Rh-positive blood cells, which could affect future pregnancies. Given this context, the choice of administering the RhoGAM shot after the first delivery or miscarriage aligns with established medical guidelines for preventing Rh sensitization. It's crucial for healthcare providers to assess the Rh status of both the mother and the newborn and administer the RhoGAM shot as necessary to avoid complications in subsequent pregnancies.

SAMPLE

7. Which component of a basic metabolic panel measures kidney function?

A. Sodium

B. Chloride

C. Creatinine

D. Potassium

The component of a basic metabolic panel that specifically measures kidney function is creatinine. Creatinine is a waste product produced from muscle metabolism and is typically filtered out of the bloodstream by the kidneys. When kidney function is impaired, the levels of creatinine in the blood increase, making it a key indicator of renal health. Measuring creatinine helps healthcare professionals evaluate how well the kidneys are working and can indicate the presence of kidney disease or dysfunction. Higher levels may suggest that the kidneys are not filtering blood effectively, thus prompting further investigation or management of kidney health. Other components of the basic metabolic panel, such as sodium, chloride, and potassium, play essential roles in various physiological processes, including fluid balance and nerve function, but they are not direct indicators of kidney function.

8. Calcium administration in pediatric patients should be considered only for which condition?

A. Hypercalcemia

B. Known/suspected hypocalcemia

C. Hyperkalemia

D. Severe dehydration

Calcium administration in pediatric patients is primarily indicated for known or suspected hypocalcemia. Hypocalcemia can lead to various symptoms, including muscle spasms, seizures, and cardiac issues. In acute situations, such as in critically ill pediatric patients, addressing low calcium levels is vital to prevent severe complications. When hypocalcemia is suspected, immediate assessment and intervention are crucial, as calcium plays a critical role in many physiological processes, including muscle contraction, nerve conduction, and cardiac function. Administering calcium can restore normal cellular function and stabilize the patient. In contrast, conditions like hypercalcemia involve elevated calcium levels, which would make further calcium administration inappropriate. Hyperkalemia requires treatment focusing on lowering potassium levels rather than increasing calcium. Severe dehydration, while serious, does not directly relate to calcium levels and is managed through fluid replacement and electrolyte balance rather than calcium administration. Therefore, recognizing hypocalcemia as a basis for calcium administration highlights the importance of addressing specific electrolyte imbalances in pediatric care.

SAMPLE

9. In elevated ICP management, what should be monitored while administering Mannitol?

A. Urine output only

B. Serum osmolality

C. Blood pressure

D. Electrolyte levels

Mannitol is an osmotic diuretic commonly used in the management of elevated intracranial pressure (ICP). When administering Mannitol, monitoring serum osmolality is essential because this medication works by drawing fluid out of the brain tissue and into the bloodstream, leading to increased serum osmolality. An increase in serum osmolality can help reduce cerebral edema and lower ICP. However, if serum osmolality becomes too high, it can result in adverse effects such as renal failure, electrolyte imbalances, and shifts in fluid balance. Therefore, it is crucial to keep serum osmolality within a safe range to prevent potential complications associated with high osmolarity, including cellular dehydration or osmotic demyelination. While monitoring urine output, blood pressure, and electrolyte levels is important during the treatment, the most critical parameter directly reflecting the effects of Mannitol and guiding its safe administration is serum osmolality. This focused monitoring helps ensure the effectiveness of the treatment while minimizing risks.

10. What are the characteristics of tension pneumothorax?

A. Tracheal deviation toward affected side

B. Decreased breath sounds and hyperresonance

C. Increased central venous pressure

D. Bradycardia and hypotension

Tension pneumothorax is characterized by the accumulation of air in the pleural space that results in increased pressure, leading to significant physiological changes. One of the hallmark features is the presence of decreased breath sounds on the affected side due to the lung being compressed by the trapped air. Additionally, hyperresonance is noted upon percussion of the chest in the area of the pneumothorax. This occurs because the presence of air in a confined space creates a sound that is more resonant than the normal lung or fluid-filled thoracic cavity. Increased central venous pressure can also occur in tension pneumothorax due to the compression of blood vessels returning to the heart, but it is not one of the primary characteristics used to identify the condition at first presentation. Similarly, while bradycardia and hypotension may result from severe cardiorespiratory compromise in a tension pneumothorax, they are more secondary manifestations rather than defining characteristics. The primary assessment findings remain decreased breath sounds and hyperresonance on examination of the affected side.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://airmethodscriticalcare.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE