AMBULANCE VICTORIA CPGS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ambulancevictoriacpgs.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. During the assessment of major trauma, how important is timely triaging to trauma care?

A. It is irrelevant as long as the patient is stable

B. It is critical for improving patient outcomes

C. It is not as important as patient comfort

D. It can be delayed without consequences

2. What is a key characteristic of blunt force trauma?

A. It involves penetrating injuries

B. It results from external impact without breaking the skin

C. It is limited to soft tissue injuries

D. It primarily affects children

3. What should be monitored in a patient showing signs of shock?

A. Brain activity and fat reserves

B. Skin condition and vital signs

C. Appetite and hydration level

D. Size of the pupils and environment

4. What is the recommended depth of chest compressions for adults during CPR?

A. At least 2 cm (0.8 inches) deep

B. At least 3 cm (1.2 inches) deep

C. At least 4 cm (1.6 inches) deep

D. At least 5 cm (2 inches) deep

5. What vital sign indicates a red flag if heart rate exceeds which number?

A. 100

B. 110

C. 120

D. 130

SAMPLE

6. What typically signifies acute respiratory failure that may require an ETT?

A. Increased oxygen saturation

B. Glasgow Coma Scale (GCS) under 10

C. Low blood sugar levels

D. Stable respiratory rate

7. What does the airway assessment mnemonic AIRWAYS include?

A. Airway, Intervention, Recovery, Wait, Assess, Yell for help, Send for assistance

B. Airway, Illness, Rescue, Wait, Assess, Yell, Send for help

C. Airway, Infection, Rescue, Wait, Assist, Yell, Sense

D. Airway, Imaging, Rescue, Wait, Alert, Seek, Support

8. What technique should be used for a patient hyperventilating due to anxiety?

A. Encourage paced breathing

B. Advise skipping meals

C. Increase physical activity

D. Introduce meditation directly

9. Which of the following is NOT considered a reversible cause of PEA?

A. Tension pneumothorax

B. Upper airway obstruction

C. Severe dehydration

D. Anaphylaxis

10. What is one common intervention in managing patients with respiratory distress?

A. Administering pain relievers

B. Providing supplemental oxygen

C. Giving fluids intravenously

D. Performing CPR

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. D
5. C
6. B
7. B
8. A
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. During the assessment of major trauma, how important is timely triaging to trauma care?

A. It is irrelevant as long as the patient is stable

B. It is critical for improving patient outcomes

C. It is not as important as patient comfort

D. It can be delayed without consequences

Timely triaging in trauma care is crucial because it significantly impacts patient outcomes. In cases of major trauma, the immediate assessment and prioritization of injuries can lead to life-saving interventions. Rapid identification of the severity of trauma allows healthcare providers to allocate resources effectively, ensuring that the most critically injured patients receive immediate care. When triaging is performed promptly, medical personnel can quickly determine the necessary level of intervention, such as advanced imaging, surgical procedures, or transfer to a specialized trauma center. This swift action can make the difference between life and death, especially in situations where time is of the essence, like major hemorrhages or head injuries. Timely triaging also facilitates communication within emergency management systems, helping to coordinate responses among multiple agencies and ensuring that the right personnel and equipment are available at the scene or hospital. This organized approach helps streamline care and reduces the risk of complications, ultimately leading to improved recovery rates and better overall patient outcomes.

2. What is a key characteristic of blunt force trauma?

A. It involves penetrating injuries

B. It results from external impact without breaking the skin

C. It is limited to soft tissue injuries

D. It primarily affects children

Blunt force trauma is characterized by external impact that does not create a break in the skin. This type of injury is often a result of collisions, falls, or strikes from blunt objects, which can lead to various forms of injury such as contusions, abrasions, and fractures. The primary characteristic that defines blunt force trauma is that it occurs without penetrating the skin; instead, the damage is generally internal and may affect underlying tissues and organs. This distinction is important because it differentiates blunt force injuries from penetrating injuries, where an object pierces the skin and opens a wound. Additionally, blunt force trauma is not limited to soft tissue injuries, as it can also result in fractures of bones and damage to internal organs. While blunt force trauma can occur in any population, including children, it is not exclusive to them, making that characteristic not universally applicable.

SAMPLE

3. What should be monitored in a patient showing signs of shock?

A. Brain activity and fat reserves

B. Skin condition and vital signs

C. Appetite and hydration level

D. Size of the pupils and environment

In a patient exhibiting signs of shock, it is essential to monitor skin condition and vital signs. Shock can lead to inadequate perfusion of tissues, which may present clinically as pallor or cyanosis of the skin, coolness, and diaphoresis, indicating the body's response to stressed circulatory status. Monitoring vital signs, including heart rate, blood pressure, respiratory rate, and oxygen saturation, is critical in assessing the severity of shock. These parameters help determine if the patient is stabilizing, worsening, or responding to treatment. Changes in heart rate and blood pressure can provide insights into the patient's hemodynamic status, allowing for timely intervention. Other options mention aspects such as brain activity, appetite, hydration, and pupil size, which, while they may hold some relevance in different contexts, do not focus directly on the immediate, critical parameters necessary to assess and manage a patient in shock. Hence, concentrating on skin condition and vital signs provides the most pertinent information in such a scenario.

4. What is the recommended depth of chest compressions for adults during CPR?

A. At least 2 cm (0.8 inches) deep

B. At least 3 cm (1.2 inches) deep

C. At least 4 cm (1.6 inches) deep

D. At least 5 cm (2 inches) deep

The recommended depth of chest compressions for adults during CPR is at least 5 cm (2 inches) deep. This depth is crucial for effectively compressing the chest and ensuring sufficient blood flow to vital organs, particularly the heart and brain, during cardiac arrest. Compressions that are too shallow may not generate enough pressure to circulate blood effectively, which can significantly reduce the chances of survival. The guideline to compress at least 5 cm is based on extensive research and evidence demonstrating that deeper compressions improve outcomes in cases of adult cardiac arrest. Proper technique, including ensuring the correct depth, is vital for the success of CPR interventions.

5. What vital sign indicates a red flag if heart rate exceeds which number?

A. 100

B. 110

C. 120

D. 130

The indication that a heart rate exceeding a certain threshold represents a red flag relates to how elevated heart rates can signify potential physiological distress or underlying issues requiring further assessment. A heart rate above 120 beats per minute is often recognized as tachycardia in adults, which can be alarming and may indicate stress on the cardiovascular system, dehydration, fever, pain, or other medical conditions. When heart rates exceed this level, especially in a clinical context, it prompts emergency responders to investigate further, taking into account other vital signs and patient history. While lower rates may still indicate concern depending on the individual and context, the threshold of 120 is commonly acknowledged in clinical guidelines as a point where further action is typically warranted. This makes it a vital sign not only to monitor but also to respond to, thereby classifying it as a "red flag" in patient assessment.

EXAMPLE

6. What typically signifies acute respiratory failure that may require an ETT?

A. Increased oxygen saturation

B. Glasgow Coma Scale (GCS) under 10

C. Low blood sugar levels

D. Stable respiratory rate

Acute respiratory failure is a critical condition marked by the inability of the respiratory system to maintain adequate gas exchange, often leading to life-threatening consequences. A Glasgow Coma Scale (GCS) score under 10 is particularly significant in this context. A decreased GCS indicates a compromised level of consciousness, often associated with neurological impairment or significant underlying medical issues. This impairment can lead to reduced respiratory drive or ineffective airway protection, increasing the risk of respiratory failure. In such scenarios, intubation via an endotracheal tube (ETT) may be necessary to secure the airway and ensure adequate ventilation and oxygenation. In contrast, increased oxygen saturation, low blood sugar levels, and a stable respiratory rate do not directly signify acute respiratory failure. Increased oxygen saturation typically suggests adequate oxygenation, low blood sugar levels can indicate metabolic issues, and a stable respiratory rate alone does not reflect the overall function of ventilation or the adequacy of gas exchange. Therefore, observing a GCS under 10 indicates a critical need for intervention such as intubation to prevent further deterioration of the patient's respiratory status.

7. What does the airway assessment mnemonic AIRWAYS include?

A. Airway, Intervention, Recovery, Wait, Assess, Yell for help, Send for assistance

B. Airway, Illness, Rescue, Wait, Assess, Yell, Send for help

C. Airway, Infection, Rescue, Wait, Assist, Yell, Sense

D. Airway, Imaging, Rescue, Wait, Alert, Seek, Support

The correct response includes components that accurately reflect the necessary steps in an airway assessment process in emergency care. The elements of "Airway, Illness, Rescue, Wait, Assess, Yell, Send for help" outline a structured approach to evaluating and managing a patient's airway. - Starting with "Airway," it emphasizes the critical first step of ensuring that the patient's airway is not obstructed and is open for breathing. - "Illness" helps responders consider the underlying condition affecting the patient that could influence airway management or other interventions. - "Rescue" implies taking immediate action if the airway is compromised, suggesting a need for urgent interventions. - "Wait" can indicate the importance of assessing the situation before acting impulsively. - "Assess" stresses the ongoing evaluation of the patient's airway status and overall condition. - "Yell" brings attention to reaching out for assistance, recognizing that effective teamwork is vital in emergency scenarios. - Finally, "Send for help" implies alerting additional resources when facing a challenging situation, reinforcing the importance of having adequate support in emergency cases. This mnemonic effectively captures the essential aspects of airway assessment and management, guiding responders in making informed decisions during critical situations. Each component plays a role in promoting patient safety

SAMPLE

8. What technique should be used for a patient hyperventilating due to anxiety?

A. Encourage paced breathing

B. Advise skipping meals

C. Increase physical activity

D. Introduce meditation directly

Encouraging paced breathing is an effective technique for a patient who is hyperventilating due to anxiety. This approach helps to regulate the breathing pattern, which can become rapid and shallow during anxiety episodes, leading to hyperventilation. By guiding the patient to slow their breathing, they can restore a more normal respiratory rate, reduce feelings of dizziness or lightheadedness, and help alleviate the panic associated with hyperventilation. Paced breathing typically involves taking slow, deep breaths, often counting to a certain number for inhalation and exhalation. This method not only provides a physical means to control breathing but also acts as a mental focus, diverting attention away from anxiety-provoking thoughts. The other options, while potentially beneficial in other contexts, do not directly address the immediate need to reduce hyperventilation. Skipping meals could lead to low blood sugar, which might worsen anxiety symptoms. Increasing physical activity may not be advisable during an acute episode of hyperventilation as it could exacerbate the situation. Introducing meditation techniques, while useful in the long term for managing anxiety, may not be practical to implement in the moment of acute hyperventilation where immediate action is necessary.

9. Which of the following is NOT considered a reversible cause of PEA?

A. Tension pneumothorax

B. Upper airway obstruction

C. Severe dehydration

D. Anaphylaxis

Severe dehydration is not typically classified as a reversible cause of pulseless electrical activity (PEA) because it does not directly lead to a mechanical or electrical obstruction of the heart or major vessels that can be immediately corrected in an emergency setting. PEA arises from multiple potential causes, many of which involve acute and potentially reversible conditions that result in inadequate cardiac output or ineffective circulation. In contrast, conditions like tension pneumothorax and upper airway obstruction involve immediate physical problems that can be quickly remedied, restoring normal circulatory function. Tension pneumothorax compresses the heart and great vessels, and when it is relieved, circulation can often be restored. Similarly, resolving upper airway obstruction can restore adequate oxygenation and blood flow. Anaphylaxis also represents a reversible cause, as it can lead to vasodilation and severe allergic reactions that can be treated effectively with appropriate interventions, such as epinephrine. While severe dehydration affects overall fluid balance and can impact circulation, it is not an acute event that can be addressed on the spot like the other conditions mentioned. In an emergency context, the immediate focus is usually on addressing causes that directly impede hemodynamic stability.

SAMPLE

10. What is one common intervention in managing patients with respiratory distress?

A. Administering pain relievers

B. Providing supplemental oxygen

C. Giving fluids intravenously

D. Performing CPR

Providing supplemental oxygen is a common and essential intervention in managing patients with respiratory distress. When a patient experiences respiratory difficulty, their body may not be receiving adequate amounts of oxygen, leading to a decrease in oxygen saturation levels in the blood. Supplemental oxygen helps to increase the delivery of oxygen to the lungs and subsequently to the bloodstream, improving the patient's overall oxygenation and helping to alleviate symptoms of distress. By administering oxygen, healthcare providers can enhance the availability of oxygen for cellular metabolism and help stabilize the patient until further assessment and treatment can be conducted. This intervention is particularly crucial in conditions such as chronic obstructive pulmonary disease (COPD), asthma exacerbations, pneumonia, and acute respiratory distress syndrome (ARDS), among others. In contrast, while administering pain relievers is important in managing pain, it does not directly address respiratory distress. Providing intravenous fluids may be necessary in specific situations involving dehydration or shock, but it is not a primary intervention for respiratory problems. Lastly, performing CPR is reserved for cases of cardiac arrest and does not generally apply to cases of respiratory distress unless the patient has stopped breathing or has no pulse.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ambulancevictoriacpgs.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE