AMERICAN ASSOCIATION OF CRITICAL-CARE NURSES (AACN) CERTIFICATION PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://aacn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **If the anticipated time frame for not using the gut is less than 2 weeks, which method of nutritional support should be used?**

   A. Central vein nutrition

   B. Peripheral vein (PPN)

   C. Enteral feeding

   D. Tube feeding

2. **Which type of shock is characterized by decreased blood volume?**

   A. Cardiogenic shock

   B. Distributive shock

   C. Hypovolemic shock

   D. Obstructive shock

3. **What action should be taken initially in the management of suspected tracheoinnominate artery fistula?**

   A. Perform a bronchoscopy

   B. Apply manual pressure

   C. Start intravenous fluid resuscitation

   D. Intubate the patient

4. **Which of the following is a category of infectious causes of fever?**

   A. Autoimmune reactions

   B. Bacterial infections

   C. Inflammatory processes

   D. Drug side effects

5. **What does the NIHSS Stroke Scale assess?**

   A. Heart rate variability

   B. Neurological status in stroke patients

   C. Pain levels in patients

   D. Muscle strength

SAMPLE

6. Which characteristic is most common in refeeding syndrome?

A. Hyperkalemia

B. Hypophosphatemia

C. Vitamin A deficiency

D. Severe dehydration

7. What is Immune Reconstitution Inflammatory Syndrome (IRIS) characterized by?

A. Improvement in immune function without any complications

B. Paradoxical worsening of pre-existing infections or malignancies

C. Immediate immune response within days of ART initiation

D. Reduction in viral load without ART

8. What syndrome is characterized by prolonged exposure to cortisol due to a pituitary tumor?

A. Adrenal insufficiency

B. Cushing's syndrome

C. Addison's disease

D. Hyperaldosteronism

9. Signs of left-sided heart failure primarily affect which part of the body?

A. Abdomen

B. Lungs

C. Limbs

D. Heart

10. What is the normal range for magnesium levels?

A. 1.0-2.0

B. 1.5-2.5

C. 2.0-3.0

D. 1.5-3.5

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. B
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. If the anticipated time frame for not using the gut is less than 2 weeks, which method of nutritional support should be used?

A. Central vein nutrition

B. Peripheral vein (PPN)

C. Enteral feeding

D. Tube feeding

When the anticipated time frame for not using the gut is less than 2 weeks, peripheral vein nutrition is often considered appropriate because it allows for providing necessary nutrients while the gastrointestinal tract is not being utilized. Peripheral parenteral nutrition (PPN) can be administered via a peripheral intravenous line, which is less invasive than central venous access and is suitable for short-term nutritional support. PPN is typically used for patients who do not require complete bowel rest but still need additional caloric intake or if they cannot consume adequate nutrition orally. It is important to note that for anticipated periods longer than 2 weeks, central vein nutrition might be preferred due to its ability to provide a more concentrated form of nutrition. In this context of short-term nutritional support, enteral feeding options such as tube feeding could also be used. However, the specific guidance suggests using a method that is less intensive and more suited for brief durations, which aligns with the benefits PPN offers for limited time frames. Therefore, in cases where the gut can be utilized for feeding over an anticipated short period, peripheral vein nutrition is the method of choice.

2. Which type of shock is characterized by decreased blood volume?

A. Cardiogenic shock

B. Distributive shock

C. Hypovolemic shock

D. Obstructive shock

Hypovolemic shock is characterized by a significant decrease in blood volume, which can result from a variety of causes such as severe bleeding, dehydration, or fluid loss due to conditions like burns or diarrhea. When the blood volume decreases, the heart does not have enough fluid to pump effectively, leading to a drop in blood pressure and inadequate perfusion of tissues and organs. In this type of shock, the body attempts to compensate for the low blood volume through mechanisms such as increasing heart rate and constricting blood vessels, but if the volume loss is significant and not corrected, it can lead to shock symptoms including confusion, lethargy, and even organ failure. The other types of shock involve different underlying mechanisms. For instance, cardiogenic shock is primarily due to the heart's inability to pump blood effectively, distributive shock involves abnormal distribution of blood flow often due to vasodilation (like in septic shock), and obstructive shock is caused by a physical obstruction of blood flow (like in tension pneumothorax or pulmonary embolism). Each of these has distinct causes and physiological responses, which underscores why hypovolemic shock is specifically linked to decreased blood volume.

SAMPLE

3. What action should be taken initially in the management of suspected tracheoinnominate artery fistula?

A. Perform a bronchoscopy

B. Apply manual pressure

C. Start intravenous fluid resuscitation

D. Intubate the patient

The initial action in the management of a suspected tracheoinnominate artery fistula is to perform a bronchoscopy. This diagnostic procedure allows the healthcare provider to visualize the airway directly and assess the extent of the injury or abnormality. In the case of a tracheoinnominate artery fistula, where there is a risk of exsanguination due to a connection between the trachea and the innominate artery, bronchoscopy can help identify the source of the bleeding and other related complications. While other interventions may be necessary after assessment, such as controlling bleeding or maintaining airway patency, recognizing the fistula through visual inspection provides critical information that dictates subsequent management. Addressing the situation through direct visualization is essential in guiding further treatment steps, such as potential surgical intervention or advanced airway management.

4. Which of the following is a category of infectious causes of fever?

A. Autoimmune reactions

B. Bacterial infections

C. Inflammatory processes

D. Drug side effects

Bacterial infections are indeed a category of infectious causes of fever. Fever is a common response of the immune system to infection, and bacterial infections are among the most prevalent causes of fever in patients. When bacteria invade the body, they can trigger an immune response that leads to the release of pyrogens, substances that cause the hypothalamus in the brain to raise the body's temperature. This process helps the body fight off infection by enhancing immune function and creating an environment less favorable for bacterial growth. In contrast, autoimmune reactions and inflammatory processes are not infectious causes, as they typically involve the body's own immune response attacking its tissues or responding to non-infectious stimuli. Drug side effects can also lead to fever, but they are not classified as infectious causes since they arise from medication rather than an infectious pathogen. Therefore, bacterial infections represent a clear and direct linkage to the physiological mechanisms that cause fever through the body's response to actual infectious agents.

SAMPLE

5. What does the NIHSS Stroke Scale assess?

A. Heart rate variability

B. Neurological status in stroke patients

C. Pain levels in patients

D. Muscle strength

The NIHSS (National Institutes of Health Stroke Scale) is specifically designed to quantify the impairment caused by a stroke and assess the neurological status of stroke patients. It evaluates various aspects of neurological function, including consciousness, motor function, and language skills, through a series of standardized tests. Each component of the NIHSS helps in determining the severity of a stroke and aids in guiding treatment decisions and predicting outcomes. The other options are not applicable in this context. For instance, while heart rate variability is an important measure in cardiology, it does not pertain to stroke assessment. Similarly, pain levels are not the primary focus of the NIHSS; the scale is more concerned with neurological deficits rather than subjective pain experiences. Although muscle strength is a component of the neurological assessment, it is only one part of the broader evaluation process that the NIHSS encompasses. Thus, the most accurate description of what the NIHSS Stroke Scale assesses is indeed the neurological status in stroke patients.

6. Which characteristic is most common in refeeding syndrome?

A. Hyperkalemia

B. Hypophosphatemia

C. Vitamin A deficiency

D. Severe dehydration

Hypophosphatemia is indeed the most common characteristic of refeeding syndrome. This condition often occurs when feeding is restarted in individuals who have been malnourished or in a state of starvation. When carbohydrates are reintroduced, insulin secretion increases, driving phosphate from the serum into the cells to be utilized in the production of ATP, which can lead to a significant decrease in serum phosphate levels. This phosphorus shift can result in a range of symptoms, including muscle weakness, respiratory failure, and cardiac dysrhythmias. Monitoring and managing phosphate levels is crucial during the refeeding period to prevent complications associated with refeeding syndrome. In contrast, while hyperkalemia can occur in other contexts, it is not a defining characteristic of refeeding syndrome. Vitamin A deficiency is unrelated to the immediate metabolic changes seen during refeeding. Severe dehydration may be present in some patients but is also not a hallmark of this syndrome. Therefore, hypophosphatemia stands out as the primary concern when addressing refeeding syndrome in clinical practice.

SAMPLE

7. What is Immune Reconstitution Inflammatory Syndrome (IRIS) characterized by?

A. Improvement in immune function without any complications

B. Paradoxical worsening of pre-existing infections or malignancies

C. Immediate immune response within days of ART initiation

D. Reduction in viral load without ART

Immune Reconstitution Inflammatory Syndrome (IRIS) is characterized by a paradoxical worsening of pre-existing infections or malignancies following the initiation of antiretroviral therapy (ART) in individuals with immune suppression, particularly those with HIV/AIDS. As the immune system begins to recover due to the effective treatment of the underlying HIV infection, it becomes more capable of mounting an inflammatory response against previously recognized pathogens or malignancies. This heightened immune response can lead to clinical deterioration, as the body reacts strongly to infections that were previously subdued or asymptomatic. This phenomenon is particularly notable in patients with significant immune recovery, which occurs when the CD4 T-cell counts increase significantly. Although ART improves overall immune function, the inflammatory responses can sometimes lead to complications that exacerbate the clinical condition of the patient, thus allowing for the paradoxical worsening characteristic of IRIS. In contrast, the other options describe scenarios that do not align with the definitions or symptoms of IRIS. For example, statements about improved immune function without complications or immediate immune responses occurring solely within days do not capture the delayed and often complicated nature of IRIS. Similarly, the notion of a reduction in viral load without ART does not pertain to the context of immune reconstitution but rather suggests an

8. What syndrome is characterized by prolonged exposure to cortisol due to a pituitary tumor?

A. Adrenal insufficiency

B. Cushing's syndrome

C. Addison's disease

D. Hyperaldosteronism

Cushing's syndrome is defined by the body being exposed to high levels of cortisol for an extended period. This condition can result from various causes, but when attributed specifically to a pituitary tumor, it is often referred to as Cushing's disease, a subtype of Cushing's syndrome. In this scenario, the pituitary gland produces an excess amount of adrenocorticotropic hormone (ACTH) due to the tumor, which in turn stimulates the adrenal glands to secrete more cortisol. The chronic elevation in cortisol levels can lead to a range of symptoms and complications, including weight gain, hypertension, diabetes, and other metabolic changes. Understanding this condition's pathology—particularly how the pituitary's role is central to the overproduction of ACTH and subsequent cortisol increase—is crucial for recognizing and diagnosing Cushing's syndrome effectively. Other conditions listed do not align with the given context. Adrenal insufficiency refers to insufficient hormone production by the adrenal glands, which is not characterized by prolonged cortisol exposure. Addison's disease is also an adrenal insufficiency scenario caused by destruction of the adrenal glands, leading to insufficient cortisol levels. Hyperaldosteronism involves excess production of aldosterone, which affects sodium and water balance rather than cortisol levels. Therefore, C

SAMPLE

9. Signs of left-sided heart failure primarily affect which part of the body?

A. Abdomen

B. Lungs

C. Limbs

D. Heart

The signs of left-sided heart failure are primarily associated with the lungs because when the left ventricle fails to effectively pump blood out to the body, it causes a backlog of blood in the pulmonary circulation. This increased pressure in the pulmonary veins leads to fluid leakage from the blood vessels into the lung tissues, resulting in pulmonary congestion and symptoms such as shortness of breath, wheezing, and cough. In the context of left-sided heart failure, while there may be systemic effects and symptoms that can involve different areas of the body, the impact on the lungs is the most pronounced and immediate due to the physiological changes that occur as a result of the heart's impaired function. Therefore, recognizing the respiratory symptoms as central to left-sided heart failure is essential for diagnosis and management.

10. What is the normal range for magnesium levels?

A. 1.0-2.0

B. 1.5-2.5

C. 2.0-3.0

D. 1.5-3.5

The normal range for magnesium levels in the bloodstream is typically between 1.5 mg/dL and 2.5 mg/dL. This range is crucial because magnesium plays a vital role in various physiological functions, including muscle and nerve function, blood glucose control, and blood pressure regulation. Values below 1.5 mg/dL may indicate hypomagnesemia, which can lead to muscle cramps, fatigue, and arrhythmias, while values above 2.5 mg/dL suggest hypermagnesemia, which can cause symptoms such as nausea, vomiting, and cardiac disturbances. Understanding the correct range not only assists healthcare professionals in diagnosing and managing potential imbalances but also emphasizes the importance of magnesium in overall health. Regular monitoring of magnesium levels is essential in patients at risk for electrolytic imbalances, particularly those with gastrointestinal or renal issues, as they may have altered absorption or excretion of this key mineral.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://aacn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE