
AMERICAN BOARD OF DENTAL EXAMINERS (ADEX) DENTAL HYGIENE LICENSING PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adex.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Who is most commonly affected by cementoma (periapical cemental dysplasia)?

A. Caucasian men

B. African American women

C. Hispanic adults

D. Asian children

2. What clinical feature is associated with Warthin tumor?

A. Usually occurs in young females

B. Occurs in the tail of the parotid gland

C. Commonly found in the anterior mandible

D. Typically debulked due to high malignancy

3. What is a common imaging appearance of fibrous dysplasia?

A. Ground glass appearance

B. Pear-shaped radiolucency

C. Heart-shaped radiolucency

D. Multilocular radiolucency

4. What is the recommended dosage of Amoxicillin for antibiotic prophylaxis?

A. 2 g 30 minutes before treatment

B. 2 g 1 hour before treatment

C. 1 g 1 hour before treatment

D. 1 g at the time of treatment

5. What is the minimum distance required between anterior-posterior palatal straps in a removable partial denture?

A. 10 mm

B. 12 mm

C. 15 mm

D. 20 mm

SAMPLE

6. What is a common feature of benign cementoblastoma?

A. Tooth is vital

B. Lesion is always painful

C. Common in females over 30

D. Typically requires no treatment

7. Mikulicz disease is primarily associated with which condition?

A. Hashimoto's thyroiditis

B. Stevens-Johnson syndrome

C. Sjogren's syndrome

D. Lupus erythematosus

8. Which feature is indicative of Acute Myelogenous Leukemia (AML) on a blood smear?

A. Immature white blood cells

B. Auer rods

C. Philadelphia chromosome

D. Increased red blood cells

9. What is the characteristic rash associated with German measles (Rubella)?

A. Petechial rash

B. Maculopapular rash

C. Vesicular rash

D. Urticarial rash

10. Which term describes the condition where normal bone structure is replaced with fibrous tissue?

A. Fibrous dysplasia

B. Gardner's syndrome

C. Calcifying epithelial odontogenic tumor

D. Dentigerous cyst

SAMPLE

Answers

SAMPLE

1. B
2. B
3. A
4. B
5. C
6. A
7. C
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Who is most commonly affected by cementoma (periapical cemental dysplasia)?

A. Caucasian men

B. African American women

C. Hispanic adults

D. Asian children

Cementoma, also known as periapical cemental dysplasia, predominantly affects a specific demographic, with research indicating that it most commonly occurs in African American women. This condition is characterized by the abnormal growth of cementum-like tissue at the apex of the teeth, typically associated with the roots of mandibular anterior teeth. Understanding the composition of the affected demographic reveals that hormonal influences and genetic predispositions may play a significant role in the prevalence of this condition among African American women. The clinical manifestations may vary, but early stages often present asymptomatically, making it essential for dental professionals to recognize the typical presentations in this group for effective diagnosis and management. In contrast, other demographics, such as Caucasian men, Hispanic adults, and Asian children, show significantly lower incidences of cementoma, highlighting the unique susceptibility of African American women to this particular dental condition. Detecting and monitoring cementoma in this population is crucial for preventing potential complications, including issues related to tooth vitality and adjacent structures.

2. What clinical feature is associated with Warthin tumor?

A. Usually occurs in young females

B. Occurs in the tail of the parotid gland

C. Commonly found in the anterior mandible

D. Typically debulked due to high malignancy

The clinical feature associated with Warthin tumor is its occurrence in the tail of the parotid gland. Warthin tumor, also known as papillary cystadenoma lymphomatosum, is a benign salivary gland tumor that most commonly arises in the parotid gland, particularly in the tail region. This location is significant as it distinguishes Warthin tumors from other types of salivary gland tumors, which may commonly occur in different areas of the salivary glands, including anterior regions or other glands entirely. In contrast, the other options describe characteristics that do not align with typical presentations of Warthin tumors. They do not occur primarily in young females, nor are they commonly found in the anterior mandible, which is more characteristic of other types of tumors. Furthermore, Warthin tumors are benign and are not typically debulked due to malignancy concerns, as they have a low rate of malignant transformation. Understanding these details reinforces the significance of the specific location of Warthin tumors within the parotid gland.

SAMPLE

3. What is a common imaging appearance of fibrous dysplasia?

A. Ground glass appearance

B. Pear-shaped radiolucency

C. Heart-shaped radiolucency

D. Multilocular radiolucency

Fibrous dysplasia is characterized by a unique imaging appearance that is often described as a "ground glass" texture. This appearance results from the abnormal proliferation of fibrous tissue and the replacement of normal bone, leading to a radiologic feature that resembles frosted glass. This texture can be observed on radiographs as a diffuse radiopacity, which may not have sharp borders. The ground glass appearance is significant because it helps differentiate fibrous dysplasia from other lesions or conditions that may be seen in similar areas. For instance, the other choices describe different radiolucency patterns that are characteristic of various pathologies. Some conditions may exhibit multilocular radiolucency or other specific shapes, but the defining ground glass characteristic of fibrous dysplasia is what sets it apart in diagnostic imaging. Recognizing this appearance is crucial for dental hygienists and other dental professionals when evaluating radiographs and determining the appropriate course of action or referral.

4. What is the recommended dosage of Amoxicillin for antibiotic prophylaxis?

A. 2 g 30 minutes before treatment

B. 2 g 1 hour before treatment

C. 1 g 1 hour before treatment

D. 1 g at the time of treatment

The recommended dosage of Amoxicillin for antibiotic prophylaxis is indeed 2 g taken 1 hour before treatment. This dosage is specifically indicated for individuals who are at risk of infective endocarditis during dental procedures. Administering the antibiotic 1 hour prior ensures that adequate serum levels are achieved in time to provide effective prophylaxis against potential bacterial dissemination that could lead to serious complications. The rationale behind taking Amoxicillin an hour before treatment instead of immediately before or at the time of treatment lies in pharmacokinetics, which studies how drugs are absorbed, distributed, metabolized, and excreted in the body. By allowing that one-hour window, the medication can be absorbed effectively, thereby maximizing its concentration in the bloodstream when the dental procedure takes place. This strategic timing is essential for those who have specific underlying health conditions that place them at higher risk for infective endocarditis, as it aids in preventing bacteria from settling in the heart tissues during invasive procedures.

5. What is the minimum distance required between anterior-posterior palatal straps in a removable partial denture?

A. 10 mm

B. 12 mm

C. 15 mm

D. 20 mm

The minimum distance required between anterior-posterior palatal straps in a removable partial denture is based on providing adequate support and stability to the denture while also ensuring comfort for the patient. A distance of 15 mm allows for effective distribution of forces during function, helping to prevent movement of the partial denture that could cause discomfort or affect its performance. This measurement also considers the anatomical features of the palate, ensuring that the straps do not interfere with the patient's normal oral functions, such as speech and swallowing. Opting for a distance shorter than 15 mm could lead to increased chances of denture dislodgement and discomfort for the wearer, while longer distances, while they may provide more support, could unnecessarily complicate the design and aesthetics of the denture. Thus, the correct choice of 15 mm balances these factors effectively.

SAMPLE

6. What is a common feature of benign cementoblastoma?

A. Tooth is vital

B. Lesion is always painful

C. Common in females over 30

D. Typically requires no treatment

A common feature of benign cementoblastoma is that the associated tooth is vital. This condition involves the formation of a tumor around the roots of a tooth where cementum, the hard tissue covering the roots, is abnormally produced. The vital status of the tooth is significant because, in benign cementoblastoma, the tooth remains alive and responds to stimuli, which contrasts with other dental lesions that can result in necrosis or non-vitality of the tooth. In benign cementoblastoma, the lesion typically does not present with pain unless there is secondary inflammation or complications. This makes the option regarding the lesion being always painful inaccurate. Additionally, while benign cementoblastoma can occur in individuals across various demographics, it is not specifically common in females over 30; it can affect a broader range of ages and genders. Lastly, treatment is often indicated to remove the lesion and prevent further complications, thus the idea that it typically requires no treatment does not hold true for most cases.

7. Mikulicz disease is primarily associated with which condition?

A. Hashimoto's thyroiditis

B. Stevens-Johnson syndrome

C. Sjogren's syndrome

D. Lupus erythematosus

Mikulicz disease is primarily associated with Sjogren's syndrome, which is characterized by dry eyes and dry mouth due to the immune system attacking moisture-producing glands. In the context of Sjogren's syndrome, Mikulicz disease refers to the swelling of the parotid and lacrimal glands, leading to the classic presentation of symptoms associated with this autoimmune condition. The condition can exist as an isolated entity or as part of a broader autoimmune process, often characterized by other systemic effects. Understanding the association with Sjogren's syndrome is crucial for diagnosis and management, as patients with this condition have an elevated risk of developing other autoimmune disorders and require comprehensive care to address their systemic symptoms and improve their quality of life. This context emphasizes the importance of recognizing the specific autoimmune mechanisms at play in Mikulicz disease.

8. Which feature is indicative of Acute Myelogenous Leukemia (AML) on a blood smear?

A. Immature white blood cells

B. Auer rods

C. Philadelphia chromosome

D. Increased red blood cells

The presence of Auer rods on a blood smear is a distinctive feature indicative of Acute Myelogenous Leukemia (AML). Auer rods are cytoplasmic inclusions formed by the fusion of granular precursors in myeloid cells, and their identification is significant in diagnosing this type of leukemia. In the context of AML, Auer rods can be seen in myeloblasts and indicate the myeloid lineage of the cancerous cells, which is a key factor in differentiating AML from other types of leukemia. While immature white blood cells might also be present in blood smears of patients with leukemia, the specific identification of Auer rods provides a more definitive association with AML. The Philadelphia chromosome is primarily linked to Chronic Myelogenous Leukemia (CML) rather than AML. Lastly, the increase in red blood cells is not characteristic of AML; in fact, patients may present with anemia due to the disrupting effects of leukemia on normal hematopoiesis. The presence of Auer rods serves as a vital diagnostic criterion in the classification and management of myeloid leukemias.

SAMPLE

9. What is the characteristic rash associated with German measles (Rubella)?

A. Petechial rash

B. Maculopapular rash

C. Vesicular rash

D. Urticarial rash

The characteristic rash associated with German measles, or rubella, is indeed a maculopapular rash. This type of rash typically presents as a combination of flat (macular) red spots and raised bumps (papular). It often begins on the face and then spreads to the rest of the body, usually starting from the hairline and moving downward, which is consistent with classic presentations of rubella. The maculopapular nature of the rash is important as it helps distinguish rubella from other viral infections that may present differently. For instance, the petechial rash is made up of small, pinpoint red or purple spots and is often associated with different conditions like thrombocytopenia or specific viral infections like meningococcemia. Vesicular rashes feature small, fluid-filled blisters and are often linked to infections like chickenpox or herpes simplex. Urticarial rashes, also known as hives, consist of raised welts on the skin and can be triggered by allergic reactions or infections but do not align with the typical presentation of rubella. Understanding the specific features of the maculopapular rash associated with rubella aids in identifying the infection correctly and differentiating it from other viral illnesses that may present

10. Which term describes the condition where normal bone structure is replaced with fibrous tissue?

A. Fibrous dysplasia

B. Gardner's syndrome

C. Calcifying epithelial odontogenic tumor

D. Dentigerous cyst

The term that describes the condition where normal bone structure is replaced with fibrous tissue is fibrous dysplasia. In this disorder, bone is progressively replaced by fibrous tissue, resulting in deformities and potential pain. This condition can affect one bone (monostotic fibrous dysplasia) or multiple bones (polyostotic fibrous dysplasia) and can lead to structural weaknesses in the affected bones. The other options refer to different conditions: Gardner's syndrome is a genetic disorder characterized by the development of various tumors and dental abnormalities, including osteomas and soft tissue tumors, rather than the replacement of bone with fibrous tissue. The calcifying epithelial odontogenic tumor, also known as a Pindborg tumor, is a benign odontogenic neoplasm characterized by the presence of calcifying epithelial cells, but it does not involve the replacement of bone. A dentigerous cyst is an odontogenic cyst associated with the crown of an unerupted tooth, which may cause localized bone expansion but does not lead to the replacement of normal bone structure with fibrous tissue.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adex.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE