
AMERICAN BOARD OF OBSTETRICS & GYNECOLOGY (ABOG) QUALIFYING PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://abogqualifying.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Which of the following is a contraindication for breastfeeding in the United States?**

   A. Asthma

   B. Active untreated tuberculosis

   C. Current use of antibiotics

   D. Parent with seasonal allergies

2. **What is the best time for determining chorionicity and amnionicity in twins?**

   A. 4-6 weeks

   B. 6-8 weeks

   C. 10-12 weeks

   D. 14-16 weeks

3. **What potential risk is associated with albumin administration?**

   A. Coagulation abnormalities

   B. Anaphylactic reactions

   C. Volume overload

   D. Hypoglycemia

4. **What is the recommended frequency of ultrasound for monitoring estimated fetal weight (EFW) in cases where EFW is between the 3rd to 10th percentile?**

   A. Every week

   B. Every 2 weeks

   C. Every 3-4 weeks

   D. Every month

5. **What is the most effective preventive measure for autonomic dysreflexia during labor?**

   A. Positioning the patient upright

   B. Administration of oral antihypertensives

   C. Neuraxial anesthesia at T10

   D. Frequent monitoring of vital signs

SAMPLE

6. Which finding signifies ventriculomegaly in relation to NTD?

A. Lemon sign

B. Olive sign

C. Banana sign

D. Bladder sign

7. Which aspect of medical ethics addresses respect for patients' rights?

A. Beneficence

B. Justice

C. Non maleficence

D. Autonomy

8. At what age does ACOG recommend beginning thyroid screening in women with risk factors or symptoms?

A. 19 years old

B. 25 years old

C. 30 years old

D. 35 years old

9. Which quad screen results indicate high risk for Down syndrome?

A. High AFP and low estriol

B. Low AFP and high hCG

C. Low AFP and estriol, high hCG and inhibin A

D. Normal AFP and estriol, low hCG

10. What could potentially happen due to decreased preload from hypotension and inferior vena cava (IVC) compression in pregnancy?

A. Increased fetal movement

B. Cardiopulmonary arrest

C. Increase in maternal heart rate

D. Uterine hypertonicity

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. C
5. C
6. A
7. D
8. A
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. Which of the following is a contraindication for breastfeeding in the United States?

A. Asthma

B. Active untreated tuberculosis

C. Current use of antibiotics

D. Parent with seasonal allergies

Active untreated tuberculosis is a contraindication for breastfeeding because the bacteria responsible for tuberculosis, Mycobacterium tuberculosis, can be transmitted to the infant through breast milk if the mother is not correctly treated. Breastfeeding in such cases poses a significant risk to the infant, particularly because tuberculosis can lead to severe illness in young children. In the United States, guidelines emphasize that mothers with active untreated tuberculosis should be isolated and treated adequately before considering breastfeeding. Once the mother is on effective anti-tuberculosis therapy and has shown signs of improvement, breastfeeding can often be safely resumed, as the treatment reduces the risk of transmission. Asthma, current use of antibiotics, and parental seasonal allergies do not pose the same level of risk to breastfeeding. Asthma can generally be managed with medications that are safe during lactation, antibiotics are often safe for breastfeeding mothers, and allergies in a parent do not affect the safety of breastfeeding. Overall, understanding these distinctions is crucial for ensuring the health of both the mother and infant.

2. What is the best time for determining chorionicity and amnionicity in twins?

A. 4-6 weeks

B. 6-8 weeks

C. 10-12 weeks

D. 14-16 weeks

Determining chorionicity and amnionicity in twin pregnancies is best performed between 6 to 8 weeks gestation. At this stage, transvaginal ultrasound is particularly effective for visualizing the early intrauterine structures. The presence of early signs, such as the number of gestational sacs and the visualization of the intermembranous septum, allows for accurate identification of whether the twins are monochorionic or dichorionic. Before 6 weeks, it may be difficult to clearly delineate the separate sacs, especially in very early pregnancies where structures are still developing. After 8 weeks, while the distinctions might still be made, the risk of misclassifying amnionicity increases as the membranes and amniotic sacs become more prominent and potentially overlap. Hence, the 6-8 week window provides the most reliable timeframe for assessment of chorionicity and amnionicity in twin pregnancies, ensuring better management and monitoring throughout the pregnancy for potential complications related to chorionic and amniotic placentation.

SAMPLE

3. What potential risk is associated with albumin administration?

A. Coagulation abnormalities

B. Anaphylactic reactions

C. Volume overload

D. Hypoglycemia

Administering albumin carries a potential risk of volume overload. Albumin is a colloid that helps maintain oncotic pressure in the vasculature, and while it can be beneficial in certain clinical situations—such as treating hypoalbuminemia or providing resuscitation in cases of shock—it can also lead to an increase in intravascular volume. This is particularly a concern in patients with compromised cardiac function or those with a history of heart failure, as the additional fluid can exacerbate their condition, potentially leading to pulmonary edema or heart failure. The other options, while relevant in different contexts, do not represent the primary concern associated with albumin administration. For instance, coagulation abnormalities are more typically linked to factors like liver disease or certain coagulation disorders rather than albumin itself. Anaphylactic reactions, although possible with any protein administration, are relatively rare with albumin compared to other blood products. Lastly, hypoglycemia is not a recognized risk associated with albumin administration, as albumin does not significantly influence blood glucose levels. Understanding these factors helps underscore the importance of monitoring and assessing patient fluid status when using albumin in clinical settings.

4. What is the recommended frequency of ultrasound for monitoring estimated fetal weight (EFW) in cases where EFW is between the 3rd to 10th percentile?

A. Every week

B. Every 2 weeks

C. Every 3-4 weeks

D. Every month

When monitoring estimated fetal weight (EFW) in cases where the EFW falls between the 3rd to 10th percentile, the recommended frequency is every 3-4 weeks. This interval allows for adequate assessment of fetal growth while minimizing unnecessary interventions or maternal anxiety. Fetuses that are small for gestational age (SGA), particularly those below the 10th percentile, require careful observation due to potential complications associated with restricted growth. Monitoring every 3-4 weeks strikes a balance between ensuring close surveillance and allowing for sufficient time to assess changes in growth patterns without overwhelming the healthcare system or the mother. More frequent monitoring, like every week or every 2 weeks, may be reserved for fetuses showing concerning growth patterns or changes in health status, but for stable cases within the 3rd to 10th percentile, every 3-4 weeks is deemed appropriate. Monthly monitoring would be too infrequent to detect important changes in fetal growth, making that option less suitable.

SAMPLE

5. What is the most effective preventive measure for autonomic dysreflexia during labor?

A. Positioning the patient upright

B. Administration of oral antihypertensives

C. Neuraxial anesthesia at T10

D. Frequent monitoring of vital signs

The most effective preventive measure for autonomic dysreflexia during labor is the administration of neuraxial anesthesia at T10. Autonomic dysreflexia is a serious condition that can occur in individuals with spinal cord injuries above the T6 level, where noxious stimuli below the level of injury lead to an exaggerated sympathetic response. This can result in increased blood pressure, bradycardia, and other significant complications. Neuraxial anesthesia works by blocking the transmission of sympathetic nerves from the lower half of the body, effectively mitigating the noxious stimuli that trigger autonomic dysreflexia. By achieving sufficient neural blockade at the T10 level, which corresponds to the upper limit of labor pain transmission, the occurrence of autonomic dysreflexia is significantly reduced. This not only helps manage pain during labor but also decreases the risk of autonomic dysreflexia by preventing the sympathetic overload that occurs in response to painful stimuli. Other measures like positioning the patient upright may help alleviate symptoms once dysreflexia occurs but are not effective preventively. The administration of oral antihypertensives could help manage elevated blood pressure, but they do not address the underlying cause related to sympathetic activation. Lastly, while frequent monitoring of vital signs is important for early

6. Which finding signifies ventriculomegaly in relation to NTD?

A. Lemon sign

B. Olive sign

C. Banana sign

D. Bladder sign

The lemon sign is indicative of ventriculomegaly, particularly when considered alongside neural tube defects (NTDs). It is an ultrasound finding characterized by a distinctive shape of the fetal skull, which appears flattened in the frontal region and pointed at the occipital area. This configuration often occurs due to dysgenesis of the cerebellum and is commonly observed in conditions such as spina bifida. Ventriculomegaly itself refers to the enlargement of the brain's ventricles, which is often associated with NTDs. The presence of the lemon sign suggests underlying structural brain changes that may occur with NTDs and indicates that further evaluation is warranted to assess the extent of the neuroanatomical anomalies. In contrast, other signs, such as the olive sign and banana sign, relate to different aspects of fetal development or other specific anomalies, whereas the bladder sign does not hold relevance in identifying ventriculomegaly in this context. Understanding the implications of these ultrasound findings is crucial for proper diagnosis and management of suspected neural tube defects and associated complications.

SAMPLE

7. Which aspect of medical ethics addresses respect for patients' rights?

A. Beneficence

B. Justice

C. Non maleficence

D. Autonomy

The aspect of medical ethics that addresses respect for patients' rights is autonomy. Autonomy recognizes the patient's right to make informed decisions about their own healthcare and treatment options. It emphasizes the importance of allowing individuals to express their values, preferences, and wishes regarding their medical care. This principle is foundational in establishing a trusting relationship between patients and healthcare providers, as it empowers patients to actively participate in their healthcare decisions. Understanding autonomy also means recognizing that patients have the right to accept or refuse treatments based on their values and beliefs. This is crucial in fostering a respectful and ethically sound medical practice where patients feel acknowledged and validated in their choices. In contrast, beneficence involves the obligation to act in the best interest of the patient, promoting their well-being. While this principle is essential in ensuring quality care, it does not specifically highlight the importance of respecting patient rights as autonomy does. Justice pertains to fairness and equitable distribution of healthcare resources but does not directly address individual patient rights. Non-maleficence focuses on the principle of "do no harm," which is essential in medical ethics but also does not emphasize patient rights as clearly as autonomy does.

8. At what age does ACOG recommend beginning thyroid screening in women with risk factors or symptoms?

A. 19 years old

B. 25 years old

C. 30 years old

D. 35 years old

The American College of Obstetricians and Gynecologists (ACOG) recommends that women with risk factors or symptoms of thyroid disease begin screening starting at age 19. This recommendation is based on the understanding that while thyroid dysfunction can occur at any age, the prevalence of thyroid disorders tends to increase with age, and early identification in younger women, especially those with specific risk factors, can lead to timely management and improved outcomes. Risk factors for thyroid disease may include a personal or family history of thyroid disorders, autoimmune diseases, or previous treatment for thyroid issues. Starting screening at 19 allows for proactive monitoring and intervention if needed, which is crucial for women, particularly those who may become pregnant, as thyroid health plays a significant role in reproductive health and fetal development. Thus, screening at this age facilitates early detection and management of potential thyroid abnormalities.

SAMPLE

9. Which quad screen results indicate high risk for Down syndrome?

A. High AFP and low estriol

B. Low AFP and high hCG

C. Low AFP and estriol, high hCG and inhibin A

D. Normal AFP and estriol, low hCG

The quad screen is a maternal serum screening test that assesses the risk of certain conditions, including Down syndrome, during pregnancy. It measures four specific markers: alpha-fetoprotein (AFP), estriol, human chorionic gonadotropin (hCG), and inhibin A. In the case of Down syndrome, a typical pattern seen in the quad screen results is a combination of low AFP, low estriol, high hCG, and elevated inhibin A. This pattern reflects the way that placental and fetal metabolism changes in pregnancies affected by Down syndrome. Low levels of AFP and estriol suggest a potential developmental issue, while high levels of hCG and inhibin A further heighten the concern for trisomy 21 (Down syndrome). The elevations in hCG and inhibin A occur due to the placental alterations associated with the condition, making these hormone levels pertinent in evaluating risk. Understanding this pattern is crucial for interpreting quad screen results effectively and recognizing when further diagnostic testing, such as amniocentesis or chorionic villus sampling, may be warranted for a more definitive assessment.

10. What could potentially happen due to decreased preload from hypotension and inferior vena cava (IVC) compression in pregnancy?

A. Increased fetal movement

B. Cardiopulmonary arrest

C. Increase in maternal heart rate

D. Uterine hypertonicity

Decreased preload, which refers to the reduced volume of blood returning to the heart, can lead to hypotension and is particularly relevant in the context of pregnancy. When the inferior vena cava (IVC) is compressed, often due to the weight of the gravid uterus, it impedes venous return and subsequently can cause significant drops in maternal blood pressure. This chronic hypotension can compromise maternal and fetal circulation. If this condition escalates, it can lead to serious outcomes such as inadequate perfusion of vital organs, eventually culminating in cardiopulmonary arrest if not addressed promptly. Maternal collapse, in this case, is a life-threatening scenario where the heart ceases to function effectively due to prolonged low blood volume and pressure, leading to decreased cardiac output. In this context, increased fetal movement and uterine hypertonicity are not direct consequences of decreased preload but rather may occur under different circumstances. An increase in maternal heart rate can occur as a compensatory mechanism to try to maintain cardiac output despite the low preload. However, it is the potential for cardiopulmonary arrest that represents the most severe risk associated with significant hypotension and IVC compression during pregnancy.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://abogqualifying.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE