AMERICAN BOARD OF PATHOLOGY (ABPATH) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://abpath.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Which congenital anomaly of the spleen is most associated with IVC anomalies and congenital heart defects?**

   A. Polysplenia

   B. Asplenia

   C. Splenomegaly

   D. Splenectomy

2. **Which of the following best describes recurrent rami?**

   A. Arteries that connect larger branches

   B. Veins draining the testis

   C. Branches that originate from the mediastinum

   D. Veins supplying the epididymis

3. **What is located in the anterolateral part of the pancreatic head?**

   A. Gastroduodenal artery

   B. Superior mesenteric artery

   C. Pancreatic duct

   D. Inferior mesenteric artery

4. **What abnormality is indicated by a tardus parvus waveform?**

   A. Increased blood flow

   B. Stenosis

   C. Normal vascular response

   D. Vasodilation

5. **The Doppler waveform of the hepatic vein typically demonstrates which pattern?**

   A. Hepatofugal

   B. Hepatopetal

   C. Reversed flow

   D. Continuous flow

SAMPLE

6. Focal fatty sparing in the liver is typically characterized as what compared to surrounding tissue?

A. Hypoechoic

B. Hyperechoic

C. Isoechoic

D. Enhanced echogenicity

7. Which condition is known to cause enlarged hepatic veins?

A. Annular pancreas

B. Right heart failure

C. Acalculous cholelithiasis

D. Splenomegaly

8. Which of the following conditions is commonly associated with hyperthyroidism?

A. Hashimoto's thyroiditis

B. Graves' disease

C. Thyroiditis

D. Goiter

9. What is the cause of the type of flow associated with an AVF?

A. Aortic Dissection

B. Arteriovenous Fistula

C. Peripheral Vascular Disease

D. The presence of a thrombus

10. What type of medical device is implanted in the inferior vena cava to prevent emboli from traveling to the lungs?

A. Stent

B. IVC filter

C. Port-a-cath

D. Catheter

SAMPLE

Answers

SAMPLE

1. B
2. A
3. A
4. B
5. A
6. A
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which congenital anomaly of the spleen is most associated with IVC anomalies and congenital heart defects?

A. Polysplenia

B. Asplenia

C. Splenomegaly

D. Splenectomy

Asplenia, or the absence of the spleen, is notably associated with a range of congenital anomalies, particularly involving the inferior vena cava (IVC) and congenital heart defects. This association can be understood through the concept of heterotaxy syndrome, which includes asplenia and forms of bilateral left-sidedness. In patients with asplenia, the absence of the spleen often correlates with structural anomalies in the major venous and arterial systems, especially in the IVC. These patients frequently present with congenital heart defects, which result from improper embryonic development leading to the abnormal placement of organs and vessels. Such abnormalities create a distinct clinical picture characterized by various cardiovascular issues as well as venous anomalies. While polysplenia is also associated with heterotaxy, it is typically linked to a different set of structural anomalies, primarily involving the presence of multiple spleens rather than the absence of one. Splenomegaly refers to an enlargement of the spleen and does not have the same associations with IVC or congenital heart anomalies. Splenectomy, the surgical removal of the spleen, although it may seem relevant, is an acquired condition rather than a congenital anomaly. Understanding these details highlights why asplenia is the correct

2. Which of the following best describes recurrent rami?

A. Arteries that connect larger branches

B. Veins draining the testis

C. Branches that originate from the mediastinum

D. Veins supplying the epididymis

Recurrent rami refer to small branches that arise from larger nerves or vessels and then return to join another structure. In the context of anatomy, especially regarding the nervous system, these branches typically extend and reconnect to larger domains, creating functional loops or circuits. The description of recurrent rami as arteries that connect larger branches accurately reflects the physiological purpose of these structures. They serve to facilitate blood supply and ensure that areas supplied by the larger arteries receive adequate flow. This connection can be crucial for maintaining appropriate circulation, especially in complex vascular networks. The other options do not align with the definition of recurrent rami. For instance, veins draining the testis, branches from the mediastinum, and veins specifically supplying the epididymis are all focused on specific anatomical structures or specific functions. However, they do not embody the general connecting and looping aspect that is characteristic of recurrent rami in the context of circulation and innervation.

SAMPLE

3. What is located in the anterolateral part of the pancreatic head?

A. Gastroduodenal artery

B. Superior mesenteric artery

C. Pancreatic duct

D. Inferior mesenteric artery

The gastroduodenal artery is indeed located in the anterolateral part of the pancreatic head. This artery branches off from the common hepatic artery and plays a significant role in supplying blood to the stomach and the duodenum, as well as providing branches that contribute to the blood supply of the pancreas. Its proximity to the pancreatic head is crucial because the area requires adequate blood flow for the functioning of both the pancreas and neighboring structures. Understanding the anatomy of the pancreatic head is important not just in terms of blood supply, but also in relation to surrounding organs and their blood vessels. The superior mesenteric artery, for instance, is located posterior to the neck of the pancreas, not in the anterolateral region. The pancreatic duct runs through the pancreas but is not located in the anterolateral part of the head. The inferior mesenteric artery, on the other hand, supplies the distal colon and is situated far from the pancreatic head. This context reinforces the understanding of vascular relationships within the upper abdomen, particularly concerning the pancreas.

4. What abnormality is indicated by a tardus parvus waveform?

A. Increased blood flow

B. Stenosis

C. Normal vascular response

D. Vasodilation

A tardus parvus waveform is indicative of stenosis, which refers to the narrowing of a blood vessel. This type of waveform is characterized by a delayed (tardus) and low-amplitude (parvus) appearance. It typically occurs in situations where there is significant obstruction in the vascular pathway, leading to reduced blood flow velocity and altered hemodynamics. In normal situations, waveforms reflect a healthy flow of blood, usually demonstrating a rapid upstroke followed by a more pronounced dicrotic notch. However, stenosis causes turbulence and a decrease in the peak flow velocity. Hence, observing a tardus parvus waveform suggests that there is a significant narrowing in the artery upstream, which restricts blood flow and subsequently changes the flow pattern observed in doppler ultrasound studies. The other options do not appropriately characterize the tardus parvus waveform. Increased blood flow would typically produce a high-velocity waveform with sharp peaks rather than a delayed and diminished pattern. Normal vascular responses would show typical biphasic or triphasic waveforms that represent good distal vascular function. Vasodilation would also not be characterized by a tardus parvus waveform, as it generally results in increased blood flow and a different waveform appearance due to the widening of blood vessels

SAMPLE

5. The Doppler waveform of the hepatic vein typically demonstrates which pattern?

A. Hepatofugal

B. Hepatopetal

C. Reversed flow

D. Continuous flow

The Doppler waveform of the hepatic vein typically demonstrates a hepatofugal pattern, characterized by flow moving away from the liver. This is significant because the hepatic veins drain deoxygenated blood from the liver into the inferior vena cava. In a healthy individual, the expected Doppler waveform will show a triphasic pattern that reflects the influence of the heart's pulsations and the respiratory cycle. However, during diastole, the predominant flow is directed hepatofugally. This understanding is critical in assessing hepatic blood flow dynamics, especially when evaluating liver conditions or portal hypertension. Recognition of the hepatofugal flow pattern can also aid in various diagnostic evaluations pertaining to hepatic function and venous obstruction. The other options, while related to different aspects of blood flow or pathological conditions, do not accurately represent the typical hepatic vein Doppler waveform.

6. Focal fatty sparing in the liver is typically characterized as what compared to surrounding tissue?

A. Hypoechoic

B. Hyperechoic

C. Isoechoic

D. Enhanced echogenicity

Focal fatty sparing in the liver occurs when certain areas of the liver are less affected by fatty infiltration compared to the surrounding liver tissue. This phenomenon manifests as regions that appear less echogenic (hypoechoic) when viewed on ultrasound. The fat-containing areas typically exhibit increased echogenicity, while the spared areas retain a relatively lower echogenicity, making them appear darker on the ultrasound image against the brighter, fatty regions. Recognizing this contrast is essential in interpreting liver imaging correctly. Other terms, such as hyperechoic or isochoic, describe different echogenic characteristics but do not accurately represent focal fatty sparing in the context of surrounding liver tissue. Enhanced echogenicity would imply greater brightness, which does not apply to the hypoechoic areas standing out due to their lower fat content. Understanding these echogenic patterns is crucial for diagnosing various liver conditions, including steatosis and diseases leading to altered liver architecture.

SAMPLE

7. Which condition is known to cause enlarged hepatic veins?

A. Annular pancreas

B. Right heart failure

C. Acalculous cholelithiasis

D. Splenomegaly

Enlarged hepatic veins are often associated with increased pressure in the venous system, particularly due to conditions affecting the right side of the heart. Right heart failure leads to a backup of blood flow in the venous circulation, which can manifest as congestion in the hepatic veins. This congestion causes the veins to enlarge due to the increased volume and pressure of blood, resulting in a condition known as hepatic venous hypertension. In contrast, conditions such as annular pancreas, acalculous cholelithiasis, and splenomegaly do not directly impact the venous return or the pressure dynamics in the liver's venous system. While they may be associated with other gastrointestinal or systemic changes, they do not specifically lead to the enlargement of the hepatic veins in the same manner as right heart failure does. Therefore, the condition that most clearly correlates with the enlargement of the hepatic veins is right heart failure, due to its direct influence on venous pressure and blood flow dynamics.

8. Which of the following conditions is commonly associated with hyperthyroidism?

A. Hashimoto's thyroiditis

B. Graves' disease

C. Thyroiditis

D. Goiter

Graves' disease is the most common condition associated with hyperthyroidism. It is an autoimmune disorder that leads to the overproduction of thyroid hormones due to the stimulation of the thyroid gland by thyroid-stimulating immunoglobulins. This results in hypermetabolic symptoms like weight loss, heat intolerance, and anxiety, which are characteristic of hyperthyroidism. The other conditions listed do not primarily or commonly present with hyperthyroidism. Hashimoto's thyroiditis is typically associated with hypothyroidism because it leads to the gradual destruction of thyroid tissue. Thyroiditis, which includes various forms of thyroid inflammation, can lead to periods of hyperthyroidism but is not predominantly characterized by it. Goiter refers to the enlargement of the thyroid gland and can occur in both hyperthyroid and hypothyroid conditions, making it less specific to any one disorder. Therefore, Graves' disease is distinctly linked with hyperthyroidism due to its mechanism of thyroid hormone overproduction.

SAMPLE

9. What is the cause of the type of flow associated with an AVF?

A. Aortic Dissection

B. Arteriovenous Fistula

C. Peripheral Vascular Disease

D. The presence of a thrombus

An arteriovenous fistula (AVF) is a direct connection between an artery and a vein, bypassing the capillary bed. This unique anatomical configuration creates a type of blood flow that is characterized by high flow and low resistance. In an AVF, the high-pressure arterial blood flows directly into the venous system, leading to increased venous return and altered hemodynamics. The creation of an AVF can occur either surgically for purposes like hemodialysis or can develop pathologically, such as in cases of trauma. The resulting flow pattern is distinct from typical arterial or venous blood flow due to the unusual connection. This condition often results in a characteristic "thrill" or "bruit" that can be detected on examination, indicative of the turbulent flow generated at the connection point. In contrast, other conditions like aortic dissection or peripheral vascular disease involve different mechanisms that do not produce the same direct communication between arteries and veins that defines an AVF. The presence of a thrombus represents a blockage, which would further disrupt normal flow and does not facilitate the unique high-flow profile observed in an AVF. Therefore, the relationship of flow specifically linked to an arteriovenous fistula is the correct choice

10. What type of medical device is implanted in the inferior vena cava to prevent emboli from traveling to the lungs?

A. Stent

B. IVC filter

C. Port-a-cath

D. Catheter

The implanted device that is specifically designed to prevent emboli from traveling to the lungs is referred to as an IVC filter. The inferior vena cava (IVC) filter is a small medical device made of metal or plastic that is inserted into the inferior vena cava, the large vein that carries deoxygenated blood from the lower body to the heart. The primary function of an IVC filter is to capture and trap blood clots that may form in the deep veins of the legs and pelvis, which could otherwise travel to the lungs and cause a pulmonary embolism—an obstructive event that can be fatal. By positioning the filter in the IVC, healthcare providers can effectively reduce the risk of this life-threatening condition in patients who are at high risk, such as those with deep vein thrombosis or certain post-surgical patients. The other devices mentioned serve different purposes: stents are used to keep blood vessels open, port-a-caths are long-term intravenous access devices for medication delivery, and catheters are used for various purposes in medicine, such as draining fluids or administering medications. None of these have the specific function of preventing emboli from traveling to the lungs, which is the critical role of the IVC filter.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://abpath.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE