

AMERICAN BOARD OF PODIATRIC MEDICINE (ABPM) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



**SAMPLE STUDY GUIDE
WITH LIMITED QUESTIONS**

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://podiatricmedabpm.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Which of the following is NOT related to the duration of local anesthetic action?**
 - A. Protein binding
 - B. Onset time
 - C. Drug metabolism
 - D. Clinical application

- 2. What angle of measurement indicates coxa valga?**
 - A. Less than 90 degrees
 - B. Between 90 and 125 degrees
 - C. Greater than 125 degrees
 - D. Exactly 125 degrees

- 3. What does enthesitis refer to?**
 - A. Inflammation of joint capsules
 - B. Inflammation of muscles
 - C. Inflammation of tissue at muscle insertion on bone
 - D. Injury of ligaments

- 4. What structure is removed last in the lateral release procedure?**
 - A. Fibular sesamoid
 - B. Deep transverse intermetatarsal ligament
 - C. Plantar fascia
 - D. Flexor hallucis brevis

- 5. Which condition is represented by the O'Donoghue knee triad or unhappy triad?**
 - A. ACL, PCL, LCL
 - B. ACL, MCL, medial meniscus
 - C. ACL, MCL, lateral meniscus
 - D. ACL, LCL, medial meniscus

- 6. What is the treatment approach for Henoch-Schönlein purpura?**
- A. Immediate surgery
 - B. Self-limited course
 - C. Long-term antibiotic therapy
 - D. Daily corticosteroids
- 7. What is another name for Christmas tree disease?**
- A. Pityriasis rosea
 - B. Psoriasis vulgaris
 - C. Atopic dermatitis
 - D. Seborrheic dermatitis
- 8. Which antibiotic is specifically recommended for Gustillo Anderson type 1 fractures?**
- A. Amikacin
 - B. Ciprofloxacin
 - C. First or second-generation cephalosporin
 - D. Gentamicin
- 9. What complication is most commonly caused by lidocaine toxicity?**
- A. Hypertension
 - B. Bradycardia
 - C. Severe headache
 - D. Nausea
- 10. What is the chance of survival for a split thickness skin graft?**
- A. Low
 - B. High
 - C. Moderate
 - D. Variable

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. B
6. B
7. A
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which of the following is NOT related to the duration of local anesthetic action?

- A. Protein binding
- B. Onset time**
- C. Drug metabolism
- D. Clinical application

The duration of action of local anesthetics is influenced by several factors, including protein binding, drug metabolism, and clinical application. Protein binding is significant because highly bound drugs tend to have a longer duration of action due to a slower release into the systemic circulation. Drug metabolism also plays a critical role, as local anesthetics that are metabolized more quickly will have a shorter duration of action. Clinical application can impact the duration of anesthesia as well, including the choice of technique and the site of administration. For instance, certain injection techniques may prolong the effect of the anesthetic or alter its metabolism. Onset time, however, does not directly affect the duration of action. It refers to how quickly the anesthetic takes effect after administration rather than how long the effects will last, making it the correct answer in this scenario. Understanding these distinctions helps to clarify the pharmacological properties of local anesthetics.

2. What angle of measurement indicates coxa valga?

- A. Less than 90 degrees
- B. Between 90 and 125 degrees
- C. Greater than 125 degrees**
- D. Exactly 125 degrees

Coxa valga is characterized by an increased angle of femoral neck-troat, typically defined as greater than 125 degrees. This condition results from a structural alteration of the hip joint that leads to an outward positioning of the femur in relation to the hip, which can contribute to various functional issues such as gait abnormalities and increased risk of hip injury. In contrast, less than 90 degrees indicates coxa vara, which is a condition where the angle is decreased, leading to an inward position of the femur. An angle between 90 and 125 degrees is considered normal. Lastly, an angle that measures exactly 125 degrees falls within the normal range and does not indicate coxa valga. Thus, the precise measurement for identifying coxa valga is specifically defined as greater than 125 degrees.

3. What does enthesitis refer to?

- A. Inflammation of joint capsules
- B. Inflammation of muscles
- C. Inflammation of tissue at muscle insertion on bone**
- D. Injury of ligaments

Enthesitis specifically refers to the inflammation that occurs at the site where tendons, ligaments, and other connective tissues insert into the bone, known as the enthesis. This condition is often associated with various forms of spondyloarthritis and can lead to pain and discomfort at these insertion points. Recognizing the pathophysiology of enthesitis is crucial for understanding how it can affect overall mobility and function, especially in individuals with chronic inflammatory conditions. Understanding the impact of enthesitis is vital for effective diagnosis and management in podiatric medicine, as it can contribute to significant morbidity if left untreated.

4. What structure is removed last in the lateral release procedure?

- A. Fibular sesamoid
- B. Deep transverse intermetatarsal ligament**
- C. Plantar fascia
- D. Flexor hallucis brevis

In the lateral release procedure, the deep transverse intermetatarsal ligament is removed last because it plays a crucial role in stabilizing the metatarsal heads and maintaining the alignment of the forefoot. This ligament connects the second, third, and fourth metatarsals and helps to keep the metatarsal heads in place during normal locomotion. By removing it last, the surgeon ensures that the forefoot has structural support until the end of the procedure, minimizing the risk of inadvertently causing excessive destabilization of the metatarsal heads or compromising the arch of the foot. This is a careful consideration in surgical technique, as it helps to achieve the desired correction without creating additional complications or instability in the foot after the release is performed. Other structures, such as the fibular sesamoid, plantar fascia, and flexor hallucis brevis, have different roles in foot mechanics and are typically addressed earlier in the procedure, ensuring that the deep transverse intermetatarsal ligament's supportive role assists through the final steps of the release.

5. Which condition is represented by the O'Donoghue knee triad or unhappy triad?

- A. ACL, PCL, LCL
- B. ACL, MCL, medial meniscus**
- C. ACL, MCL, lateral meniscus
- D. ACL, LCL, medial meniscus

The O'Donoghue knee triad, commonly referred to as the unhappy triad, involves three specific knee injuries that occur simultaneously during a traumatic event, typically involving a forceful impact to the knee. The correct answer, which identifies this triad as ACL (anterior cruciate ligament), MCL (medial collateral ligament), and the medial meniscus, is significant for several reasons. Firstly, the anterior cruciate ligament is critical for knee stability and preventing anterior translation of the tibia relative to the femur. When injured, it often occurs with rotational forces that can also stress the other structures of the knee. Secondly, the medial collateral ligament is located on the inner part of the knee and provides stability against lateral forces. A simultaneous injury to the MCL is common in scenarios where there's a direct hit to the lateral aspect of the knee, causing a combination of stress on both the ACL and the MCL. Finally, the medial meniscus is the crescent-shaped cartilage that provides cushioning and stability within the knee joint. It is susceptible to tearing during injuries that affect the ACL and MCL due to its anatomical proximity and the rotational forces involved. This disorder commonly manifests in athletes, especially during sports involving sudden stops, jumping

6. What is the treatment approach for Henoch-Schönlein purpura?

- A. Immediate surgery
- B. Self-limited course**
- C. Long-term antibiotic therapy
- D. Daily corticosteroids

Henoch-Schönlein purpura (HSP) is a small-vessel vasculitis that primarily affects children, characterized by a tetrad of symptoms: purpura, typically on the buttocks and legs, joint pain, abdominal pain, and renal involvement. The treatment approach is largely supportive, as HSP often follows a self-limited course. Patients typically experience resolution of symptoms within weeks to months without necessitating aggressive interventions. In many cases, supportive care includes pain management and monitoring for any complications, particularly concerning the kidneys. While corticosteroids may be used in severe cases or if there's significant abdominal pain, they are not routinely required for most patients, further underscoring the self-limiting nature of the condition. Surgery is not indicated as a treatment for HSP, as the condition often resolves spontaneously. Long-term antibiotic therapy is unnecessary, as HSP is not caused by an infection, and daily corticosteroids are not a standard treatment regimen, as they do not significantly alter the course of the disease for most patients. Therefore, the most accurate approach to managing Henoch-Schönlein purpura is recognizing its self-limited nature.

7. What is another name for Christmas tree disease?

- A. Pityriasis rosea**
- B. Psoriasis vulgaris
- C. Atopic dermatitis
- D. Seborrheic dermatitis

Christmas tree disease is commonly known as pityriasis rosea. This skin condition is characterized by the appearance of a distinctive rash that often starts with a single oval or round "herald patch," followed by smaller lesions that tend to form a pattern resembling a Christmas tree on the back. This name comes from the arrangement and appearance of the rashes as they spread outward in a shape that resembles a tree. The other conditions mentioned, such as psoriasis vulgaris, atopic dermatitis, and seborrheic dermatitis, have different characteristics and presentations. Psoriasis vulgaris involves silvery scales and can affect various parts of the body, while atopic dermatitis is often chronic and associated with itchiness and inflammation. Seborrheic dermatitis typically appears as greasy, scaly patches primarily on the scalp and face. Each of these conditions is distinct and does not exhibit the hallmark "Christmas tree" pattern seen in pityriasis rosea.

8. Which antibiotic is specifically recommended for Gustilo Anderson type 1 fractures?

- A. Amikacin
- B. Ciprofloxacin
- C. First or second-generation cephalosporin**
- D. Gentamicin

Gustilo-Anderson type 1 fractures are open fractures characterized by a clean wound that is less than 1 cm in size. The primary concern in managing such injuries is preventing infection, particularly due to skin flora and the potential for contamination with bacteria from the surrounding environment. First or second-generation cephalosporins are specifically recommended for type 1 fractures because they target common pathogens associated with clean skin lacerations, including Staphylococcus aureus and Streptococcus species. These antibiotics are effective against the typical flora found on the skin, which is crucial when dealing with a clean wound that has been exposed to the outside environment. The effectiveness of cephalosporins in preventing deep infection and their favorable safety profile make them a suitable choice for empirical therapy in this type of fracture. They can also be easily administered in a surgical or outpatient setting, which aligns with the management protocols for such injuries. Other antibiotics listed do not specifically align with the typical management criteria for Gustilo-Anderson type 1 fractures. For instance, amikacin and gentamicin are more commonly used for infections requiring coverage against more resistant Gram-negative organisms, while ciprofloxacin is not a first-line choice for standard prophylaxis in open fractures. Thus, using

9. What complication is most commonly caused by lidocaine toxicity?

- A. Hypertension
- B. Bradycardia**
- C. Severe headache
- D. Nausea

Lidocaine toxicity can lead to various cardiovascular effects, with bradycardia being one of the most significant and commonly observed complications. Bradycardia refers to an abnormally slow heart rate, which can result from lidocaine's mechanism of action as a sodium channel blocker. At toxic levels, lidocaine can interfere with the normal conduction pathways in the heart, leading to impaired cardiac excitability and rhythmicity. This can manifest as bradycardia and, in more severe cases, can progress to more critical cardiac arrhythmias. While other options like hypertension, severe headache, and nausea can occur due to various factors in different contexts, they are not as directly linked to lidocaine toxicity as bradycardia is. Hypertension is not a typical consequence of lidocaine toxicity; in fact, lidocaine can often lower blood pressure due to its vasodilatory properties. Severe headache and nausea can also occur for various other reasons, but they do not have the same direct relationship with lidocaine toxicity that bradycardia has. This makes bradycardia the most appropriate and common complication to associate with lidocaine toxicity.

10. What is the chance of survival for a split thickness skin graft?

- A. Low
- B. High**
- C. Moderate
- D. Variable

The chance of survival for a split thickness skin graft is considered high because this type of graft is designed to maintain a significant portion of the dermis along with the epidermis. This allows for better integration and healing when the graft is placed on a well-vascularized recipient site. The presence of dermal elements contributes to the graft's ability to revascularize and survive post-operatively, leading to a more favorable outcome compared to other types of grafts. A high chance of survival is particularly enhanced in cases where the recipient site is prepared properly and the graft is applied meticulously, allowing for optimal healing conditions. Factors such as the overall health of the patient, the presence of infection, and the area's vascularity can influence the graft's success, but generally, split thickness skin grafts have a good success rate in clinical practice. Other options, such as low, moderate, or variable, do not accurately reflect the typical outcomes associated with split thickness skin grafting. A low chance of survival would not align with the known efficacy of this graft type, while describing the survival rate as variable or moderate does not capture the generally positive prognosis seen with this surgical technique.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://podiatricmedabpm.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE