AMERICAN MIDWIFERY CERTIFICATION BOARD (AMCB) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://midwiferyamcb.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is one of the risk factors associated with placental abruption?

A. History of previous vaginal birth

B. Smoking tobacco

C. Age under 25

D. Excessive exercise

2. When does implantation typically occur after fertilization?

A. 3-5 days

B. 6-7 days

C. 8-10 days

D. 11-14 days

3. What physical examination finding may indicate trichomoniasis?

A. Pelvic pain

B. This discharge is odorless

C. Punctate lesions on the cervix

D. Normal vaginal flora

4. What is the typical percentage of spontaneous abortions that occur in the first 12 weeks of pregnancy?

A. 30%

B. 50%

C. 70%

D. 80%

5. What is the recommended calories and protein intake per day for pregnancy?

A. 2000 kcal/day and 40g/day of protein

B. 2500 kcal/day and 60g/day of protein

C. 3000 kcal/day and 80g/day of protein

D. 1800 kcal/day and 30g/day of protein

SAMPLE

6. Which factors may contribute to weakness of pelvic support structures?

A. Increased estrogen

B. Neuromuscular injury during childbirth

C. Improved pelvic strength exercises

D. Regular hormonal treatments

7. In the treatment of community-acquired pneumonia with risk factors for DRSP, which antibiotic is recommended?

A. Penicillin

B. Ciprofloxacin

C. Respiratory fluoroquinolone

D. Doxycycline

8. Which of the following is one of Amsel's Criteria for diagnosing Bacterial Vaginosis?

A. Vaginal pH of less than 4.5

B. Positive whiff test

C. Presence of yeast cells

D. Thick white discharge

9. What defines anterior asynclitism in fetal presentation?

A. When the sagittal suture is aligned centrally

B. When the anterior parietal bone is presenting first

C. Presence of multiple angles of descent

D. When the occiput is closer to the sacrum

10. What is the common duration of menstrual flow considered normal?

A. 2-4 days

B. 4.5 to 8 days

C. 5-10 days

D. 3-7 days

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. D
5. B
6. B
7. C
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is one of the risk factors associated with placental abruption?

A. History of previous vaginal birth

B. Smoking tobacco

C. Age under 25

D. Excessive exercise

Smoking tobacco is indeed recognized as one of the significant risk factors associated with placental abruption. The harmful substances in tobacco smoke can lead to various complications during pregnancy, including vascular constriction and reduced blood flow to the placenta. This can increase the likelihood of the placenta separating from the uterine wall prematurely, resulting in placental abruption. The impact of smoking on placental health and overall pregnancy outcomes highlights the importance of addressing tobacco use in expectant mothers to reduce the risk of complications such as abruption. In contrast, while a history of previous vaginal birth, being under age 25, or engaging in excessive exercise might be relevant to other aspects of pregnancy, they are not specifically linked to an increased risk of placental abruption in the same direct manner as smoking is.

2. When does implantation typically occur after fertilization?

A. 3-5 days

B. 6-7 days

C. 8-10 days

D. 11-14 days

Implantation typically occurs 6 to 7 days after fertilization. Following fertilization, the zygote begins its journey down the fallopian tube, undergoing several divisions as it travels. By the time it reaches the uterus, it has developed into a blastocyst, which is the stage that is ready for implantation. The blastocyst usually arrives in the uterus about 5 days after fertilization. Once in the uterine cavity, it continues to develop and prepares for implantation into the uterine lining. The actual process of implantation, where the blastocyst attaches to the endometrium, takes place between the 6th and 7th day after fertilization, marking a critical stage in establishing a pregnancy. This timing is important for successful implantation and subsequent development of the embryo.

3. What physical examination finding may indicate trichomoniasis?

A. Pelvic pain

B. This discharge is odorless

C. Punctate lesions on the cervix

D. Normal vaginal flora

The presence of punctate lesions on the cervix is a noteworthy physical examination finding that may indicate trichomoniasis. In women with trichomoniasis, the cervix can appear erythematous and may have these specific lesions, often referred to as "strawberry cervix," due to the inflammation and capillary dilation that occur as a result of the infection. This distinct appearance is not commonly seen with other vaginitis etiologies and can be a key clinical sign for diagnosis. Other findings associated with trichomoniasis may include a profuse, frothy, greenish-yellow discharge, often accompanied by a foul odor, which further differentiates it from other conditions. This context is essential to recognize, as it helps in clinical assessments and guiding appropriate treatment protocols for the infection.

SAMPLE

4. What is the typical percentage of spontaneous abortions that occur in the first 12 weeks of pregnancy?

A. 30%

B. 50%

C. 70%

D. 80%

The correct percentage of spontaneous abortions that typically occur in the first 12 weeks of pregnancy is 80%. This high percentage can be attributed to several factors. Early pregnancy losses are often linked to chromosomal abnormalities, which account for a significant portion of first-trimester miscarriages. Additionally, the risk of miscarriage decreases as the pregnancy progresses past this initial stage, where many embryos that are not viable are lost. It's important to understand that while estimates can vary depending on the population studied and specific circumstances, the consensus in obstetric and reproductive health literature generally identifies that a substantial number of spontaneous abortions occur within the first trimester. The first 12 weeks of gestation is a critical period for embryonic development, and factors such as hormonal support, uterine environment, and maternal health all play essential roles in the viability of the pregnancy during this time. In contrast, percentages lower than this do not accurately reflect the broader evidence in medical literature regarding the incidence of early pregnancy loss.

5. What is the recommended calories and protein intake per day for pregnancy?

A. 2000 kcal/day and 40g/day of protein

B. 2500 kcal/day and 60g/day of protein

C. 3000 kcal/day and 80g/day of protein

D. 1800 kcal/day and 30g/day of protein

During pregnancy, nutritional needs increase to support the developing fetus and maintain the health of the mother. The recommended caloric intake for pregnant women typically rises to around 2500 kcal per day. This additional energy helps support fetal growth, amniotic fluid production, and increased metabolic demands on the body. In terms of protein, the recommendation is to consume about 60 grams of protein per day. Protein is essential for the growth of fetal tissues, including the brain, as well as the expansion of maternal tissues such as the uterus and breasts. Adequate protein intake is critical for optimal developing fetus health and maternal wellbeing during pregnancy. This particular choice aligns well with nutritional guidelines that focus on the importance of both sufficient caloric and protein intake during pregnancy, ensuring that both the mother and baby receive the necessary nutrients to thrive.

SAMPLE

6. Which factors may contribute to weakness of pelvic support structures?

A. Increased estrogen

B. Neuromuscular injury during childbirth

C. Improved pelvic strength exercises

D. Regular hormonal treatments

Neuromuscular injury during childbirth is a significant factor that can contribute to the weakness of pelvic support structures. During childbirth, especially in vaginal deliveries, there can be trauma to the pelvic floor muscles and nerves, leading to issues such as pelvic organ prolapse or incontinence. The strain placed on the pelvic muscles during labor and delivery can result in damage, altering the support that these structures provide to the pelvic organs. As a result, this can weaken the overall integrity of the pelvic support system and may increase the likelihood of subsequent complications. On the other hand, increased estrogen is generally associated with improvements in pelvic tissue elasticity and strength, while regular hormonal treatments can help maintain pelvic health. Improved pelvic strength exercises are designed to enhance muscle tone and support structures, helping to mitigate weakness rather than contributing to it. Thus, none of these options align with factors that specifically contribute to the weakening of pelvic support structures in the same way that neuromuscular injury during childbirth does.

7. In the treatment of community-acquired pneumonia with risk factors for DRSP, which antibiotic is recommended?

A. Penicillin

B. Ciprofloxacin

C. Respiratory fluoroquinolone

D. Doxycycline

In the treatment of community-acquired pneumonia (CAP) when there are risk factors for drug-resistant Streptococcus pneumoniae (DRSP), respiratory fluoroquinolones are the recommended antibiotic choice due to their broad spectrum of activity against prevalent pathogens, including DRSP. Respiratory fluoroquinolones, such as moxifloxacin and gemifloxacin, are particularly effective because they not only target typical bacterial pathogens but also have increased activity against atypical organisms and are better tolerated in patients with specific risk factors that might contribute to the development of antibiotic resistance. When patients present with risk factors for DRSP—such as recent antibiotic use, age over 65, or comorbidities—broad-spectrum treatment is essential to ensure that resistant strains are adequately covered. Respiratory fluoroquinolones offer the necessary efficacy while also addressing potential resistance patterns that could complicate treatment. The other antibiotics listed may not provide the same level of coverage needed in these situations. Penicillin is more suited for infections caused by penicillin-susceptible strains, which may not apply when resistance is a concern. Ciprofloxacin, while effective against some types of bacteria, is not typically used for respiratory infections due to insufficient coverage of Streptococcus

SAMPLE

8. Which of the following is one of Amsel's Criteria for diagnosing Bacterial Vaginosis?

A. Vaginal pH of less than 4.5

B. Positive whiff test

C. Presence of yeast cells

D. Thick white discharge

Amsel's Criteria for diagnosing Bacterial Vaginosis includes specific clinical findings that help differentiate it from other types of vaginal infections. The positive whiff test is a key component of this diagnostic criteria. In this test, a fishy odor is produced when potassium hydroxide is added to a sample of vaginal fluid, indicating the presence of amines released by certain bacteria associated with Bacterial Vaginosis. This symptom is particularly compelling because the odor often becomes more pronounced after sexual intercourse or during menstruation, which can help clinicians identify the condition more reliably. The whiff test is a simple yet effective clinical tool that, when combined with other criteria, aids in the diagnosis of Bacterial Vaginosis, making it a critical element of Amsel's Criteria.

9. What defines anterior asynclitism in fetal presentation?

A. When the sagittal suture is aligned centrally

B. When the anterior parietal bone is presenting first

C. Presence of multiple angles of descent

D. When the occiput is closer to the sacrum

Anterior asynclitism is characterized by the position of the fetal head during labor, specifically in relation to the mother's pelvis. In this scenario, the term refers to a situation where the anterior parietal bone of the fetal head is positioned in a leading or presenting manner. This occurs when the fetal head is tilted in such a way that the anterior fontanelle is closer to the pubic bone compared to the posterior fontanelle. This presentation can significantly affect the progress of labor and may lead to a prolonged labor if the fetal position needs correction. When the anterior parietal bone is presenting first, it indicates a positional variation which can complicate the descent through the birth canal unless managed appropriately. Understanding this definition is crucial for midwives and obstetricians as it guides them in monitoring and potentially correcting fetal position during labor.

10. What is the common duration of menstrual flow considered normal?

A. 2-4 days

B. 4.5 to 8 days

C. 5-10 days

D. 3-7 days

The common duration of menstrual flow typically considered normal is between 3 to 7 days. This time frame encompasses the average length of menstruation experienced by many individuals. A flow duration of 2-4 days may be considered shorter than average, while a duration of 5-10 days extends beyond the usual norm and might indicate underlying health issues such as hormonal imbalances or other conditions. The range of 4.5 to 8 days, while inclusive of some variations, is less commonly cited in clinical guidelines than the range of 3 to 7 days. Thus, the option specifying 3-7 days aligns more closely with established medical standards for menstrual flow.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://midwiferyamcb.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE