ANCC FAMILY NURSE PRACTITIONER PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://anccfamilynurse.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which condition is associated with an increased risk of ectopic pregnancy?

A. Normal pelvic anatomy

B. PID and prior abortions

C. Regular menstrual cycles

D. Previous vaginal births

2. What is a first-line treatment option for hereditary essential tremor?

A. Atenolol

B. Primidone

C. Propranolol

D. Timolol

3. What type of tissue is affected by a growth plate fracture?

A. Cartilage

B. Muscle

C. Growing tissue

D. Fatty tissue

4. What is the primary goal of using steroid injections for lateral epicondylitis?

A. To improve strength

B. To reduce inflammation

C. To increase mobility

D. To provide long-term relief

5. What is sialolithiasis more commonly known as?

A. Salivary calculi

B. Oral thrush

C. Dry mouth syndrome

D. Salivary gland infection

6. Which of the following is NOT a sign of anticholinergic toxicity?

A. Mydriasis

B. Diarrhea

C. Flushed skin

D. Increased body temperature

SAMPLE

7. Which type of medication is indicated for managing hypertension in a patient with diabetic neuropathy?

A. Thiazide

B. Calcium channel blockers

C. ACE inhibitors or ARBs

D. Beta-blockers

8. What is the recommended diagnostic tool for ruling out an Abdominal Aortic Aneurysm (AAA) in a patient aged 65 or older?

A. X-ray

B. CT scan

C. Ultra sound

D. MRI

9. What effect do NSAIDs have on cardiac function?

A. Decrease heart rate

B. Inhibit prostaglandins reaction in blood vessels

C. Increase blood flow to the heart

D. None of the above

10. Which laboratory finding is indicative of an acute Hepatitis B infection?

A. Negative Anti-HBs

B. Positive HBsAg

C. Positive Anti-HBc

D. Positive Anti-HBe

SAMPLE

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. A
6. B
7. C
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which condition is associated with an increased risk of ectopic pregnancy?

A. Normal pelvic anatomy

B. PID and prior abortions

C. Regular menstrual cycles

D. Previous vaginal births

The association of pelvic inflammatory disease (PID) and prior abortions with an increased risk of ectopic pregnancy is well-documented in medical literature. PID often results in scarring and damage to the fallopian tubes, which can impede the passage of a fertilized egg. If a fertilized egg is unable to move properly through the uterine tubes due to such damage, it may implant outside the uterus, commonly within the tube itself, leading to an ectopic pregnancy. Additionally, prior abortions, especially if they involve surgical procedures, can similarly result in changes to the uterine or tubal anatomy, increasing the likelihood of ectopic implantation. These factors combined contribute significantly to the risk profile for ectopic pregnancies, emphasizing the importance of a detailed gynecological history in assessing a patient's risk factors. In contrast, normal pelvic anatomy, regular menstrual cycles, and previous vaginal births do not present the same risk of obstructing the normal passage of the fertilized egg, thereby not significantly increasing the likeliness of ectopic pregnancies. Understanding these underlying mechanisms helps in recognizing the importance of thorough assessments in patients who may be at risk.

2. What is a first-line treatment option for hereditary essential tremor?

A. Atenolol

B. Primidone

C. Propranolol

D. Timolol

In the management of hereditary essential tremor, propranolol is recognized as a first-line treatment option. Propranolol, a non-selective beta-blocker, works by dampening the physiological response involved in tremors. It reduces the amplitude of the tremor and can improve overall control of tremulous movements, particularly in patients with essential tremor who experience significant functional impairment. When considering the mechanism of action, propranolol focuses on the beta-adrenergic receptors, leading to decreased sympathetic nervous system activity, which is implicated in the exacerbation of tremors. The effectiveness of propranolol has been supported by numerous studies, and it is often the medication of choice for both primary and secondary essential tremor, particularly when the tremor is affecting daily activities such as writing or eating. The other potential treatment options listed may also play a role in managing tremor syndromes, but they do not have the same level of established effectiveness as propranolol specifically for hereditary essential tremor. For instance, while atenolol and timolol are beta-blockers, they are less commonly used in this context. Primidone, an anticonvulsant, can also be used to treat essential tremor but is generally considered when

SAMPLE

3. What type of tissue is affected by a growth plate fracture?

A. Cartilage

B. Muscle

C. Growing tissue

D. Fatty tissue

A growth plate fracture primarily affects the areas of developing tissue at the ends of long bones, known as the epiphyseal plate or growth plate. This structure is made up of cartilage, which is crucial for bone growth and development during childhood and adolescence. When injuries occur in this area, they can significantly impact the growth potential of the bone if not treated appropriately. The concept of "growing tissue" encompasses the cartilage present in the growth plate, but it also refers to the overall physiological processes occurring in the area. This makes it the most encompassing and accurate description of the tissue affected by a growth plate fracture. Fractures in this region can disrupt normal growth patterns, leading to potential complications, such as differential growth or deformities. Other tissues, like muscle and fatty tissue, are not directly involved in the growth plates and do not adapt or change in response to growth plate injuries in the same way. Therefore, while cartilage is an important component, the term "growing tissue" more comprehensively captures the nature and implications of a growth plate fracture.

4. What is the primary goal of using steroid injections for lateral epicondylitis?

A. To improve strength

B. To reduce inflammation

C. To increase mobility

D. To provide long-term relief

The primary goal of using steroid injections for lateral epicondylitis is to reduce inflammation. Lateral epicondylitis, commonly known as tennis elbow, involves pain and inflammation around the outer part of the elbow due to overuse or repetitive strain injuries. Steroid injections contain corticosteroids, which are powerful anti-inflammatory agents. By directly injecting these medications into the affected area, the intention is to alleviate pain and reduce inflammatory processes, thereby promoting a more favorable environment for healing and recovery. While improving strength and increasing mobility are important aspects of rehabilitation following lateral epicondylitis, they are not the immediate goals of steroid injections. These injections serve primarily to address the acute symptoms of pain and inflammation. Providing long-term relief is typically a result of a comprehensive treatment approach that may include physical therapy, activity modification, and gradual strengthening exercises after managing the inflammation. Therefore, the use of steroid injections specifically targets the reduction of inflammation to alleviate pain in the short term.

SAMPLE

5. What is sialolithiasis more commonly known as?

A. Salivary calculi

B. Oral thrush

C. Dry mouth syndrome

D. Salivary gland infection

Sialolithiasis is more commonly known as salivary calculi, which refers to the formation of stones or calcified structures within the salivary glands or ducts. These stones can block the flow of saliva, leading to swelling, pain, and potential infection in the affected gland. Understanding this term is essential for healthcare professionals, as it helps in diagnosing conditions related to salivary gland dysfunction. The other options represent different conditions that are not synonymous with sialolithiasis. Oral thrush is a fungal infection in the mouth, dry mouth syndrome refers to xerostomia, or a lack of salivary production, and a salivary gland infection is often a result of sialolithiasis but does not directly define the presence of stones in the glands. The specificity of the term "salivary calculi" accurately captures the nature of the problem in sialolithiasis.

6. Which of the following is NOT a sign of anticholinergic toxicity?

A. Mydriasis

B. Diarrhea

C. Flushed skin

D. Increased body temperature

Anticholinergic toxicity is characterized by a range of symptoms that result from the inhibition of acetylcholine in the parasympathetic nervous system. Common signs of this condition include mydriasis (dilated pupils), flushed skin, and increased body temperature due to decreased sweating and disrupted thermoregulation. Mydriasis occurs as a direct effect of anticholinergic agents blocking the action of acetylcholine on the iris sphincter muscle, leading to pupil dilation. Flushed skin results from peripheral vasodilation, another outcome of disrupted cholinergic function, while increased body temperature is a consequence of reduced sweating and the overheating that can occur in this state. In the context of gastrointestinal symptoms, diarrhea is typically associated with an increase in acetylcholine activity, particularly in the enteric nervous system, which governs the gastrointestinal tract. Thus, diarrhea does not align with the symptoms of anticholinergic toxicity. Instead, one would expect to see symptoms like constipation, as anticholinergic agents slow down gastrointestinal motility. Therefore, identifying diarrhea as a non-sign of anticholinergic toxicity highlights a key understanding of how these medications influence the autonomic nervous system and reflects a more comprehensive knowledge of the drug's effects on

SAMPLE

7. Which type of medication is indicated for managing hypertension in a patient with diabetic neuropathy?

A. Thiazide

B. Calcium channel blockers

C. ACE inhibitors or ARBs

D. Beta-blockers

In patients with diabetic neuropathy, ACE inhibitors or angiotensin receptor blockers (ARBs) are particularly beneficial for managing hypertension due to their renal protective effects. Diabetic neuropathy often involves underlying renal complications; therefore, using medications that can mitigate these issues is crucial. ACE inhibitors work by inhibiting the angiotensin-converting enzyme, which results in vasodilation and a reduction in blood pressure. These medications have the added benefit of reducing proteinuria and slowing the progression of diabetic nephropathy, making them a first-line choice for hypertensive patients with diabetes. ARBs share similar efficacy and are often used in patients who may not tolerate ACE inhibitors due to side effects such as cough. Both classes of medications improve cardiovascular outcomes and provide renal protection in individuals with diabetes-related complications, reinforcing their role as a preferred choice in this patient population. Other classes of antihypertensive medications, while effective at lowering blood pressure, do not offer the same renal protective benefits that ACE inhibitors and ARBs provide, making them less suitable for managing hypertension specifically related to diabetic neuropathy.

8. What is the recommended diagnostic tool for ruling out an Abdominal Aortic Aneurysm (AAA) in a patient aged 65 or older?

A. X-ray

B. CT scan

C. Ultra sound

D. MRI

The recommended diagnostic tool for ruling out an Abdominal Aortic Aneurysm (AAA) in a patient aged 65 or older is an ultrasound. This non-invasive imaging technique is safe, quick, and particularly effective for this purpose. An ultrasound can provide real-time imaging of the abdominal aorta, allowing healthcare providers to visualize any dilation or abnormal enlargement indicative of an AAA. This demographic, particularly men aged 65 and older, is at increased risk for AAAs due to factors such as age, hypertension, and a history of smoking. Therefore, screening with an ultrasound is recommended as part of preventive care in this population. While CT scans and MRIs can also be used for a more detailed evaluation of known AAAs, ultrasound remains the first-line screening tool because of its accessibility, low cost, and the absence of ionizing radiation. X-rays are not used for this purpose as they do not provide sufficient detail to evaluate vascular structures, making them ineffective for ruling out an AAA.

SAMPLE

9. What effect do NSAIDs have on cardiac function?

A. Decrease heart rate

B. Inhibit prostaglandins reaction in blood vessels

C. Increase blood flow to the heart

D. None of the above

Non-steroidal anti-inflammatory drugs (NSAIDs) primarily work by inhibiting the enzyme cyclooxygenase (COX), which is essential for the formation of prostaglandins. Prostaglandins have various roles in the body, including promoting inflammation and regulating vascular tone. By inhibiting prostaglandin synthesis, NSAIDs can have significant effects on blood vessels, often leading to vasoconstriction rather than vasodilation. This indicates their potential to affect hemodynamics, such as increasing blood pressure, and can have implications for cardiac function, particularly in patients with preexisting cardiovascular conditions. Understanding this dynamic is crucial, especially for clinicians managing patients with cardiovascular risk factors or those on NSAIDs. While the inhibition of prostaglandins can lead to decreased inflammation and pain relief, it may also compromise vascular function and cardiac health by altering blood vessel reactivity. Thus, the option regarding the inhibition of prostaglandin reactions in blood vessels accurately reflects the impact of NSAIDs on cardiac function and overall circulatory dynamics.

10. Which laboratory finding is indicative of an acute Hepatitis B infection?

A. Negative Anti-HBs

B. Positive HBsAg

C. Positive Anti-HBc

D. Positive Anti-HBe

A positive HBsAg indicates that the Hepatitis B virus is actively replicating in the body. HBsAg, or Hepatitis B surface antigen, is the first serological marker that becomes detectable during an acute HBV infection and typically appears within 1-10 weeks after exposure. The presence of HBsAg is essential for the diagnosis of acute hepatitis B, as it confirms that the virus is currently present in the bloodstream and the individual is potentially infectious. In contrast, the other markers listed provide different information that does not directly indicate an acute infection. For example, negative Anti-HBs suggests that there is no immunity to Hepatitis B, while a positive Anti-HBc would imply past or current infection but does not specifically confirm active infection as HBsAg does. Positive Anti-HBe indicates low levels of viral replication and may be seen in chronic infections but is not a primary indicator of an acute infection. Therefore, HBsAg is the definitive laboratory finding for confirming an acute Hepatitis B infection.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://anccfamilynurse.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE