
ANESTHESIA KNOWLEDGE TEST 24 (AKT-24) PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://anesthesiaakt24.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the osmolarity of Lactated Ringer's solution?

A. 154 mOsm

B. 274 mOsm

C. 252 mOsm

D. 330 mOsm

2. Which of the following can lead to increased myocardial oxygen consumption (MVo2)?

A. Bradycardia

B. Shivering

C. Low blood pressure

D. Hypothermia

3. Which part of the larynx is manipulated during the BURP maneuver?

A. Cricoid cartilage

B. Epiglottis

C. Thyroid cartilage

D. Arytenoid cartilage

4. Which of the following is a specific risk factor for pulmonary complications following minimally invasive thymectomy?

A. BMI greater than 28

B. Age less than 30 years

C. History of hypertension

D. Severe anxiety disorders

5. What is a shunt in the context of respiratory function?

A. Ventilation without perfusion

B. Perfusion without ventilation

C. Normal ventilation

D. Excessive ventilation

SAMPLE

6. Which cranial nerve is responsible for the sensory afferent limb of the gag reflex?

A. CN 7

B. CN 9

C. CN 10

D. CN 12

7. Why does lidocaine in tumescent anesthesia have reduced systemic toxicity?

A. It dilutes in the bloodstream

B. It is combined with local vasoconstrictors

C. It is lipophilic and sequestered in adipose tissue

D. It is rapidly metabolized by the liver

8. In pediatric anesthesia, which condition is most likely to increase the risk of apnea?

A. Premature birth

B. Bacterial infections

C. Non-anesthetic medications

D. Low blood pressure

9. How does PEEP affect Functional Residual Capacity (FRC)?

A. It decreases FRC.

B. It has no effect on FRC.

C. It increases FRC.

D. It causes FRC to become non-calculable.

10. Which block provides a reliable block of the orbicularis oculi muscle for eye surgery?

A. Retrobulbar block

B. Sphenopalatine block

C. Peribulbar block

D. Topical anesthetic only

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. A
5. B
6. B
7. C
8. A
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. What is the osmolarity of Lactated Ringer's solution?

A. 154 mOsm

B. 274 mOsm

C. 252 mOsm

D. 330 mOsm

Lactated Ringer's solution is an isotonic intravenous fluid commonly used for fluid resuscitation. It contains a mixture of electrolytes, including sodium, potassium, calcium, and lactate as the buffering agent, which collectively contribute to its osmolarity. The osmolarity of Lactated Ringer's solution is approximately 274 mOsm/L. This value reflects the total concentration of dissolved particles in the solution, which includes not just the sodium and chloride ions, but also potassium, calcium, and lactate. It is significant that the osmolarity of Lactated Ringer's solution is designed to be similar to that of plasma, making it an effective choice for volume replacement while maintaining isotonicity, thus preventing hemolysis or complications associated with the administration of hypotonic or hypertonic solutions. The other options provided do not reflect the correct osmolarity of Lactated Ringer's solution and typically underrepresent or overrepresent the concentration of solutes in this solution. Understanding the precise osmolarity is crucial for clinical applications, especially in perioperative fluid management and resuscitation protocols.

2. Which of the following can lead to increased myocardial oxygen consumption (MVo2)?

A. Bradycardia

B. Shivering

C. Low blood pressure

D. Hypothermia

Increased myocardial oxygen consumption (MVo2) occurs when the heart muscle requires more oxygen to meet the demands of the body. Shivering is the body's response to cold temperatures, leading to increased muscle activity to generate heat. This muscle activity raises the metabolic rate, which in turn increases cardiac output and heart rate to supply more oxygen to the working muscles. As a consequence, the heart consumes more oxygen to support this heightened metabolic state. In contrast, bradycardia typically lowers MVo2 due to reduced heart rate, which decreases the workload on the heart. Low blood pressure may also lead to inadequate perfusion and oxygen delivery, resulting in a non-demanding state for the myocardium. Hypothermia can lower metabolism and therefore decrease MVo2 as the heart rate and overall body demands are reduced. By understanding the physiological mechanisms of shivering and its impact on metabolism and cardiovascular function, it's clear why this response significantly increases myocardial oxygen consumption compared to the other options.

SAMPLE

3. Which part of the larynx is manipulated during the BURP maneuver?

A. Cricoid cartilage

B. Epiglottis

C. Thyroid cartilage

D. Arytenoid cartilage

The BURP maneuver involves the manipulation of the thyroid cartilage. This technique is used to improve the visibility of the glottic opening during intubation by adjusting the position of the larynx. When the thyroid cartilage is pushed backward and upward, it often moves the larynx into a more favorable position, allowing for easier access to the vocal cords. This maneuver is particularly useful in cases where the airway is difficult to visualize, as it can help align the trachea with the line of sight for the intubating device. The manipulation of other parts of the larynx, such as the cricoid cartilage, epiglottis, or arytenoid cartilage, does not achieve the same purpose in the context of the BURP maneuver. While the cricoid cartilage is important in airway management and can be subject to pressure during various techniques, it does not facilitate the improvement of the view in the same way as the thyroid cartilage. The epiglottis and arytenoids serve different roles in the airway and are not the primary focus in this particular maneuver aimed at enhancing the approach to the vocal cords.

4. Which of the following is a specific risk factor for pulmonary complications following minimally invasive thymectomy?

A. BMI greater than 28

B. Age less than 30 years

C. History of hypertension

D. Severe anxiety disorders

Having a Body Mass Index (BMI) greater than 28 is recognized as a specific risk factor for pulmonary complications following minimally invasive thymectomy. Higher BMI is often associated with obesity, which can lead to a number of respiratory issues. Obese patients may have decreased lung volumes due to reduced chest wall compliance, increased airway resistance, and impaired gas exchange due to changes in pulmonary mechanics. These factors can significantly complicate the perioperative management of patients undergoing thoracic surgery, leading to a higher incidence of postoperative pulmonary complications such as atelectasis, pneumonia, and respiratory failure. When considering other risk factors listed, age less than 30 years typically is not associated with increased risk of pulmonary complications in this specific surgical population, as younger patients generally tend to have better respiratory function and lower comorbidity rates. A history of hypertension may have various implications for cardiovascular health but does not specifically indicate a heightened risk for pulmonary complications. Severe anxiety disorders can impact a patient's overall well-being and may affect perioperative care but do not directly correlate with an increased risk of pulmonary issues related specifically to thymectomy. Thus, the evidence supports the notion that high BMI is a key concern when evaluating patients for this procedure.

SAMPLE

5. What is a shunt in the context of respiratory function?

A. Ventilation without perfusion

B. Perfusion without ventilation

C. Normal ventilation

D. Excessive ventilation

A shunt in respiratory function refers to a situation where blood moves from the right side of the heart to the left side without being oxygenated in the lungs. This occurs when there is perfusion without ventilation, meaning that blood is flowing through the pulmonary capillaries but is not exposed to fresh oxygen because the alveoli are not receiving air. This can happen in various clinical scenarios, such as in conditions like pneumonia or acute respiratory distress syndrome (ARDS), where alveolar collapse or fluid-filled alveoli leads to ineffective gas exchange. Understanding shunts is crucial because they indicate poor oxygenation of blood, which can lead to systemic hypoxia. In contrast, ventilation without perfusion, normal ventilation, and excessive ventilation do not represent the mechanism of a shunt as they either denote adequate or impaired exchanges of air and blood in a more specific manner that does not lead to the same consequences as shunting.

6. Which cranial nerve is responsible for the sensory afferent limb of the gag reflex?

A. CN 7

B. CN 9

C. CN 10

D. CN 12

The cranial nerve responsible for the sensory afferent limb of the gag reflex is the glossopharyngeal nerve, also known as cranial nerve 9. This nerve plays a crucial role in providing sensory innervation to the posterior one-third of the tongue and the oropharynx. When these areas are stimulated, sensory information is sent via the glossopharyngeal nerve to the brainstem, which processes this information and coordinates the gag reflex response. The gag reflex itself involves both a sensory input and a motor output. The sensory input is crucial for detecting potentially harmful stimuli in the throat, and this specific role is fulfilled by the glossopharyngeal nerve. Once the sensory receptors in this area are activated, the response is mediated by the efferent pathways involving other cranial nerves, but the initial sensory trigger comes from cranial nerve 9. Understanding this mechanism highlights the importance of cranial nerve 9 in protecting the airway and preventing aspiration, making it essential in clinical settings, particularly during procedures that involve the throat or when assessing a patient's ability to protect their airway.

SAMPLE

7. Why does lidocaine in tumescent anesthesia have reduced systemic toxicity?

A. It dilutes in the bloodstream

B. It is combined with local vasoconstrictors

C. It is lipophilic and sequestered in adipose tissue

D. It is rapidly metabolized by the liver

In tumescent anesthesia, lidocaine exhibits reduced systemic toxicity primarily due to its lipophilic properties, which allow it to be sequestered in adipose tissue. This retention in the fat compartments decreases the concentration of lidocaine in the bloodstream, effectively minimizing the potential for systemic toxic effects. By residing in the adipose tissue, lidocaine is gradually released, leading to a prolonged duration of action while mitigating any immediate harmful systemic effects. This mechanism is particularly beneficial in cosmetic and surgical procedures where large volumes of anesthetic might be used, as it permits the safe administration of larger doses than would typically be permissible if delivered into the systemic circulation directly. Additionally, the body's ability to manage and metabolize the drug is more favorable when it is not circulating freely in the bloodstream at high concentrations, thus emphasizing the protective role of lipophilicity in reducing systemic toxicity during tumescent anesthesia procedures.

8. In pediatric anesthesia, which condition is most likely to increase the risk of apnea?

A. Premature birth

B. Bacterial infections

C. Non-anesthetic medications

D. Low blood pressure

Premature birth is indeed associated with an increased risk of apnea in pediatric patients. This condition is often due to the immaturity of the central nervous system in premature infants, which can lead to disruptions in the control of breathing. Apnea, which is defined as a pause in breathing that lasts for more than 20 seconds, can occur in neonates and infants who are born before the term, particularly in those who are less than 28 weeks of gestation. The underdevelopment of the respiratory centers in the brain of premature infants means they are less capable of maintaining stable breathing patterns. This can be exacerbated by other factors common in this population, such as respiratory distress syndrome and the need for resuscitation immediately after birth. Therefore, healthcare providers in anesthesiology and pediatrics must be particularly vigilant when managing anesthesia in premature infants, as they have a higher propensity for apneic events, especially when exposed to anesthetic agents. In contrast, while bacterial infections can potentially impact respiratory function, the direct correlation with apnea risk is less established compared to the effects of prematurity. Non-anesthetic medications may influence a variety of physiological responses but do not have a specific or consistent link to increased apnea risk like prematurity does. Low blood pressure

SAMPLE

9. How does PEEP affect Functional Residual Capacity (FRC)?

A. It decreases FRC.

B. It has no effect on FRC.

C. It increases FRC.

D. It causes FRC to become non-calculable.

The correct choice highlights that Positive End-Expiratory Pressure (PEEP) increases Functional Residual Capacity (FRC). This occurs because PEEP maintains airway pressure at the end of expiration, thereby preventing alveolar collapse and enhancing lung recruitment. By increasing the volume of air remaining in the lungs at the end of expiration, PEEP effectively raises FRC. Higher FRC improves oxygenation and ventilation-perfusion matching, making it a valuable tool in the management of patients with conditions such as acute respiratory distress syndrome (ARDS) or other forms of hypoxemia. The increased FRC allows for a greater surface area available for gas exchange, aiding in improved oxygen delivery and carbon dioxide removal. Other options suggest a decrease in FRC, no effect, or an uncalculable status, which does not align with the physiological principles associated with PEEP. Therefore, understanding how PEEP interacts with lung mechanics is crucial for optimizing ventilatory support in clinical settings.

10. Which block provides a reliable block of the orbicularis oculi muscle for eye surgery?

A. Retrobulbar block

B. Sphenopalatine block

C. Peribulbar block

D. Topical anesthetic only

The peribulbar block is designed to anesthetize the globe and the surrounding structures, providing a reliable block of the orbicularis oculi muscle, which is critical for eye surgeries. This technique involves the injection of local anesthetic around the orbit, allowing for sensory and motor nerve blockade in this area, including the eyelid muscles. In eye surgery, particularly when muscle relaxation is essential, the peribulbar block effectively reduces intraoperative movement and provides adequate anesthesia for both muscle and sensory nerves, enabling a more controlled surgical environment. This method also minimizes the risk of intraocular complications compared to other techniques. The retrobulbar block, while it can also provide anesthesia for the eye, has associated risks such as intracranial injection and optic nerve damage, which may make it less favorable for specific procedures. The sphenopalatine block primarily targets the palate and does not provide adequate anesthesia to the eye region. Topical anesthetics may not provide sufficient depth of anesthesia or muscle block for surgical procedures, thus limiting their effectiveness compared to the peribulbar approach.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://anesthesiaakt24.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE