ASCP MOLECULAR BIOLOGY (MB) TECHNOLOGIST PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ascpmolecularbiology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the main application of Western Blotting in clinical diagnostics?

A. Detection of RNA levels

B. Cancer gene identification

C. Disease diagnosis through protein detection

D. Virus quantification

2. What can minor histocompatibility antigens (mHags) cause after a transplant?

A. Increased graft vs. tumor effects

B. Enhanced immune tolerance

C. Graft-vs-host disease and graft failure

D. Successful long-term graft acceptance

3. What is another term for a negative control in PCR?

A. Contamination control

B. Amplification control

C. Positive control

D. Sample control

4. What is the wobble effect in relation to tRNA?

A. The ability to bind only to specific codons

B. The tolerance of variations in the third base of codons

C. The mechanism that reduces mutation rates

D. The secondary structure of tRNA molecules

5. Which amplification method is included in target amplification methods?

A. Gel electrophoresis

B. Transcription-Based Amplification

C. Isolation techniques

D. Sequencing

SAMPLE

6. Multiplex Ligation-Dependent Probe Amplification (MLPA) is used to detect what type of mutational changes?

A. Point mutations

B. Copy number changes

C. Translocations

D. Sequence variations

7. What is the role of DNA Polymerase γ in eukaryotes?

A. Base excision repair

B. Mitochondrial DNA replication

C. Replication initiation

D. Leading strand synthesis

8. Which of the following oncogenes have tyrosine kinase activity?

A. EGFR

B. ALL of the above

C. BCR-Abl

D. HER2

9. What does Northern Blotting primarily analyze?

A. DNA sequences

B. Proteins

C. RNA sequences

D. gene mutations

10. In the context of gene expression analysis, which blotting technique is preferred to study mRNA levels?

A. Southern Blot

B. Northern Blot

C. Western Blot

D. PCR amplification

SAMPLE

Answers

SAMPLE

1. C
2. C
3. A
4. B
5. B
6. B
7. B
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What is the main application of Western Blotting in clinical diagnostics?

A. Detection of RNA levels

B. Cancer gene identification

C. Disease diagnosis through protein detection

D. Virus quantification

The main application of Western Blotting in clinical diagnostics is disease diagnosis through protein detection. This technique is widely used to identify specific proteins in a sample, which are often indicative of various medical conditions. For example, Western Blotting is crucial in the diagnosis of infectious diseases like HIV, where the presence of certain viral proteins can confirm infection. In addition to infectious diseases, Western Blotting can also be applied to detect autoantibodies in autoimmune disorders, as these antibodies target specific proteins in the body. The method is characterized by its specificity and ability to provide detailed information about protein expression, making it an important tool in the clinic for diagnosing diseases based on protein markers rather than nucleic acid levels or gene mutations. Other techniques serve different roles in molecular biology; for instance, the detection of RNA levels aligns more closely with techniques such as RT-PCR. Cancer gene identification typically involves methods like sequencing or other genomic techniques rather than protein detection. Virus quantification is adeptly addressed by techniques like quantitative PCR instead of Western Blotting, which focuses on proteins rather than viral RNA or DNA.

2. What can minor histocompatibility antigens (mHags) cause after a transplant?

A. Increased graft vs. tumor effects

B. Enhanced immune tolerance

C. Graft-vs-host disease and graft failure

D. Successful long-term graft acceptance

Minor histocompatibility antigens (mHags) play a significant role in transplant immunology, particularly in the context of mismatched grafts. These antigens can trigger immune responses because they are recognized as foreign by the recipient's immune system. When mHags are present and not matched between the donor and recipient, the recipient's immune cells can mount an attack against the transplanted tissue. This immune response can lead to graft-versus-host disease (GVHD), where the donor immune cells recognize the recipient's tissues as foreign, causing inflammation and damage. Furthermore, the reaction to mHags can also contribute to the potential failure of the graft due to rejection processes initiated by the recipient's immune system. In contrast to other options, the presence of mHags typically does not enhance immune tolerance or contribute positively to graft acceptance. Instead, they are more associated with adverse outcomes such as graft-versus-host disease and the risk of graft failure. Understanding the influence of minor histocompatibility antigens is crucial for improving transplant outcomes and managing patient care post-transplantation.

SAMPLE

3. What is another term for a negative control in PCR?

A. Contamination control

B. Amplification control

C. Positive control

D. Sample control

A negative control in PCR is often referred to as a contamination control. This type of control is essential for ensuring the specificity and accuracy of the PCR results. The purpose of a negative control is to confirm that any observed amplification is due to the target nucleic acid and not from contamination or non-specific binding. In practice, this means that a sample containing all the PCR reagents except the template DNA is run alongside the experimental samples. If amplification occurs in the negative control, it indicates contamination or issues with the reagents. While amplification control, positive control, and sample control serve important roles in a PCR experiment, they do not specifically refer to the absence of a target nucleic acid that defines a negative control. Amplification controls validate that the PCR conditions are suitable for amplifying DNA if present, while positive controls demonstrate that the system is capable of producing results when a known target is present. Sample control is typically associated with the assessment of the specimens being tested rather than serving as a baseline comparison for contamination.

4. What is the wobble effect in relation to tRNA?

A. The ability to bind only to specific codons

B. The tolerance of variations in the third base of codons

C. The mechanism that reduces mutation rates

D. The secondary structure of tRNA molecules

The wobble effect refers specifically to the flexible base-pairing that occurs at the third position of a codon in mRNA, allowing a single tRNA molecule to recognize multiple codons that code for the same amino acid. This phenomenon is particularly important because it helps to explain why there are fewer tRNA molecules than there are codons. The ability for the third nucleotide in a codon to form non-standard base pairs means that the genetic code can be efficiently translated without needing a unique tRNA for every codon, thus increasing the efficiency of protein synthesis. In the context of the options provided, the correct choice correctly identifies this tolerance for variations in the third base of codons, which is a hallmark of the wobble effect. This feature plays a critical role in maintaining the efficiency and fidelity of the translation process in protein synthesis.

SAMPLE

5. Which amplification method is included in target amplification methods?

A. Gel electrophoresis

B. Transcription-Based Amplification

C. Isolation techniques

D. Sequencing

Transcription-Based Amplification is a form of target amplification that specifically focuses on increasing the quantity of a particular nucleic acid target. This method involves the use of RNA polymerase to synthesize RNA from a DNA template, leading to a significant increase in the amount of target RNA. This amplification can then be further analyzed or detected using various techniques. Target amplification methods are crucial because they enable researchers and clinicians to detect low-abundance nucleic acids, making them essential in applications like viral load testing or the detection of pathogens in clinical samples. By amplifying the target, transcription-based methods ensure that there is enough material to analyze, allowing for increased sensitivity and specificity in molecular diagnostics. In contrast, gel electrophoresis serves as a separation technique, isolation techniques focus on obtaining specific molecules from a mixture, and sequencing is used to determine the exact sequence of nucleotides in a DNA or RNA molecule. While all these methods are important in molecular biology, they do not fall under the category of target amplification methods like Transcription-Based Amplification does.

6. Multiplex Ligation-Dependent Probe Amplification (MLPA) is used to detect what type of mutational changes?

A. Point mutations

B. Copy number changes

C. Translocations

D. Sequence variations

Multiplex Ligation-Dependent Probe Amplification (MLPA) is specifically designed to detect and quantify copy number changes in DNA, which includes deletions and duplications of genomic regions. This technique utilizes specific probes that bind to target sequences in the DNA, followed by a ligation step that joins adjacent probes. Once ligated, these products are amplified through PCR, allowing for the identification and quantification of specific DNA segments. The strength of MLPA lies in its ability to simultaneously analyze multiple target sequences, making it an efficient tool for assessing copy number variations across several genes or genomic loci in a single reaction. This is crucial in various clinical applications, such as diagnosing genetic disorders, detecting chromosomal abnormalities, and identifying potential cancer-related mutations. In contrast, point mutations are typically analyzed using sequencing techniques or specific mutation detection methods, translocations are generally evaluated using methods such as fluorescence in situ hybridization (FISH) or karyotyping, and sequence variations encompass changes in nucleotide sequences that might not result in copy number alterations. Therefore, the use of MLPA is particularly relevant to understanding and addressing copy number changes, confirming that this choice is the most accurate representation of what MLPA is intended to detect.

SAMPLE

7. What is the role of DNA Polymerase γ in eukaryotes?

A. Base excision repair

B. Mitochondrial DNA replication

C. Replication initiation

D. Leading strand synthesis

DNA Polymerase γ plays a crucial role in the replication of mitochondrial DNA in eukaryotic cells. This enzyme is specifically responsible for synthesizing the mitochondrial genome, which is distinct from nuclear DNA and is essential for the proper functioning of mitochondria. Since mitochondria have their own unique DNA, DNA Polymerase γ is tailored to meet their specific needs, facilitating both the replication of mitochondrial DNA and, to some extent, its repair. The focus on mitochondrial DNA highlights the importance of DNA Polymerase γ in supporting cellular energy production and metabolism, as mitochondria are the powerhouse of the cell. This specialization distinguishes it from other DNA polymerases in the cell that are involved in nuclear DNA processes such as repair, replication initiation, and leading strand synthesis, which are carried out by different polymerases with their own specialized functions.

8. Which of the following oncogenes have tyrosine kinase activity?

A. EGFR

B. ALL of the above

C. BCR-Abl

D. HER2

The correct answer is that all of the mentioned oncogenes exhibit tyrosine kinase activity. Tyrosine kinases are crucial players in cellular signaling pathways, particularly those related to growth and differentiation. Each of the oncogenes listed is known for its role in cancer through mechanisms involving tyrosine kinase activity. EGFR (epidermal growth factor receptor) is a receptor tyrosine kinase that, when activated by binding to its ligand, undergoes autophosphorylation. This phosphorylation activates several downstream signaling pathways that promote cell proliferation and survival, connecting it to various cancers. BCR-Abl is a fusion protein resulting from the Philadelphia chromosome translocation associated with chronic myeloid leukemia (CML). This fusion combines the breakpoint cluster region (BCR) gene with the Abelson murine leukemia viral oncogene (Abl), creating a constitutively active tyrosine kinase that leads to uncontrolled cell division. HER2 (human epidermal growth factor receptor 2) is another receptor tyrosine kinase that, when overexpressed, is implicated in breast cancer. Like EGFR, HER2 can trigger signaling cascades promoting cell division and survival through its tyrosine kinase activity. The presence of ty

9. What does Northern Blotting primarily analyze?

A. DNA sequences

B. Proteins

C. RNA sequences

D. gene mutations

Northern Blotting is a laboratory technique specifically designed to detect and analyze RNA molecules within a sample. This method involves the separation of RNA by gel electrophoresis, followed by transfer to a membrane and hybridization with a complementary labeled probe. The primary application of Northern Blotting is to assess the size, abundance, and expression levels of specific RNA transcripts, which provides valuable information regarding gene expression. By focusing on RNA rather than DNA or proteins, Northern Blotting enables researchers to study various aspects of RNA biology, including alternative splicing, developmental regulation, and responses to environmental signals. The technique is particularly useful for investigating gene expression profiles in different tissues or under varying conditions, thereby shedding light on the functional roles of specific genes.

SAMPLE

10. In the context of gene expression analysis, which blotting technique is preferred to study mRNA levels?

A. Southern Blot

B. Northern Blot

C. Western Blot

D. PCR amplification

The preferred blotting technique to study mRNA levels is the Northern Blot. This technique is specifically designed to detect and quantify RNA transcripts, allowing researchers to analyze gene expression by measuring the amount of specific mRNA in a sample. During a Northern Blot, RNA is extracted from cells or tissues and then separated by size through gel electrophoresis. The RNA is subsequently transferred to a membrane and hybridized with a labeled probe complementary to the target mRNA. The presence and intensity of the signal from the probe indicate the abundance of the specific mRNA, providing insights into gene expression levels. In contrast, other techniques mentioned serve different purposes: the Southern Blot is used for DNA analysis, primarily to detect specific DNA sequences; the Western Blot is focused on proteins, allowing for the detection and quantification of specific proteins in a sample; and PCR amplification is a method for amplifying specific segments of DNA, not RNA. Each of these techniques is suited to its respective type of molecule, making the Northern Blot the appropriate choice for mRNA study.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ascpmolecularbiology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE