ASCP SPECIALIST IN HEMATOLOGY (SH) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ascpspecialistinhematology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What type of blood cells are primarily affected in aplastic anemia?

A. White blood cells only

B. Red blood cells only

C. Platelets only

D. All types of blood cells (red cells, white cells, and platelets)

2. What happens if a sodium citrate sample is filled an inch below the fill line?

A. Unacceptable for coagulation studies

B. Acceptable for coagulation studies

C. Acceptable if additional blood can fill the tube

D. Acceptable if only the PT test is performed

3. Which test is used to differentiate monoblasts from myeloblasts?

A. Pappenheimer stain

B. Leukocyte alkaline phosphatase stain

C. Esterase stains

D. Hemoglobin electrophoresis

4. What is the significance of a high white blood cell count?

A. Indicates anemia

B. Suggests a possible infection or inflammation

C. Indicates dehydration

D. Suggests a state of hypoxia

5. Which organ is primarily responsible for recycling iron from old red blood cells?

A. Spleen

B. Liver

C. Kidneys

D. Bone marrow

SAMPLE

6. Which factor is primarily associated with the intrinsic pathway of coagulation?

A. Factor VII

B. Factor VIII

C. Factor X

D. Factor XI

7. In what type of patients would you expect to see a “left shift" in the white blood cell differential?

A. In patients experiencing chronic inflammation

B. In patients with allergies

C. In patients with acute infection or stress

D. In patients recovering from chemotherapy

8. What type of leukemia is characterized by the presence of promyelocytes with heavy granulation?

A. Acute myeloid leukemia

B. Acute promyelocytic leukemia

C. Chronic myeloid leukemia

D. Acute lymphoblastic leukemia

9. Which condition may lead to secondary thrombocytopenia?

A. Infections

B. Hemophilia

C. Polycythemia vera

D. Sideroblastic anemia

10. In patients with thalassemia, what is typically seen in red blood cells?

A. Increased MCV and decreased RDW

B. Increased RDW and normal MCV

C. Decreased MCV with increased RDW

D. Both increased MCV and RDW

SAMPLE

Answers

SAMPLE

1. D
2. A
3. C
4. B
5. A
6. B
7. C
8. B
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. What type of blood cells are primarily affected in aplastic anemia?

A. White blood cells only

B. Red blood cells only

C. Platelets only

D. All types of blood cells (red cells, white cells, and platelets)

Aplastic anemia is characterized by the failure of the bone marrow to produce sufficient blood cells, which can affect all hematopoietic lineages. The condition leads to a significant reduction in the production of red blood cells, white blood cells, and platelets, resulting in pancytopenia. In this disorder, the bone marrow is typically hypoplastic or even aplastic, which means it contains fewer blood-forming stem cells to produce all types of blood cells—erythrocytes (red blood cells), leukocytes (white blood cells), and thrombocytes (platelets). The lack of red blood cells can cause anemia, leading to fatigue and weakness. The decrease in white blood cells increases the risk of infections, while the reduction in platelets can lead to bleeding and bruising. Thus, the impact of aplastic anemia is systemic, affecting the production of all major blood components, making the correct answer encompass all types of blood cells.

2. What happens if a sodium citrate sample is filled an inch below the fill line?

A. Unacceptable for coagulation studies

B. Acceptable for coagulation studies

C. Acceptable if additional blood can fill the tube

D. Acceptable if only the PT test is performed

When a sodium citrate sample is not filled to the appropriate fill line, it can lead to an unacceptable blood to anticoagulant ratio. Sodium citrate acts as an anticoagulant, and the correct volume of blood is crucial for accurate coagulation results. An inch below the fill line significantly reduces the volume of blood relative to the amount of anticoagulant in the tube, which can artificially prolong coagulation times and lead to misleading test results. In coagulation studies, maintaining the correct blood to citrate ratio is essential to ensure that the clotting factors are adequately preserved for accurate testing. If the volume is insufficient, the anticoagulant will not function as intended, thus rendering the sample unsuitable for reliable testing. Consequently, the sample is deemed unacceptable for coagulation studies when it is not filled to the specified line.

3. Which test is used to differentiate monoblasts from myeloblasts?

A. Pappenheimer stain

B. Leukocyte alkaline phosphatase stain

C. Esterase stains

D. Hemoglobin electrophoresis

The differentiation of monoblasts from myeloblasts is an important aspect of diagnosing and categorizing hematological disorders, particularly acute leukemias. Esterase stains are employed for this purpose because they target specific enzymes present in different cell types. Monoblasts typically express non-specific esterase, which indicates their monocytic lineage, while myeloblasts show less activity for this enzyme. As a result, when a sample is subjected to esterase staining, the presence of non-specific esterase activity helps to identify monoblasts, thus facilitating the distinction from myeloblasts which may express different esterases or show different staining patterns. Other stains listed, despite their utility in different contexts, do not specifically aid in distinguishing between monoblasts and myeloblasts. For instance, the Pappenheimer stain is more associated with iron detection and identifying sideroblasts, while the leukocyte alkaline phosphatase stain assesses leukocyte activity but is not utilized for distinguishing between these two blast types. Hemoglobin electrophoresis is primarily employed to separate different hemoglobin types rather than for identifying specific white blood cell lineages.

SAMPLE

4. What is the significance of a high white blood cell count?

A. Indicates anemia

B. Suggests a possible infection or inflammation

C. Indicates dehydration

D. Suggests a state of hypoxia

A high white blood cell (WBC) count is particularly significant because it often suggests the presence of an infection or ongoing inflammation in the body. White blood cells are crucial components of the immune system, playing a key role in identifying and combating pathogens like bacteria and viruses. When the body detects an infection or inflammation, it typically increases the production of white blood cells to help counteract the threat, leading to a higher count observed in laboratory tests. In cases of infection, the elevation in white blood cells can be quite pronounced, manifesting as a leukocytosis where the body mobilizes neutrophils and other immune cells to the site of infection. Additionally, inflammatory conditions, such as autoimmune diseases or even stress responses, can also result in increased white blood cell counts as the body reacts to various stimuli. This is why a high WBC count is a critical indicator and often triggers further investigation to identify the underlying cause of the immune response. Anemia, dehydration, and hypoxia are associated with different physiological processes and do not directly correlate with an elevated white blood cell count. Instead, they involve other markers and clinical findings that guide their diagnosis and management. Therefore, understanding the implications of a high WBC count leads clinicians to investigate potential infections or

5. Which organ is primarily responsible for recycling iron from old red blood cells?

A. Spleen

B. Liver

C. Kidneys

D. Bone marrow

The spleen plays a critical role in the recycling of iron from old red blood cells. This organ acts as a filter for the blood, removing aged and damaged erythrocytes. When red blood cells reach the end of their life span, they are phagocytized by macrophages in the spleen, where hemoglobin is broken down. This process releases heme, which then undergoes further breakdown to release iron. The iron can then be stored for future use or transported back to the bone marrow for the production of new red blood cells. While other organs, such as the liver, are involved in the overall metabolism and storage of iron, they do not primarily handle the recycling of iron from old red blood cells. The kidneys focus on filtering blood and excreting waste but do not play a significant role in iron recycling. Bone marrow is crucial for the production of new red blood cells but is not involved in the recycling process of iron from older cells. This distinct function of the spleen underscores its importance in maintaining iron homeostasis and supporting hematopoiesis.

SAMPLE

6. Which factor is primarily associated with the intrinsic pathway of coagulation?

A. Factor VII

B. Factor VIII

C. Factor X

D. Factor XI

The correct choice highlights Factor VIII as a key component of the intrinsic pathway of coagulation. The intrinsic pathway is one of the two pathways that lead to blood coagulation, with the other being the extrinsic pathway. It is activated when blood comes into contact with negatively charged surfaces, such as exposed collagen in a damaged vessel. Factor VIII is specifically involved in the intrinsic pathway as it acts as a cofactor for Factor IXa, which in turn activates Factor X in the presence of calcium ions. This action is crucial for the proper functioning of the intrinsic pathway and ensures the formation of the clotting factor complexes necessary for effective hemostasis. Factors VII and X are part of the extrinsic and common pathways respectively. Factor XI is an additional component of the intrinsic pathway, but its role is secondary to that of Factor VIII, which is critical for the amplification of the coagulation cascade initiated by the intrinsic pathway. This distinction ultimately underlines why Factor VIII is primarily associated with this pathway.

7. In what type of patients would you expect to see a “left shift” in the white blood cell differential?

A. In patients experiencing chronic inflammation

B. In patients with allergies

C. In patients with acute infection or stress

D. In patients recovering from chemotherapy

A "left shift" in the white blood cell differential refers to the presence of an increased number of immature white blood cells, particularly neutrophils, in circulation. This phenomenon is commonly observed in conditions characterized by an acute inflammatory response, primarily due to infection or significant physiological stress. In cases of acute infection, the body rapidly responds to the presence of pathogens by increasing the production and release of white blood cells from the bone marrow. As the infection progresses, there is not only a rise in the total number of white blood cells but also a relative increase in immature forms, such as band neutrophils. This state signifies that the body is trying to mobilize its defenses quickly to combat the infectious agent, hence the term "left shift," which is a reference to how these cells are positioned on the white blood cell differential chart. For acute infection or stress, this response is essential as it indicates an active fight against potential threats, showcasing the body's efforts to maintain homeostasis. While chronic inflammation and allergies can lead to various changes in white blood cell counts and types, they typically do not present with a classic "left shift" as seen in acute processes. Similarly, during recovery from chemotherapy, the bone marrow is regaining its normal function, which may

SAMPLE

8. What type of leukemia is characterized by the presence of promyelocytes with heavy granulation?

A. Acute myeloid leukemia

B. Acute promyelocytic leukemia

C. Chronic myeloid leukemia

D. Acute lymphoblastic leukemia

Acute promyelocytic leukemia (APL) is characterized specifically by the presence of promyelocytes that exhibit heavy granulation. This type of leukemia is a subtype of acute myeloid leukemia (AML) and is particularly noted for its distinct morphological features, including the presence of the promyelocyte cells that have abundant cytoplasmic granules and often multiple Auer rods. In APL, these promyelocytes can also sometimes be found in bundles known as "faggot cells," which further helps in identifying this leukemia. The heavy granulation is a hallmark finding in cytogenetic and morphological examinations, which is critical for diagnosis. Additionally, APL is commonly associated with specific genetic abnormalities, particularly the translocation t(15;17), which leads to the fusion of the promyelocytic leukemia (PML) gene and the retinoic acid receptor alpha (RARA) gene, confirming its identity as a distinct clinical entity within the broader category of acute myeloid leukaemias. Understanding these features is important for hematopathologists and medical professionals in order to provide appropriate treatment strategies, including the use of all-trans retinoic acid (ATRA), which has dramatically improved outcomes for patients with this specific type

9. Which condition may lead to secondary thrombocytopenia?

A. Infections

B. Hemophilia

C. Polycythemia vera

D. Sideroblastic anemia

Secondary thrombocytopenia refers to a condition where platelet levels are decreased due to underlying factors or diseases, rather than from a primary bone marrow disorder. Infections can lead to secondary thrombocytopenia through various mechanisms, such as the destruction of platelets by the immune system, consumption of platelets in response to infections, or suppression of platelet production in the bone marrow due to inflammatory cytokines. Infections can induce a significant immune response, leading to thrombocytopenia. For instance, viral infections (like HIV and hepatitis) are known to affect platelet production and survival. Additionally, sepsis can cause thrombocytopenia through mechanisms such as disseminated intravascular coagulation (DIC), where platelets are consumed excessively. Other options do not typically lead to secondary thrombocytopenia. Hemophilia primarily affects the coagulation pathway and leads to bleeding tendencies rather than directly impacting platelet levels. Polycythemia vera is a myeloproliferative disorder characterized by an increase in red blood cells and usually presents with normal or elevated platelet counts. Sideroblastic anemia is primarily an issue of ineffective erythropoiesis and iron metabolism, which does not commonly result in thrombocytopenia. Thus, infections are the

SAMPLE

10. In patients with thalassemia, what is typically seen in red blood cells?

A. Increased MCV and decreased RDW

B. Increased RDW and normal MCV

C. Decreased MCV with increased RDW

D. Both increased MCV and RDW

In thalassemia, the hallmark red blood cell changes include a decreased mean corpuscular volume (MCV) and an increased red cell distribution width (RDW). Thalassemia is characterized by ineffective erythropoiesis and microcytic anemia, which results in smaller than normal red blood cells, thus leading to a reduced MCV. Additionally, the increased RDW reflects the variability in red blood cell sizes present in thalassemia, which occurs due to the presence of both normal and smaller cells produced by the bone marrow. This variability is significant because it indicates a mixture of red blood cell populations, often seen in genetically inherited conditions like thalassemia. The other options presented do not align with the typical hematological findings in thalassemia. Increased MCV would suggest macrocytic anemia, which is not characteristic of this condition, while a decreased RDW might indicate a more homogenous population of red blood cells rather than the variation seen in thalassemia.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ascpspecialistinhematology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE