ASSOCIATE DEGREE IN NURSING (ADN) LEVEL 3 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://adultnursinglvl3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the significance of a hard, rigid abdomen in the postoperative client?

A. It indicates normal healing

B. It suggests the presence of peritonitis

C. It is a common postoperative finding

D. It indicates muscle spasm

2. When evaluating patients for potential complications, which clinical presentation requires urgent intervention?

A. A client with an anxiety attack before surgery

B. A client showing signs of superficial edema

C. A client with rigid abdomen indicating possible peritonitis

D. A client rating postoperative pain as an "8"

3. Which electrolyte imbalance is most commonly associated with cardiac arrhythmias?

A. Calcium imbalance

B. Sodium imbalance

C. Magnesium imbalance

D. Potassium imbalance

4. If a nurse finds a client with a "do not resuscitate" order unresponsive, what should the nurse do first?

A. Page the client's physician.

B. Push the emergency alarm to call a code.

C. Pull the curtain and leave the room.

D. Stay in the room and call the nursing team for assistance.

5. What is one of the most important aspects of patient education regarding medication side effects?

A. Patients should never discuss side effects with other healthcare providers

B. Patients should be advised to stop taking medications if side effects occur

C. Patients should be informed about potential side effects and when to seek help

D. Patients need to be reassured that all medications are risk-free

SAMPLE

6. Which patient complaint should lead the nurse to perform a thorough assessment first?

A. Complaints of nausea and vomiting

B. Intense abdominal pain rating "8"

C. Mild discomfort at the incision site

D. General fatigue after surgery

7. What does a nurse assess when checking for peripheral pulses?

A. Heart rate

B. Circulation and blood flow to the extremities

C. Skin temperature

D. Capillary refill time

8. In what patient population is polypharmacy most frequently seen?

A. Elderly patients

B. Teenagers

C. Newborns

D. Middle-aged adults

9. What statement by the client at 32 weeks' gestation indicates an understanding of when to call the physician?

A. "If I see any bleeding, even if I have no pain."

B. "If I experience leg cramps at night."

C. "If my baby seems more active than usual."

D. "If I have a pounding headache that goes away."

10. How often should vital signs be assessed for a post-operative patient?

A. Every 30 minutes for the first hour

B. Every 15 minutes for the first hour

C. Every hour for the first 4 hours

D. Every 10 minutes for the first 30 minutes

SAMPLE

Answers

SAMPLE

1. B
2. C
3. D
4. D
5. C
6. B
7. B
8. A
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What is the significance of a hard, rigid abdomen in the postoperative client?

A. It indicates normal healing

B. It suggests the presence of peritonitis

C. It is a common postoperative finding

D. It indicates muscle spasm

A hard, rigid abdomen in a postoperative client is significant because it suggests the presence of peritonitis, a serious condition that can occur following surgical procedures. Peritonitis is the inflammation of the peritoneum, which can result from various factors, including infection, leakage of intestinal contents, or lateral complications from the surgery, such as perforation or abscess formation. The rigid nature of the abdomen indicates increased intra-abdominal pressure and may reflect an underlying inflammatory process. In the postoperative setting, this finding should raise concern about potential complications that require immediate assessment and intervention, as untreated peritonitis can lead to severe consequences, such as sepsis or organ failure. In contrast, normal healing typically presents with a soft and non-tender abdomen, while muscle spasms may cause localized tightness but do not generally result in a hard, rigid abdomen. Additionally, while some changes to abdominal firmness can occur post-surgery, a rigid abdomen is not considered a common postoperative finding and warrants further investigation to rule out significant complications.

2. When evaluating patients for potential complications, which clinical presentation requires urgent intervention?

A. A client with an anxiety attack before surgery

B. A client showing signs of superficial edema

C. A client with rigid abdomen indicating possible peritonitis

D. A client rating postoperative pain as an "8"

The clinical presentation denoted by a client with a rigid abdomen is highly indicative of potential peritonitis, which is a serious condition that occurs when the peritoneum, a membrane lining the abdominal wall and covering the abdominal organs, becomes inflamed. This inflammation may be due to a variety of causes, including infection from perforation of an abdominal organ, and can lead to severe complications such as sepsis. In such cases, the rigidity of the abdomen suggests that there may be an underlying infection or another critical issue requiring immediate medical assessment and intervention. Untreated peritonitis can rapidly progress to a life-threatening condition, making it essential for healthcare providers to act swiftly to stabilize the patient, conduct necessary diagnostic tests, and initiate treatment. In contrast, while anxiety attacks, superficial edema, and a high pain rating following surgery are concerning and warrant attention, they do not present the same level of immediate risk to the patient's life as the rigid abdomen associated with potential peritonitis. Addressing these other issues is important for patient comfort and management, but they do not necessitate the same urgency as a rigid abdomen would.

SAMPLE

3. Which electrolyte imbalance is most commonly associated with cardiac arrhythmias?

A. Calcium imbalance

B. Sodium imbalance

C. Magnesium imbalance

D. Potassium imbalance

The association between potassium imbalance and cardiac arrhythmias is well-established in medical literature. Potassium is a critical electrolyte that plays a vital role in maintaining the electrical conductivity of the heart. It is essential for the proper functioning of cardiac myocytes (heart muscle cells) and helps regulate the heart's rhythm. When potassium levels are either too high (hyperkalemia) or too low (hypokalemia), it can significantly disrupt the electrical impulses in the heart, leading to various types of arrhythmias. For example, high potassium levels can cause phase 4 depolarization to be affected, leading to slower heart rates or even cardiac arrest in severe cases. On the other hand, low potassium levels can lead to increased myocardial excitability and the potential for arrhythmias like premature ventricular contractions and even more serious issues like ventricular fibrillation. While calcium, sodium, and magnesium are also important electrolytes and can influence cardiac function, potassium's direct effect on the heart's electrical conduction makes it the most commonly associated electrolyte with cardiac arrhythmias. This is why, among the various electrolyte imbalances, potassium imbalance is often highlighted in the context of cardiac health and arrhythmias.

4. If a nurse finds a client with a "do not resuscitate" order unresponsive, what should the nurse do first?

A. Page the client's physician.

B. Push the emergency alarm to call a code.

C. Pull the curtain and leave the room.

D. Stay in the room and call the nursing team for assistance.

In a situation where a nurse discovers a client who is unresponsive and has a "do not resuscitate" (DNR) order, the most appropriate action is to stay in the room and call the nursing team for assistance. This response indicates that the nurse acknowledges the client's wishes as stated in the DNR order, which means that resuscitation attempts should not be initiated. By staying in the room, the nurse can monitor the client's condition and ensure that they provide supportive care as needed, respecting the client's dignity and decisions. Calling for the nursing team ensures that there is an adequate response to the situation and that other healthcare professionals are informed, enabling them to assist as necessary without taking unnecessary measures that contradict the DNR order. The other options would not be appropriate in this context. For example, contacting the physician or calling a code could lead to interventions that go against the client's established wishes. Pulling the curtain and leaving the room would neglect the duty of care the nurse has towards the client, both ethically and professionally. Therefore, remaining with the client and seeking assistance from the nursing team is the best course of action to respect the DNR order while ensuring the client is not left unattended.

SAMPLE

5. What is one of the most important aspects of patient education regarding medication side effects?

A. Patients should never discuss side effects with other healthcare providers

B. Patients should be advised to stop taking medications if side effects occur

C. Patients should be informed about potential side effects and when to seek help

D. Patients need to be reassured that all medications are risk-free

Providing patients with comprehensive information about potential side effects of their medications, as well as guidance on when to seek help, is essential for promoting safe medication use and empowering patients to manage their health. When patients are aware of possible side effects, they can more readily identify any adverse reactions that may occur and understand the importance of reporting these issues to their healthcare providers promptly. This knowledge enables patients to differentiate between normal side effects and serious reactions that require immediate medical attention. By fostering an open dialogue about side effects, patients can feel more comfortable communicating with their healthcare team about their experiences and concerns, leading to better health outcomes and adherence to their medication regimen. Proper education serves to demystify the medication process, ensuring that patients are both informed and engaged in their own care.

6. Which patient complaint should lead the nurse to perform a thorough assessment first?

A. Complaints of nausea and vomiting

B. Intense abdominal pain rating "8"

C. Mild discomfort at the incision site

D. General fatigue after surgery

The most acute and concerning complaint among those listed is intense abdominal pain rated as "8". This level of pain indicates a significant distress that may suggest serious underlying issues, such as an injury, a rupture, or a complication that could require immediate intervention. In a clinical setting, a high pain rating acts as a crucial indicator for the need for prompt assessment and potential treatment. Intense abdominal pain can have a wide range of causes that may necessitate urgent care, including but not limited to post-operative complications like hemorrhage, infection, or bowel obstruction. Swift evaluation of the patient's condition can help determine the necessary interventions to prevent further deterioration and manage pain effectively. In contrast, although complaints of nausea and vomiting may indicate a need for assessment, they are typically less urgent than the severe abdominal pain. Mild discomfort at the incision site or general fatigue after surgery, while important to monitor, are less immediate in nature compared to a patient experiencing severe abdominal pain. Attending to the most pressing complaints first ensures that nursing care prioritizes patient safety and addresses the most critical needs accordingly.

SAMPLE

7. What does a nurse assess when checking for peripheral pulses?

A. Heart rate

B. Circulation and blood flow to the extremities

C. Skin temperature

D. Capillary refill time

When a nurse assesses peripheral pulses, the primary focus is on circulation and blood flow to the extremities. Peripheral pulses are the pulsations of arteries in the limbs, and their strength, rhythm, and quality provide crucial information regarding the vascular status of the extremities. By palpating these pulses, the nurse can determine if there is adequate blood flow reaching the arms and legs, which is essential for tissue perfusion. Monitoring peripheral pulses helps identify conditions such as peripheral artery disease, blood clots, or vascular occlusion, which can compromise circulation. Observations of the pulse's characteristics, combined with other assessments, can indicate the overall effectiveness of the cardiovascular system. Although heart rate can be indirectly inferred from peripheral pulse assessment, its main purpose is to assess the adequacy of circulation rather than solely measuring how fast the heart beats. Similarly, skin temperature and capillary refill time are useful peripheral circulation assessments but are separate evaluations that can complement the overall understanding of circulation status.

8. In what patient population is polypharmacy most frequently seen?

A. Elderly patients

B. Teenagers

C. Newborns

D. Middle-aged adults

Polypharmacy, which refers to the use of multiple medications by a patient, is most frequently seen in elderly patients. This tendency is largely attributable to several factors inherent to this population. As individuals age, they often develop multiple chronic conditions such as hypertension, diabetes, and arthritis, which typically require various treatments. Additionally, the elderly may visit multiple healthcare providers who prescribe separate medications that can lead to overlapping treatments or drug interactions. Moreover, age-related physiological changes can influence how the body processes medications, increasing the risk of adverse effects and complicating management. Because older adults are more likely to experience issues related to compliance, side effects, and potential drug interactions, the phenomenon of polypharmacy becomes particularly concerning in this demographic. In contrast, other populations such as teenagers, newborns, and middle-aged adults generally do not have the same prevalence of chronic diseases or the complex health needs that typically result in polypharmacy. This contributes to the observed higher rates of polypharmacy in the elderly, making this group the most frequently affected by this issue.

SAMPLE

9. What statement by the client at 32 weeks' gestation indicates an understanding of when to call the physician?

A. "If I see any bleeding, even if I have no pain."

B. "If I experience leg cramps at night."

C. "If my baby seems more active than usual."

D. "If I have a pounding headache that goes away."

The statement indicating an understanding of when to call the physician is the one regarding bleeding, even in the absence of pain. During pregnancy, particularly in the third trimester, any instance of bleeding is considered significant and warrants immediate medical attention. This is crucial as it may point to potential complications such as placental abruption, placenta previa, or other serious issues that can affect both the mother and the fetus. Recognizing this symptom and having a proactive approach to seek medical advice demonstrates awareness of the warning signs that could signify serious health concerns. Other statements reflect various concerns, but they do not indicate the same level of urgency. For instance, leg cramps at night can be a common and usually benign issue during pregnancy; it's often related to changes in blood flow and can typically be managed with simple interventions like stretching. Increased fetal movement, while noteworthy, does not always indicate a problem and should be evaluated in conjunction with overall patterns of activity. A headache that resolves on its own may be common in pregnancy, but a pounding headache that persists could require evaluation—however, without more context indicating severity or duration, it's not as critical as the indication of bleeding.

10. How often should vital signs be assessed for a post-operative patient?

A. Every 30 minutes for the first hour

B. Every 15 minutes for the first hour

C. Every hour for the first 4 hours

D. Every 10 minutes for the first 30 minutes

For a post-operative patient, assessing vital signs every 15 minutes for the first hour is considered the appropriate standard of care. This frequency is crucial because the post-operative period is a time when patients are at increased risk for complications such as bleeding, respiratory distress, or changes in vital signs that may indicate instability or distress. By monitoring vital signs at this interval, healthcare providers can quickly detect any changes that might require immediate intervention, ensuring the safety and well-being of the patient. Following the initial hour, the frequency can typically be reduced, allowing for a structured and capable approach to post-operative care while still remaining vigilant. The other choices suggest intervals that may not provide adequate monitoring during this critical initial recovery phase. For instance, assessing every 30 minutes or hourly may allow for too much time to pass without checking the patient, potentially missing early signs of complications.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://adultnursinglvl3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE