ASSOCIATION OF SOCIAL WORK BOARDS (ASWB) CLINICAL PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://aswb-clinical.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 9

Explanations 11

Next Steps 17

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **If a client shows symptoms of physical illness but is rooted in emotional issues, this could indicate:**

   A. Somatization

   B. Anxiety disorders

   C. Conversion disorder

   D. Adjustment disorder

2. **When engaging in physical contact with clients, what must be true?**

   A. Contact must be with adult clients only

   B. Contact can occur if it causes no psychological harm

   C. Contact must be culturally sensitive

   D. Rules for contact must be clearly articulated

3. **A client prescribed Nardil for depression is likely required to:**

   A. Get regular blood monitoring

   B. Limit certain foods high in tyramine

   C. Watch her caloric intake

   D. Have her blood pressure taken frequently

4. **A client shows visible distress discussing their family. When the worker notes the client's discomfort, the client claims "everything is fine". This is an example of:**

   A. Lack of congruence in communication

   B. Denial of psychic conflict

   C. Desire to please the social worker

   D. Symptomatology of a psychiatric diagnosis

5. **Which assessment tool should be used to determine a client's global functioning according to the DSM-5?**

   A. Global Assessment of Functioning Scale (GAF)

   B. World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0)

   C. Myers-Briggs Type Indicator

   D. Thematic Apperception Test

SAMPLE

6. Which of the following Personality Disorders is clustered with Antisocial Personality Disorder in a grouping characterized by dramatic, emotional, and erratic behavior?

A. Schizoid Personality Disorder

B. Narcissistic Personality Disorder

C. Obsessive-Compulsive Personality Disorder

D. Paranoid Personality Disorder

7. Which of the following are stages of group development in sequential order?

A. Preaffiliation, differentiation, intimacy, power and control, and separation/termination.

B. Differentiation, power and control, preaffiliation, intimacy, and separation/termination.

C. Preaffiliation, power and control, intimacy, differentiation, and separation/termination.

D. Differentiation, preaffiliation, power and control, intimacy, and separation/termination.

8. Which of the following is NOT one of the therapeutic advantages of injectable antipsychotic medications?

A. They reduce medication noncompliance because they are needed less frequently than oral medications.

B. They are less likely to be used for suicide or result in overdose because the dosing is controlled.

C. They are associated with decreased hospitalizations due to fewer symptom relapses.

D. They are less expensive than oral medications.

9. What defines delirium tremens?

A. Genetic conditions that can cause intellectual disabilities

B. Physical symptoms associated with alcohol withdrawal

C. Psychotic features linked with certain mental health diagnoses

D. Neurological deterioration that can occur in older adults

SAMPLE

10. Which of the following is a condition for further study in the DSM-5?

A. Sexual Masochism Disorder.

B. Caffeine Use Disorder.

C. Frotteuristic Disorder.

D. Pica.

SAMPLE

Answers

SAMPLE

1. C
2. C
3. B
4. A
5. B
6. B
7. C
8. D
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. If a client shows symptoms of physical illness but is rooted in emotional issues, this could indicate:

A. Somatization

B. Anxiety disorders

C. Conversion disorder

D. Adjustment disorder

The scenario described—where a client exhibits physical symptoms that are fundamentally linked to emotional issues—aligns closely with the concept of conversion disorder. This disorder involves the presence of neurological symptoms that cannot be explained by medical conditions, and these symptoms are typically triggered by psychological factors such as stress or emotional conflict. In conversion disorder, clients may present with symptoms like paralysis, weakness, or sensory deficits. These manifestations are not intentional and are a way for the individual to express psychological distress. Understanding this connection helps in recognizing that the physical ailments are not merely medical but are instead stemming from deeper emotional problems. The other options consider various disorders, with some related to emotional distress manifesting physically, but they do not specifically address the mechanism whereby psychological conflict leads to a significant change in motor or sensory functioning, which is characteristic of conversion disorder. Thus, this makes conversion disorder the most appropriate choice in the context provided.

2. When engaging in physical contact with clients, what must be true?

A. Contact must be with adult clients only

B. Contact can occur if it causes no psychological harm

C. Contact must be culturally sensitive

D. Rules for contact must be clearly articulated

When engaging in physical contact with clients, it is essential that the contact is culturally sensitive. This means recognizing and respecting the diverse cultural backgrounds of clients, as well as their individual beliefs and practices regarding physical touch. Cultural sensitivity in social work practice is crucial because different cultures have varying norms concerning physical interaction. Some clients may view physical contact, such as hugs or handshakes, as supportive and affirming, while others might find it intrusive or uncomfortable. Understanding these cultural nuances helps to build trust and rapport between the client and the social worker. When practitioners prioritize cultural sensitivity, they demonstrate respect for the client's identity, which can enhance the therapeutic relationship and increase the effectiveness of interventions. This consideration goes beyond mere adherence to guidelines; it involves a commitment to understanding the client's perspective and responding accordingly. In contrast, the other options may lack the comprehensive approach needed to ensure that physical contact is appropriate and beneficial. For example, restricting contact to adult clients does not address the complexities of individual client needs, and stating that contact should cause no psychological harm could be subjective without clear guidelines. While articulating rules for contact is important, it does not encompass the broader necessity of understanding and adapting to the client's cultural context. Ultimately, prioritizing cultural sensitivity ensures that social

SAMPLE

3. A client prescribed Nardil for depression is likely required to:

A. Get regular blood monitoring

B. Limit certain foods high in tyramine

C. Watch her caloric intake

D. Have her blood pressure taken frequently

The recommendation to limit certain foods high in tyramine is crucial for clients prescribed Nardil, which is a monoamine oxidase inhibitor (MAOI) used in the treatment of depression. MAOIs work by inhibiting the enzyme monoamine oxidase, which breaks down neurotransmitters like serotonin and norepinephrine, contributing to mood elevation. However, this inhibition also affects the breakdown of tyramine, a naturally occurring amino acid found in various foods. When a client on Nardil consumes foods high in tyramine, it can lead to a dangerous increase in blood pressure, known as a hypertensive crisis. Symptoms of a hypertensive crisis include severe headache, neck stiffness, nausea, and can even lead to stroke or other serious health complications. Therefore, it's imperative for clients on Nardil to avoid foods such as aged cheeses, cured meats, fermented products, and certain alcoholic beverages to prevent such adverse reactions. While options mentioning blood monitoring or blood pressure checks may seem relevant, they are more closely associated with other types of medications or conditions rather than the specific dietary restrictions required for those on MAOIs like Nardil. Monitoring caloric intake is also not a primary concern related to Nardil therapy, as the critical issue is

4. A client shows visible distress discussing their family. When the worker notes the client's discomfort, the client claims "everything is fine". This is an example of:

A. Lack of congruence in communication

B. Denial of psychic conflict

C. Desire to please the social worker

D. Symptomatology of a psychiatric diagnosis

The situation described highlights a lack of congruence in communication. Congruence refers to the alignment between verbal and non-verbal communication. In this scenario, the client's observable distress contradicts their verbal assertion that "everything is fine." This discrepancy indicates that, while the client is expressing one thing verbally, their emotional state and body language tell a different story. Lack of congruence can often arise from various underlying issues, such as fear of vulnerability, a desire to avoid discussing uncomfortable topics, or even internalized stigma regarding mental health issues. Understanding this conflict is crucial for the social worker, as it can provide insight into the client's emotional struggles and pave the way for deeper therapeutic conversation. Other options, while relevant to various psychological concepts, do not accurately describe this particular scenario. For example, denial of psychic conflict suggests a broader avoidance of internal struggles rather than a specific mismatch in communication. The desire to please the social worker might explain the client's denial but does not fully encapsulate the observed dissonance. Lastly, symptomatology of a psychiatric diagnosis would imply a clear-cut set of symptoms associated with a specific disorder, which does not directly pertain to the phenomenon of mismatched verbal and non-verbal cues in this context

SAMPLE

5. Which assessment tool should be used to determine a client's global functioning according to the DSM-5?

A. Global Assessment of Functioning Scale (GAF)

B. World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0)

C. Myers-Briggs Type Indicator

D. Thematic Apperception Test

The World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0) is the most appropriate assessment tool for determining a client's global functioning according to the DSM-5. This tool provides a comprehensive framework for measuring disability across various dimensions of functioning, including cognition, mobility, self-care, and interpersonal relationships. It is designed to assess the impact of health conditions on functional activities and is aligned with the biopsychosocial model, which encompasses biological, psychological, and social factors contributing to a person's overall functioning. WHODAS 2.0 captures both the severity of disability and the individual's perceived ability to function in daily life, making it a valuable asset for clinicians in assessing global functioning. Its standardized format allows for consistent and reliable evaluation across different populations and settings, thus helping clinicians to tailor treatment approaches and monitor progress over time. Other assessment tools listed, such as the Global Assessment of Functioning Scale (GAF), while previously utilized, are not used in the DSM-5. The Myers-Briggs Type Indicator is focused on personality types rather than functioning. The Thematic Apperception Test is a projective psychological test that explores themes and emotions rather than providing a measure of global functioning.

6. Which of the following Personality Disorders is clustered with Antisocial Personality Disorder in a grouping characterized by dramatic, emotional, and erratic behavior?

A. Schizoid Personality Disorder

B. Narcissistic Personality Disorder

C. Obsessive-Compulsive Personality Disorder

D. Paranoid Personality Disorder

The correct answer is Narcissistic Personality Disorder, which, along with Antisocial Personality Disorder, is part of the dramatic, emotional, and erratic cluster of personality disorders (Cluster B). This cluster is characterized by an array of symptoms and behaviors that demonstrate unstable emotions, interpersonal relationships, and self-image. Narcissistic Personality Disorder involves patterns of grandiosity, a need for admiration, and a lack of empathy, all of which can lead to erratic and dramatic interactions with others. Individuals with this disorder might engage in manipulative or exploitative behavior, similar to those diagnosed with Antisocial Personality Disorder, increasing the dramatic and emotional nature of their actions. The other personality disorders listed are classified differently. Schizoid Personality Disorder is characterized by a pervasive pattern of detachment from social relationships and a restricted range of emotional expression, placing it in Cluster A, which encompasses odd or eccentric behaviors. Obsessive-Compulsive Personality Disorder entails a preoccupation with orderliness, perfectionism, and control, making it a part of Cluster C, associated with anxious and fearful behaviors. Lastly, Paranoid Personality Disorder is also a Cluster A disorder, characterized by pervasive distrust and suspicion of others, markedly different from the dramatic and emotional qualities of Cluster

SAMPLE

7. Which of the following are stages of group development in sequential order?

A. Preaffiliation, differentiation, intimacy, power and control, and separation/termination.

B. Differentiation, power and control, preaffiliation, intimacy, and separation/termination.

C. Preaffiliation, power and control, intimacy, differentiation, and separation/termination.

D. Differentiation, preaffiliation, power and control, intimacy, and separation/termination.

The correct response details the stages of group development in a specific sequential order that aligns with established theories in social work and group dynamics. The stages identified—preaffiliation, power and control, intimacy, differentiation, and separation/termination—represent a journey a group typically undergoes as it forms and matures. Initially, in the preaffiliation stage, group members are often focused on getting to know each other and establishing trust. This is followed by the power and control phase, where conflicts may arise as individuals assert their needs and desires, leading to power dynamics within the group. The intimacy stage is characterized by deepening relationships and a sense of closeness among members, fostering collaboration and support. After the intimacy phase, groups move toward differentiation, in which members start to recognize and honor individual differences while still functioning cohesively as a group. Finally, the separation or termination stage marks the dissolution of the group or the end of its purpose, where members reflect on their experiences and gather closure. By adhering to this sequence, the answer exemplifies a comprehensive understanding of how these stages interconnect and contribute to the overall dynamics and development of a group in a clinical or social work context.

8. Which of the following is NOT one of the therapeutic advantages of injectable antipsychotic medications?

A. They reduce medication noncompliance because they are needed less frequently than oral medications.

B. They are less likely to be used for suicide or result in overdose because the dosing is controlled.

C. They are associated with decreased hospitalizations due to fewer symptom relapses.

D. They are less expensive than oral medications.

The therapeutic advantages of injectable antipsychotic medications primarily revolve around their effectiveness in promoting adherence to treatment, minimizing the risks of overdose, and ultimately reducing the need for hospitalizations due to symptom relapse. Injectable forms of medication provide a more consistent therapeutic level in the bloodstream, which can enhance compliance as patients do not have to remember daily doses. The first three options highlight significant benefits: reducing medication noncompliance addresses a common concern in mental health treatment, as the injectable form decreases the frequency of administration. Furthermore, by having controlled dosing, there is a diminished risk of overdose, making these medications safer in acute scenarios. Additionally, the management of symptoms can lead to fewer episodes requiring hospitalization, contributing to overall better health outcomes. In contrast, the idea that injectable antipsychotics are less expensive than oral medications is not inherently true. While oral medications may come with ongoing costs due to daily dosing, injectable medications can involve higher up-front costs related to the administration, healthcare professional involvement, and potential need for special storage and handling. Thus, cost does not reflect a therapeutic advantage but rather a financial consideration, making it the correct identification of what is not a therapeutic advantage in the context of injectable antipsychotic medications.

SAMPLE

9. What defines delirium tremens?

A. Genetic conditions that can cause intellectual disabilities

B. Physical symptoms associated with alcohol withdrawal

C. Psychotic features linked with certain mental health diagnoses

D. Neurological deterioration that can occur in older adults

Delirium tremens is characterized by severe physical symptoms that can arise during alcohol withdrawal. It typically presents 48 to 72 hours after the last drink in individuals with a history of heavy alcohol use. Key symptoms include intense agitation, confusion, hallucinations, and autonomic instability, which can manifest as rapid heartbeat and elevated blood pressure. The distinction between delirium tremens and other conditions is crucial for proper diagnosis and treatment. While the other options refer to various medical or psychosocial issues—such as genetic conditions, psychotic features of mental disorders, or neurological degeneration in older persons—delirium tremens specifically pertains to the physiological response to the abrupt cessation of alcohol intake, making it a life-threatening condition that requires immediate medical attention and intervention.

10. Which of the following is a condition for further study in the DSM-5?

A. Sexual Masochism Disorder.

B. Caffeine Use Disorder.

C. Frotteuristic Disorder.

D. Pica.

Caffeine Use Disorder is classified as a condition that requires further study in the DSM-5, meaning that it is recognized as a potential issue but is not yet fully validated as a formal psychiatric diagnosis. This designation indicates that the symptoms and criteria for the disorder need more empirical research to establish their reliability and validity. The DSM-5 takes a nuanced approach to mental health and substance issues, acknowledging that certain conditions may be prevalent in the population but lack sufficient evidence to warrant formal categorization as mental disorders. In the case of Caffeine Use Disorder, while caffeine consumption is widespread and can lead to problematic use, the current understanding and research on its impact and clinical significance are still evolving. In contrast, the other disorders listed, such as Sexual Masochism Disorder, Frotteuristic Disorder, and Pica, have more established diagnostic criteria and are recognized under the DSM-5 as distinct mental health issues with clearer understanding and implications for treatment.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://aswb-clinical.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE