
ATI NURSING CARE OF CHILDREN PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://atinursingcareofchildren.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which method is suggested for treating diaper rash?

A. Cover with talcum powder

B. Use an excessive amount of soap

C. Expose to air and use a zinc oxide barrier

D. Keep the area constantly wet

2. What is a late sign of head injury in children?

A. Delayed or impaired pupillary responses

B. High-pitched cry

C. Pallor and irritability

D. Increased sleepiness

3. What is a common physical change in school-age children as they begin to develop a sense of industry?

A. Increased anxiety

B. Improvements in cooperative play

C. Greater need for privacy

D. Decrease in friendships

4. What is a key difference between viral and bacterial meningitis?

A. Viral often resolves with supportive care, while bacterial is more dangerous

B. Viral meningitis requires hospitalization, but bacterial does not

C. Both types present with similar symptoms and treatment

D. Viral meningitis only occurs in infants and children

5. How long does the Babinski reflex typically last in infants?

A. Until 3 months

B. Usually until a year

C. Until 18 months

D. Until 2 years

6. What is the expected pulse rate of a child aged 3 months to 2 years?

A. 60 to 100/min

B. 70 to 150/min

C. 80 to 120/min

D. 90 to 130/min

SAMPLE

7. What is a common complication associated with cleft palate that may require specialist intervention?

A. Speech difficulties

B. Dental problems

C. Ear infections and hearing loss

D. Gastroesophageal reflux disease

8. What is often a sign of a ruptured appendix in children?

A. Improved pain

B. Severe diarrhea

C. High fever

D. Absence of vomiting

9. What type of play is common for toddlers facing hospitalization?

A. Cooperative play

B. Parallel play

C. Constructive play

D. Associative play

10. Which of the following is true regarding the inheritance of cystic fibrosis?

A. It follows an autosomal dominant inheritance pattern

B. Both parents must carry the recessive trait

C. It is inherited from the father only

D. It only affects males

SAMPLE

Answers

SAMPLE

1. C
2. A
3. B
4. A
5. B
6. B
7. C
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which method is suggested for treating diaper rash?

A. Cover with talcum powder

B. Use an excessive amount of soap

C. Expose to air and use a zinc oxide barrier

D. Keep the area constantly wet

Using exposure to air along with a zinc oxide barrier is an effective method for treating diaper rash because it addresses both the moisture and the physical irritation that can exacerbate the condition. Diaper rash often occurs due to prolonged exposure to moisture, friction, and irritants from urine and feces. By allowing the affected area to be exposed to air, the moisture that contributes to the rash can evaporate, promoting a healing environment. Additionally, applying a zinc oxide barrier serves multiple purposes; it not only provides a protective layer that shields the skin from further irritation from urine and feces but also has soothing properties that can reduce inflammation and promote healing of the skin. The other methods may pose risks or not provide the intended relief. For example, using talcum powder can sometimes exacerbate problems by contributing to respiratory issues if inhaled and may not effectively address the underlying moisture problem. Excessive soap usage can strip natural oils from the skin, leading to further irritation instead of relief. Keeping the area constantly wet would create an environment that can worsen the rash rather than allowing it to dry and recover.

2. What is a late sign of head injury in children?

A. Delayed or impaired pupillary responses

B. High-pitched cry

C. Pallor and irritability

D. Increased sleepiness

Delayed or impaired pupillary responses serve as a late sign of head injury in children because they indicate significant neurological impairment. When there is an injury to the brain or skull, it can affect the functioning of cranial nerves, particularly the third cranial nerve responsible for pupillary reactions. As the injury progresses, the ability of the pupils to respond to light may diminish or become uneven, reflecting increased intracranial pressure or brainstem involvement. Identifying pupillary changes is crucial in assessing the severity of a head injury. Late signs, such as those involving pupillary responses, often suggest a more serious level of brain injury, necessitating immediate medical intervention. In contrast, the other options listed are generally considered earlier signs of head injury or symptoms that may occur with various conditions rather than specifically indicating late-stage complications related to head trauma.

SAMPLE

3. What is a common physical change in school-age children as they begin to develop a sense of industry?

A. Increased anxiety

B. Improvements in cooperative play

C. Greater need for privacy

D. Decrease in friendships

As school-age children develop a sense of industry, they often show significant improvements in cooperative play. This developmental stage, which generally encompasses children aged 6 to 12 years, is marked by a growing ability to work with peers towards common goals and to engage in more complex group activities. Enhanced cooperative play not only helps children build friendships but also fosters important social skills such as sharing, teamwork, and problem-solving. During this time, children begin to take pride in their accomplishments and learned skills, which enhances their confidence and encourages them to collaborate with others. This sense of industry is crucial for their emotional and social development, allowing them to navigate the challenges of school and peer relationships more effectively. The option regarding increased anxiety generally signifies challenges that children may face rather than a typical change associated with developing a sense of industry. Similarly, the need for privacy and the dynamics of friendships also evolve during this period, but they do not represent the hallmark of social development that cooperative play does during the school-age years. Thus, improvements in cooperative play best exemplify the physical and social changes that accompany the development of a sense of industry in school-age children.

4. What is a key difference between viral and bacterial meningitis?

A. Viral often resolves with supportive care, while bacterial is more dangerous

B. Viral meningitis requires hospitalization, but bacterial does not

C. Both types present with similar symptoms and treatment

D. Viral meningitis only occurs in infants and children

The distinction between viral and bacterial meningitis is significant, particularly in terms of treatment and prognosis. Viral meningitis typically resolves with supportive care, as it is often less severe than bacterial meningitis. Patients diagnosed with viral meningitis usually recover fully without the need for aggressive medical intervention. In contrast, bacterial meningitis is a much more serious condition that can lead to severe complications, including neurological damage or death if not treated promptly with antibiotics. The emphasis on the danger of bacterial meningitis highlights its critical nature; this type of meningitis requires immediate medical attention and often hospitalization. While both types can present with similar symptoms, the treatment approaches and the urgency of care differ greatly between the two. Understanding the differences in management and outcomes is crucial for both healthcare professionals and caregivers when handling cases of meningitis. The other options present inaccuracies regarding the nature of these diseases. For example, while hospitalization may be typically necessary for bacterial meningitis, viral meningitis can often be managed on an outpatient basis. Additionally, it is not correct to claim that viral meningitis only occurs in infants and children, as it can affect individuals of all ages. Recognizing these key differences aids in the timely and effective treatment of meningitis.

SAMPLE

5. How long does the Babinski reflex typically last in infants?

A. Until 3 months

B. Usually until a year

C. Until 18 months

D. Until 2 years

The Babinski reflex is a normal neurological response observed in infants, characterized by the fanning of the toes when the sole of the foot is stroked. This reflex typically lasts until the child is about one year old. After this time, as the nervous system matures and brain development progresses, the reflex typically fades, and a more adult-like response occurs, where the toes curl inward instead. This timeframe aligns with significant developmental milestones during the first year of life, when the infant's motor skills and neurological pathways are rapidly maturing. The fact that the Babinski reflex can be assessed as a sign of neurological health in infants reinforces the importance of monitoring its presence and duration. Understanding the timing of this reflex helps caregivers and healthcare providers differentiate between normal development and potential neurological concerns.

6. What is the expected pulse rate of a child aged 3 months to 2 years?

A. 60 to 100/min

B. 70 to 150/min

C. 80 to 120/min

D. 90 to 130/min

The expected pulse rate for a child aged 3 months to 2 years typically ranges from 70 to 150 beats per minute. This age group is in a rapid growth and developmental phase, and their heart rates are generally higher than those of older children and adults. In infants and young toddlers, physiological factors such as their smaller body size, higher metabolic rate, and the demands of their developing organs contribute to this higher heart rate. It's important for healthcare providers to recognize this normal range when assessing pediatric patients, as deviations from it can indicate potential health issues. By understanding these normal physiological parameters, nurses and caregivers can more accurately identify anomalies that may require further evaluation or intervention.

7. What is a common complication associated with cleft palate that may require specialist intervention?

A. Speech difficulties

B. Dental problems

C. Ear infections and hearing loss

D. Gastroesophageal reflux disease

A common complication associated with cleft palate that may require specialist intervention is ear infections and hearing loss. Cleft palate often impacts the structure and function of the Eustachian tubes, which connect the middle ear to the throat. This can lead to a decreased ability to ventilate the middle ear properly, increasing the risk of fluid accumulation and infections. Without timely intervention, these ear infections can lead to persistent middle ear fluid and potentially result in hearing loss. Children with cleft palate often need regular monitoring, and they may require interventions such as tympanostomy tube placement to help prevent or manage these problems. Addressing ear infections and maintaining good hearing are crucial for a child's speech development and overall communication skills.

SAMPLE

8. What is often a sign of a ruptured appendix in children?

A. Improved pain

B. Severe diarrhea

C. High fever

D. Absence of vomiting

When considering the signs of a ruptured appendix in children, improved pain can be indicative of a change in the condition. Initially, a child with appendicitis typically experiences increasing abdominal pain, especially in the lower right quadrant. However, if the appendix ruptures, the pain may temporarily decrease due to the release of pressure within the abdomen. This change can lead to a false sense of relief for the patient, but it is a critical sign that the condition has worsened significantly and can lead to further complications such as peritonitis. In contrast, other symptoms related to a ruptured appendix generally point to more severe illness. Severe diarrhea may occur due to various gastrointestinal issues but is not a typical sign of appendicitis or its rupture. A high fever can be associated with many infections, including appendicitis, but does not specifically indicate rupture. Lastly, the absence of vomiting is not a clear sign of a ruptured appendix, as vomiting is common in appendicitis but may not be present in all cases. Understanding these nuances helps in recognizing the critical signs that might show a progression from appendicitis to a more severe condition like a rupture, which is vital for timely medical intervention.

9. What type of play is common for toddlers facing hospitalization?

A. Cooperative play

B. Parallel play

C. Constructive play

D. Associative play

Toddlers who are hospitalized often engage in parallel play. This is a developmental stage where children play alongside each other but do not necessarily interact with or influence one another's play. Given that toddlers are typically around 1 to 3 years of age, their play patterns are characterized by a growing awareness of others while still focusing on their own activities. In the hospital setting, this type of play can be particularly beneficial as it allows them to feel a sense of companionship and normalcy in an otherwise stressful environment without the pressure of social interaction. During this time, toddlers might play with similar toys or engage in similar activities without direct collaboration. This helps them express themselves and work through their experiences in the hospital context. Parallel play can also provide comfort and a sense of security, as they see other children nearby, which is crucial in making sense of their situation. In this context, the emphasis is on how toddlers use play to explore their environment and cope with the challenges of hospitalization. The development of more complex forms of play, such as associative or cooperative play, typically occurs as children grow older and becomes more proficient in social skills.

SAMPLE

10. Which of the following is true regarding the inheritance of cystic fibrosis?

A. It follows an autosomal dominant inheritance pattern

B. Both parents must carry the recessive trait

C. It is inherited from the father only

D. It only affects males

Cystic fibrosis is a genetic disorder that follows an autosomal recessive inheritance pattern. When both parents are carriers of the recessive gene for cystic fibrosis, they can pass this gene on to their children. For a child to be affected by cystic fibrosis, they must inherit two copies of the mutated gene, one from each parent. This means that if both parents are carriers, each child has a 25% chance of inheriting the disorder, a 50% chance of being a carrier like the parents, and a 25% chance of being unaffected. This understanding is key, as it highlights that both parents play a crucial role in the inheritance of this condition, which is why the assertion that both parents must carry the recessive trait is correct in the context of cystic fibrosis. The other options do not reflect the genetic mechanisms involved with cystic fibrosis: the disorder does not follow an autosomal dominant pattern, it can be inherited from either parent, and it can affect individuals of any gender.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://atinursingcareofchildren.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE