ATI PHARMACOLOGICAL AND PARENTERAL THERAPIES PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://
atipharmacologyparenteraltherapies.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which vital signs should be closely monitored when giving opioids?

A. Blood glucose levels and oxygen saturation

B. Respiratory rate, blood pressure, heart rate, and consciousness

C. Body temperature and electrolyte levels

D. Weight and height

2. What should a nurse assess before administering insulin?

A. Blood glucose levels and the patient's dietary intake

B. Patient's weight and height

C. Respiratory rate and blood pressure

D. Skin condition and temperature

3. When administering 1,000 mL of Lactated Ringer's IV over 6 hours with a drop factor of 10 gtt/mL, what is the required rate in gtt/min?

A. 20 gtt/min

B. 28 gtt/min

C. 25 gtt/min

D. 30 gtt/min

4. When monitoring a postoperative client, which behavior could suggest they may be experiencing pain?

A. Increased sleepiness

B. Crying

C. Clenching teeth

D. Relaxed posture

5. When administering liquid medications through an NG tube with intermittent suction, what action should the nurse take?

A. Attach the medication syringe immediately

B. Pinch the tube prior to attaching the medication syringe

C. Flush the tube with water first

D. Administer medications all at once

SAMPLE

6. How does pharmacogenomics impact patient care?

A. It limits the options of medications available

B. It allows for personalized medicine based on genetic makeup

C. It standardizes treatment across all patients

D. It decreases the frequency of medication use

7. What should a nurse observe for after administering chemotherapy?

A. Increased energy levels

B. Signs of side effects and adverse reactions

C. Changes in appetite

D. Request for additional pain medication

8. Which legislation governs the approval and regulation of medications in the United States?

A. The Health Insurance Portability and Accountability Act (HIPAA)

B. The Food, Drug, and Cosmetic Act (FDCA)

C. The Controlled Substances Act (CSA)

D. The Drug Enforcement Administration Policy (DEA)

9. What nonverbal behavior should the nurse identify as an indication that a postoperative client is experiencing pain?

A. Smiling

B. Restlessness

C. Speaking softly

D. Calm demeanor

10. What factors influence a medication's bioavailability?

A. Patient's age and weight

B. Route of administration, first-pass metabolism, and drug formulation

C. Duration of medication use

D. Patient's overall health condition

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. C
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Which vital signs should be closely monitored when giving opioids?

A. Blood glucose levels and oxygen saturation

B. Respiratory rate, blood pressure, heart rate, and consciousness

C. Body temperature and electrolyte levels

D. Weight and height

When administering opioids, it is essential to closely monitor respiratory rate, blood pressure, heart rate, and consciousness. Opioids are potent analgesics that can significantly depress the central nervous system, leading to respiratory depression, which is one of the most concerning side effects. Monitoring respiratory rate helps to ensure that the patient is not experiencing inadequate breathing, which can lead to serious complications. Blood pressure and heart rate are also monitored because opioids can cause hypotension and bradycardia in some patients; understanding these parameters allows for quicker interventions if the patient experiences adverse effects. Furthermore, monitoring consciousness is crucial to assess the patient's level of sedation and prevent overdose, which can lead to respiratory arrest or other severe outcomes. The other options include vital signs that are not as critical when specifically giving opioids. Blood glucose levels and oxygen saturation might be relevant in different contexts but are not the primary concerns associated with opioid administration. Body temperature and electrolyte levels, while important in overall patient assessment, do not directly pertain to the immediate effects of opioid therapy. Similarly, weight and height are unrelated to the monitoring needs during opioid administration. Therefore, focusing on respiratory rate, blood pressure, heart rate, and consciousness provides a comprehensive understanding of the patient's safety during opioid treatment

2. What should a nurse assess before administering insulin?

A. Blood glucose levels and the patient's dietary intake

B. Patient's weight and height

C. Respiratory rate and blood pressure

D. Skin condition and temperature

In the context of administering insulin, it is essential for the nurse to assess blood glucose levels and the patient's dietary intake. Monitoring blood glucose levels provides critical information on the patient's current glycemic status and helps determine the appropriate dosage of insulin needed, ensuring that it aligns with their metabolic needs. Additionally, understanding the patient's dietary intake is vital because insulin administration is closely related to food consumption. Insulin helps to manage the rise in blood glucose that occurs after eating, making it important to know what and when the patient has eaten to prevent both hyperglycemia (if insulin is given without sufficient food) and hypoglycemia (if food is not adequately timed with insulin administration). This careful assessment aids in personalizing diabetes management and promoting patient safety and effectiveness in treatment, making it the most pertinent choice in this context.

SAMPLE

3. When administering 1,000 mL of Lactated Ringer's IV over 6 hours with a drop factor of 10 gtt/mL, what is the required rate in gtt/min?

A. 20 gtt/min

B. 28 gtt/min

C. 25 gtt/min

D. 30 gtt/min

To determine the required rate in drops per minute when administering 1,000 mL of Lactated Ringer's solution over 6 hours with a drop factor of 10 gtt/mL, it's essential to follow these calculations. First, convert the total volume to mL: you have a total of 1,000 mL to be infused. Next, convert the infusion time from hours to minutes, since the final answer must be in gtt/min. There are 60 minutes in an hour, so for 6 hours, you multiply: 6 hours × 60 minutes/hour = 360 minutes. Now, to find the infusion rate in mL/minute, divide the total volume by the total time: 1,000 mL ÷ 360 minutes = approximately 2.78 mL/min. Next, convert this mL/min rate to gtt/min. Since the drop factor is given as 10 gtt/mL, you can convert mL/min to gtt/min by multiplying: 2.78 mL/min × 10 gtt/mL = 27.8 gtt/min. When rounding 27.8 gtt/min to the nearest whole number, it becomes approximately 28 g

4. When monitoring a postoperative client, which behavior could suggest they may be experiencing pain?

A. Increased sleepiness

B. Crying

C. Clenching teeth

D. Relaxed posture

Clenching teeth is a significant indicator that a patient may be experiencing pain, especially in a postoperative setting. This behavior often signifies tension and discomfort, reflecting the body's response to pain. Patients may subconsciously tighten their jaw muscles in response to pain, as clenching can be a way to cope with or guard against discomfort. In postoperative care, managing pain is critical to facilitate recovery, so recognizing non-verbal cues such as tooth clenching can be vital for timely intervention. In contrast, while increased sleepiness or crying might sometimes be associated with discomfort or distress, they are not as definitive as clenching teeth in indicating pain. Additionally, a relaxed posture indicates a state of comfort, which would usually be a sign that the patient is not currently experiencing significant pain. Thus, clenching teeth stands out as the most specific behavior linked to the experience of pain in this scenario.

5. When administering liquid medications through an NG tube with intermittent suction, what action should the nurse take?

A. Attach the medication syringe immediately

B. Pinch the tube prior to attaching the medication syringe

C. Flush the tube with water first

D. Administer medications all at once

When administering liquid medications through a nasogastric (NG) tube with intermittent suction, pinching the tube prior to attaching the medication syringe is crucial. This action helps to prevent air from entering the tube, which can disrupt the suction and cause complications such as improper medication delivery or aspiration. By pinching the tube, the nurse ensures a closed system while the medication is being administered. Additionally, this technique allows for a more controlled way to deliver the medication, ensuring that the full dose reaches the intended area without interference from the suction mechanism. After administering the medication, the nurse can then safely release the pinch to resume suction as needed. This method is essential for maintaining proper medication administration and patient safety while managing the NG tube.

SAMPLE

6. How does pharmacogenomics impact patient care?

A. It limits the options of medications available

B. It allows for personalized medicine based on genetic makeup

C. It standardizes treatment across all patients

D. It decreases the frequency of medication use

Pharmacogenomics significantly enhances patient care by tailoring medication choices to the individual's genetic profile. This discipline studies how a person's genes affect their response to drugs, allowing healthcare providers to predict which medications will be most effective or might cause adverse reactions. By considering an individual's genetic makeup, clinicians can select therapies that are more likely to succeed and reduce the trial-and-error approach often associated with prescribing. This personalized medicine approach helps to optimize treatment plans, improve medication adherence, minimize side effects, and ultimately lead to better health outcomes for patients. In contrast, other options do not capture the essence of pharmacogenomics. For instance, suggesting that it limits medication options overlooks the customization aspect, which actually enhances therapeutic possibilities. The idea that it standardizes treatment fails to acknowledge the personalized nature of care that pharmacogenomics promotes. Finally, claiming that it decreases medication frequency overlooks the potential for more effective treatments leading to improved management of conditions, which could stabilize a patient's health status and potentially decrease the number of medications needed over time.

7. What should a nurse observe for after administering chemotherapy?

A. Increased energy levels

B. Signs of side effects and adverse reactions

C. Changes in appetite

D. Request for additional pain medication

After administering chemotherapy, it is critical for a nurse to observe for signs of side effects and adverse reactions due to the nature of these drugs. Chemotherapy can have a wide range of effects on the body because it targets rapidly dividing cells, which includes not only cancer cells but also healthy cells in the bone marrow, digestive tract, and hair follicles. Common side effects may include nausea, vomiting, fatigue, hair loss, and changes in blood counts, which can lead to increased risk of infection or bleeding. Early recognition of these side effects allows for timely interventions, such as administering antiemetics for nausea or providing supportive care to manage fatigue. Monitoring for adverse reactions is crucial because some patients may experience severe or unexpected reactions that can compromise their safety or treatment effectiveness. By focusing on these potential complications, nurses can ensure that patients receive comprehensive care and that any necessary adjustments to their treatment regimen are made promptly. Other options such as increased energy levels, changes in appetite, or requests for additional pain medication might occur based on individual patient experiences and are important to assess on their own; however, they are not as immediate or critical as observing for side effects and adverse reactions following chemotherapy administration.

SAMPLE

8. Which legislation governs the approval and regulation of medications in the United States?

A. The Health Insurance Portability and Accountability Act (HIPAA)

B. The Food, Drug, and Cosmetic Act (FDCA)

C. The Controlled Substances Act (CSA)

D. The Drug Enforcement Administration Policy (DEA)

The Food, Drug, and Cosmetic Act (FDCA) establishes the foundation for the regulation of medications in the United States. This legislation grants the Food and Drug Administration (FDA) the authority to oversee the safety, efficacy, and labeling of drugs and medical devices. It requires that all new medications undergo rigorous testing and approval processes before they can be marketed to the public, ensuring that only safe and effective products enter the consumer marketplace. The FDCA was enacted in 1938, significantly expanding the FDA's responsibilities to include not only the regulation of food and drugs but also cosmetics and medical devices. The act has undergone several amendments, which have further strengthened regulatory processes, including provisions for expedited approvals for certain drugs, especially during public health emergencies. In contrast, other options like the Health Insurance Portability and Accountability Act primarily concern patient privacy and health information security rather than drug approval processes. The Controlled Substances Act focuses specifically on the regulation of controlled substances and their distribution, while the Drug Enforcement Administration Policy relates to the enforcement of laws surrounding controlled substances rather than the general approval and regulation of all medications.

9. What nonverbal behavior should the nurse identify as an indication that a postoperative client is experiencing pain?

A. Smiling

B. Restlessness

C. Speaking softly

D. Calm demeanor

Restlessness is a critical nonverbal behavior that indicates a client may be experiencing pain, especially in a postoperative setting. When patients are in discomfort or pain, they may exhibit behaviors such as fidgeting, shifting positions frequently, or being unable to stay still. This restlessness often reflects their inability to find a comfortable position and their struggle to cope with the discomfort they are feeling. In contrast, behaviors such as smiling or maintaining a calm demeanor typically suggest that a person is feeling well or is not in significant pain. Speaking softly does not convey any definitive indication of pain; patients can speak softly for numerous reasons unrelated to their pain levels. Recognizing restlessness as an indicator of pain allows nurses to intervene appropriately and provide necessary pain management strategies, improving the client's postoperative experience and comfort.

SAMPLE

10. What factors influence a medication's bioavailability?

A. Patient's age and weight

B. Route of administration, first-pass metabolism, and drug formulation

C. Duration of medication use

D. Patient's overall health condition

The correct factor that influences a medication's bioavailability is the route of administration, first-pass metabolism, and drug formulation. Bioavailability refers to the proportion of a drug that enters the circulation when introduced into the body and is available for action. The route of administration significantly affects how much of the drug is absorbed. For example, oral medications may undergo considerable first-pass metabolism, where the drug is metabolized by the liver before it reaches systemic circulation, thus reducing the bioavailability. In contrast, drugs administered intravenously bypass the digestive system and liver, leading to 100% bioavailability. Additionally, drug formulation plays a crucial role because the physical and chemical properties of the drug (like solubility and stability) and how it's formulated (tablet, liquid, etc.) can affect absorption rates and the extent to which the drug is available in the bloodstream. While patient factors, such as age, weight, health, and duration of medication use, can influence overall treatment effectiveness and how a patient responds to a drug, they do not directly determine the inherent pharmacokinetic properties of bioavailability. Thus, the specified factors—route of administration, first-pass metabolism, and drug formulation—are fundamental determinants of a medication's bioavailability.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://atipharmacologyparenteraltherapies.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE