AUSTRALIAN FIRST AID PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ausfirstaid.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. If a casualty with a head injury is unresponsive, what should you do next?

A. Leave them as they are
B. Put them in the recovery position
C. Give them water to drink
D. Start CPR immediately

2. When treating a casualty for asthma, what position should they be in?

A. Supine
B. Upside down
C. Sitting upright
D. Reclining

3. What is the immediate treatment for a burn?

A. Apply ice directly
B. Cover with a bandage
C. Cool the burn under running water
D. Use butter or oil

4. What is anaphylaxis?

A. A mild allergic reaction
B. A severe allergic reaction
C. A type of asthma attack
D. An infection of the airways

5. Which of the following is a common symptom of croup?

A. High fever
B. Barking cough
C. Blue lips
D. Severe distress

6. When performing CPR, what is the critical importance of the compression rate?

A. To ensure sufficient blood flow to vital organs
B. To minimize the risk of rib fractures
C. To prolong the duration of resuscitation
D. To allow time for rescue breaths

SAMPLE

7. How often should puffs without spacers be administered for asthma?

A. Two puffs every four minutes

B. Four puffs every four minutes

C. Six puffs every two minutes

D. One puff every minute

8. Which of the following statements about severe burns is correct?

A. They can be treated at home with band-aids

B. They require immediate medical assistance

C. They are usually not painful

D. They heal quickly without medical coverage

9. What action should be taken if a patient is still experiencing anaphylaxis symptoms after using an EpiPen?

A. Use a second EpiPen after five minutes

B. Wait for further instructions from a doctor

C. Start CPR immediately

D. Encourage them to move around

10. What action should be taken if a casualty is unconscious from hypothermia?

A. Check breathing every 30-45 seconds

B. Give them water immediately

C. Wait for them to wake up

D. Move them to a colder location

SAMPLE

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. B
6. A
7. B
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. If a casualty with a head injury is unresponsive, what should you do next?

A. Leave them as they are

B. Put them in the recovery position

C. Give them water to drink

D. Start CPR immediately

When a casualty with a head injury is unresponsive, the most appropriate action is to place them in the recovery position. This position helps maintain an open airway, which is crucial since unresponsive individuals may have a compromised ability to breathe properly. By placing the casualty on their side, any fluid or vomit that may obstruct the airway can drain away, reducing the risk of choking or aspiration. Moreover, the recovery position allows for better access to the airway and ensures that the casualty's breathing can be monitored closely. It's essential to avoid any unnecessary movement of the head or neck, as a head injury may also involve spinal injury, and improper handling can exacerbate these injuries. Other options listed are not suitable in this context. Leaving the casualty as they are fails to address potential airway issues. Providing water could lead to choking and is inappropriate due to their unresponsive state. Starting CPR is not indicated unless there are signs of cardiac arrest or the casualty is not breathing. Therefore, placing the casualty in the recovery position is the most effective and responsible action to take in response to an unresponsive individual with a head injury.

2. When treating a casualty for asthma, what position should they be in?

A. Supine

B. Upside down

C. Sitting upright

D. Reclining

When treating a casualty for asthma, the most effective position for them to be in is sitting upright. This position helps to open the airways, allowing for easier breathing. When a person is upright, gravity assists in lung expansion and can reduce the pressure on the diaphragm, which is crucial during an asthma attack when the airway becomes narrowed due to inflammation and bronchoconstriction. In this position, the casualty can take deeper breaths and utilize their accessory muscles of respiration more effectively, which is vital during an asthma episode. It also helps them remain calm and focused, which can further aid in their ability to breathe. Other positions, such as supine (lying flat on the back) or reclining, can restrict lung capacity and make breathing more difficult. Being upside down would be counterproductive and could lead to further distress or complications. Hence, sitting upright is the most appropriate choice for facilitating effective breathing in an asthmatic emergency.

SAMPLE

3. What is the immediate treatment for a burn?

A. Apply ice directly

B. Cover with a bandage

C. Cool the burn under running water

D. Use butter or oil

The immediate treatment for a burn is to cool the burn under running water. This is important as cooling the burn effectively helps to reduce the temperature of the skin, which minimizes tissue damage and alleviates pain. Running cool water over the burn for at least 20 minutes helps to halt the burning process and prevent further injury. Additionally, cooling the burn can reduce inflammation and swelling, providing essential pain relief. It is recommended to use cooler, not ice-cold, water as extreme cold can further damage the skin tissue. Using a bandage is not the immediate first step because the priority is to cool the burn first to address the injury effectively. Applying ice directly can also cause frostbite and additional harm to the tissue, and using butter or oil is outdated practice; it can trap heat in the skin and promote infection rather than assist in recovery. Therefore, cooling the burn under running water is the most effective and safest method for immediate treatment.

4. What is anaphylaxis?

A. A mild allergic reaction

B. A severe allergic reaction

C. A type of asthma attack

D. An infection of the airways

Anaphylaxis is recognized as a severe allergic reaction that can occur rapidly and can be life-threatening if not treated promptly. This condition is characterized by symptoms such as difficulty breathing, swelling of the throat and tongue, hives, rapid or weak pulse, and a drop in blood pressure. In the context of your options, it's important to understand that anaphylaxis is not simply a mild allergic reaction; mild reactions typically involve local symptoms such as itching or mild rash rather than the systemic responses seen in anaphylaxis. It is also distinct from a type of asthma attack, which may involve breathing problems but is not necessarily triggered by allergy-related mechanisms in the same way anaphylaxis is. Finally, it is unrelated to infections of the airways, as infections would involve pathogens and distinct symptoms associated with illness rather than the immune system's overreaction to an allergen. Therefore, anaphylaxis represents a critical medical emergency that requires immediate intervention, often with the administration of epinephrine and subsequent medical care.

5. Which of the following is a common symptom of croup?

A. High fever

B. Barking cough

C. Blue lips

D. Severe distress

A barking cough is a classic symptom of croup, which is often associated with viral infections in young children. The cough gets its characteristic sound due to the inflammation and swelling of the airways, specifically the larynx and trachea. This inflammation narrows the airway, producing a distinctive cough that resembles the sound of a seal barking. Other symptoms of croup may include a hoarse voice and difficulty breathing, especially when the child is agitated or lying down. These symptoms can worsen at night, leading to increased concern for parents and caregivers. While high fever, blue lips, and severe distress can be associated with various medical conditions, they are not typical presentations specifically highlighting croup itself. The barking cough serves as a hallmark sign, making it a critical factor in the diagnosis and management of this condition.

SAMPLE

6. When performing CPR, what is the critical importance of the compression rate?

A. To ensure sufficient blood flow to vital organs

B. To minimize the risk of rib fractures

C. To prolong the duration of resuscitation

D. To allow time for rescue breaths

The critical importance of the compression rate during CPR lies in its role in ensuring sufficient blood flow to vital organs, particularly the heart and brain. When someone experiences a cardiac arrest, their heart stops beating effectively, which means that blood is not being pumped to sustain vital organs. The purpose of performing chest compressions is to manually circulate blood and maintain perfusion to these organs until professional medical help arrives or until an automated external defibrillator (AED) can be used. A compression rate of 100 to 120 compressions per minute is recommended because it is effective at maintaining blood flow. This rate helps to provide adequate oxygenation and nutrient delivery to organs while also allowing for sufficient time for the chest to recoil between compressions, which is crucial for maximizing blood return to the heart. Other options relate to concerns that are less crucial in the overall response to cardiac arrest. Minimizing rib fractures, prolonging resuscitation duration, and allowing time for rescue breaths, while important in their own contexts, do not directly address the primary objective of restoring blood circulation effectively during CPR. The urgency of maintaining a proper compression rate prioritizes brain and heart function, making it the most critical component of the CPR process.

7. How often should puffs without spacers be administered for asthma?

A. Two puffs every four minutes

B. Four puffs every four minutes

C. Six puffs every two minutes

D. One puff every minute

For managing asthma symptoms, particularly during an attack, it is generally recommended to administer four puffs of a reliever medication (such as a short-acting beta-agonist) every four minutes if symptoms do not improve. This approach is designed to provide a rapid response to acute asthma symptoms, allowing sufficient medication to be delivered into the lungs to help open the airways. The frequency and number of puffs work to maintain adequate bronchodilation, ensuring that the individual receives enough medication for effective relief. Using a spacer is often encouraged as it can enhance medication delivery and effectiveness; however, the question specifically addresses the situation without a spacer. It's important to follow established guidelines and protocols for asthma management to ensure patient safety and effective treatment. Other options that suggest fewer puffs or an increased frequency may not provide adequate relief and could place the person at risk of inadequate treatment of acute asthma symptoms. Overall, administering four puffs every four minutes is a widely accepted standard in emergency and first aid practice for asthma management.

SAMPLE

8. Which of the following statements about severe burns is correct?

A. They can be treated at home with band-aids

B. They require immediate medical assistance

C. They are usually not painful

D. They heal quickly without medical coverage

Severe burns are indeed a medical emergency and require immediate professional medical assistance. This is because severe burns can affect deeper layers of skin and potentially damage underlying tissues, which may lead to complications such as infection, fluid loss, and shock. Timely intervention is critical to ensure appropriate treatment, minimize complications, and promote healing. In contrast, treating severe burns at home with simple remedies, like band-aids, is inappropriate due to the level of damage and risk involved. These types of burns typically also involve significant pain due to nerve damage and require professional care for effective pain management. Furthermore, severe burns do not heal quickly without proper medical intervention; they often need extensive treatment, including specialized wound care and possibly surgical procedures, to facilitate recovery.

9. What action should be taken if a patient is still experiencing anaphylaxis symptoms after using an EpiPen?

A. Use a second EpiPen after five minutes

B. Wait for further instructions from a doctor

C. Start CPR immediately

D. Encourage them to move around

Using a second EpiPen after five minutes is the appropriate action if a patient continues to experience symptoms of anaphylaxis after the first injection. Anaphylaxis is a severe, life-threatening allergic reaction that can progress rapidly. The initial dose of adrenaline (epinephrine) from the EpiPen can sometimes be insufficient to counteract the reaction fully, especially in cases of severe symptoms or when the patient has additional risk factors, such as a high body mass or a significant allergenic exposure. Administering a second dose of adrenaline can help to stabilize the patient by further constricting blood vessels, increasing heart rate, and opening the air passages in the lungs, which counteracts the symptoms of anaphylaxis. The five-minute interval is advised because it allows time for the adrenaline to take effect, and further doses may be necessary in cases of severe, ongoing symptoms. In contrast, waiting for further instructions from a doctor may leave the patient in a critical state without appropriate intervention. Starting CPR immediately is not warranted unless the patient is unresponsive and not breathing, as the priority should be dealing with the anaphylaxis symptoms first. Encouraging movement can be dangerous as it can increase the heart rate and worsen the patient's condition, therefore it is not

SAMPLE

10. What action should be taken if a casualty is unconscious from hypothermia?

A. Check breathing every 30-45 seconds

B. Give them water immediately

C. Wait for them to wake up

D. Move them to a colder location

In a situation where a casualty is unconscious from hypothermia, checking their breathing every 30-45 seconds is crucial because it allows you to monitor their airway and respiratory function. Hypothermia can lead to severe complications, including respiratory failure, so it's important to ensure the individual is breathing adequately. If their breathing stops, timely intervention, such as performing CPR, may be necessary to prevent further harm or death. Constant monitoring provides critical information that can guide any immediate first aid actions while waiting for professional help to arrive or while preparing to take them to a warmer environment. Maintaining awareness of their breathing status ensures that you are prepared to act quickly if their condition changes. Providing water is not appropriate in this scenario since the casualty is unconscious, which raises the risk of choking. Moving them to a colder location would worsen their condition, and simply waiting for them to wake up neglects the need for active monitoring and intervention.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ausfirstaid.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE