AWHONN PERINATAL ORIENTATION AND EDUCATION PROGRAM (POEP) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://awhonnpoep.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which is true regarding vital signs monitoring post-operatively?

A. Only blood pressure needs to be monitored

B. Vital signs are less critical if the patient is stable

C. Continuous monitoring is needed for the first hour

D. Vital signs should be taken hourly

2. Cardiac output in the immediate postpartum period is generally:

A. Decreased

B. Increased

C. Only of concern with a history of cardiac disease

D. Not affected during the recovery period

3. What type of cardiac disease is associated with advanced maternal age, hypertension, and diabetes in pregnancy?

A. Acquired

B. Congenital

C. Hemorrhagic

D. Ischemic

4. What is a potential outcome for an obese primiparous woman with unmanaged glucose levels during pregnancy?

A. Postpartum hemorrhage

B. Delayed fetal development

C. Gestational diabetes

D. Increased incidence of cesarean delivery

5. What is the most accurate sign to monitor for early signs of complications in a postpartum cesarean patient?

A. Urine output

B. Fundal height

C. Reflex responses

D. Respiratory rate

SAMPLE

6. What is the initial drug of choice for excessive bleeding in the postpartum period?

A. Methergine, IM

B. Oxytocin, IV

C. Prostaglandin, PR

D. Terbutaline, SC

7. What is the initial nonpharmacologic intervention for postpartum hemorrhage caused by involution failure?

A. External or bimanual uterine massage

B. Insertion of an indwelling Foley catheter

C. Pulse oximetry and cardiac monitoring

D. Reverse Trendelenburg position

8. How can a nurse assess recovery from neuraxial anesthesia?

A. By checking sensory awareness on one side

B. By observing for movement in both legs

C. By evaluating reaction to blunt needle prick

D. By measuring blood pressure

9. Women undergoing induction of labor are at an increased risk for which of the following?

A. Postpartum hemorrhage

B. Umbilical cord prolapse

C. Longer second stage of labor

D. Fetus in occiput posterior position

10. In addition to hepatitis B, which other type of hepatitis virus may also be prevented with the hepatitis B vaccine?

A. Hepatitis A

B. Hepatitis C

C. Hepatitis D

D. Hepatitis G

SAMPLE

Answers

SAMPLE

1. C
2. B
3. D
4. D
5. B
6. B
7. A
8. A
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. Which is true regarding vital signs monitoring post-operatively?

A. Only blood pressure needs to be monitored

B. Vital signs are less critical if the patient is stable

C. Continuous monitoring is needed for the first hour

D. Vital signs should be taken hourly

Post-operative vital signs monitoring is crucial for ensuring patient safety and identifying early signs of complications. Continuous monitoring for the first hour is essential because this period is when patients are most vulnerable and may experience changes in their condition due to anesthesia effects or surgical complications. During this time, healthcare providers can promptly detect alterations in heart rate, respiratory rate, blood pressure, and temperature, allowing for immediate intervention if needed. Monitoring vital signs continuously helps in assessing the patient's response to surgery and anesthesia, and can provide vital data that informs the clinical team about the patient's recovery progress. Continuous or more frequent monitoring during the initial recovery phase is a standard practice to ensure that any potential issues are addressed quickly, reducing the risk of serious complications. In contrast, while monitoring during stable periods is important, the focus immediately post-operatively remains on ensuring that the patient is stable through close observation. The approach of monitoring vital signs only on an hourly basis or focusing solely on blood pressure would be inadequate during this critical recovery window. Therefore, initial continuous monitoring is a best practice in the post-operative setting.

2. Cardiac output in the immediate postpartum period is generally:

A. Decreased

B. Increased

C. Only of concern with a history of cardiac disease

D. Not affected during the recovery period

In the immediate postpartum period, cardiac output increases due to several physiological changes that occur in the body following childbirth. After the delivery of the placenta, there is a significant decrease in intra-abdominal pressure, which allows the blood that was previously sequestered in the uterine vascular bed to return to the systemic circulation. This increase in circulating blood volume contributes to a rise in cardiac output. Additionally, hormonal changes and the restoration of blood volume can also enhance cardiac performance. This increase is a normal physiological response as the body adapts to the sudden changes and transitions from pregnancy to the postpartum state. It is important to recognize that this increase in cardiac output is a part of the body's effort to stabilize hemodynamics and ensure adequate perfusion to vital organs after the stress of labor and delivery. Understanding this physiological adaptation is crucial, especially when monitoring postpartum patients for signs of complications such as hemorrhage or cardiovascular instability. The normal elevation of cardiac output in this period helps provide insights into maternal health and recovery, necessitating close observation and assessment.

SAMPLE

3. What type of cardiac disease is associated with advanced maternal age, hypertension, and diabetes in pregnancy?

A. Acquired

B. Congenital

C. Hemorrhagic

D. Ischemic

The correct answer is ischemic heart disease, which pertains to conditions that arise due to a lack of blood flow to the heart muscle, typically caused by blockages in the coronary arteries. In the context of pregnancy, advanced maternal age, hypertension, and diabetes significantly increase the risk of cardiovascular complications, including ischemic heart disease. Advanced maternal age is linked to multiple factors that affect heart health, including pre-existing conditions like hypertension and diabetes. These conditions can lead to changes in vascular health and increased myocardial oxygen demand, setting the stage for ischemic events. Hypertension can cause increased workload on the heart, while diabetes can lead to vascular damage over time, making individuals more susceptible to ischemic episodes. During pregnancy, the physical changes and demands placed on the mother's body can exacerbate these pre-existing conditions, thus elevating the risk of ischemic heart disease. This highlights the importance of careful monitoring and management of cardiovascular health in pregnant individuals, particularly those presenting with these risk factors.

4. What is a potential outcome for an obese primiparous woman with unmanaged glucose levels during pregnancy?

A. Postpartum hemorrhage

B. Delayed fetal development

C. Gestational diabetes

D. Increased incidence of cesarean delivery

The potential outcome of increased incidence of cesarean delivery in an obese primiparous woman with unmanaged glucose levels during pregnancy stems from the complications often associated with obesity and gestational diabetes. Obesity can lead to a variety of pregnancy-related issues, including larger fetal size (macrosomia), which increases the likelihood of a cesarean delivery. This is because larger babies can complicate the labor process, making vaginal delivery more challenging or even unsafe, resulting in the need for surgical intervention. In addition to the mechanical difficulties of delivering a larger fetus, obesity is linked to various other factors, such as increased risk of hypertension and preeclampsia, which may further contribute to the decision to perform a cesarean section. Given these factors, unmanaged glucose levels can exacerbate these risks, leading healthcare providers to opt for a cesarean delivery for the safety of both mother and child. Understanding the direct relationship between obesity, unmanaged glucose levels, and the increased risk of cesarean delivery highlights the importance of proper management of maternal health during pregnancy.

SAMPLE

5. What is the most accurate sign to monitor for early signs of complications in a postpartum cesarean patient?

A. Urine output

B. Fundal height

C. Reflex responses

D. Respiratory rate

Monitoring for early signs of complications in a postpartum cesarean patient is crucial for timely intervention and ensuring patient safety. One of the most reliable indicators in this scenario is fundal height. The fundus is the top portion of the uterus, and its position provides important information about uterine involution following delivery. After a cesarean section, it is essential to assess fundal height regularly. A fundus that is higher than expected, particularly if rising in the abdomen instead of descending, can indicate conditions such as uterine atony, retention of placental fragments, or hemorrhage. Early detection of these issues can significantly impact patient outcomes and facilitate prompt clinical intervention. Other options such as urine output, reflex responses, and respiratory rate provide valuable information but might not be as direct or sensitive for assessing uterine health or complications specifically associated with the postpartum period following a cesarean delivery. While urine output can indicate kidney function and fluid balance, reflex responses are more generalized neurological assessments, and respiratory rate, although important, is more related to respiratory function rather than directly monitoring uterine complications. Hence, the assessment of fundal height stands out as the most critical parameter in this context.

6. What is the initial drug of choice for excessive bleeding in the postpartum period?

A. Methergine, IM

B. Oxytocin, IV

C. Prostaglandin, PR

D. Terbutaline, SC

Oxytocin administered intravenously is recognized as the initial drug of choice for managing excessive bleeding, or postpartum hemorrhage (PPH), following childbirth. The primary mechanism of action for oxytocin in this context is its ability to stimulate uterine contractions, which helps to constrict uterine blood vessels and reduce bleeding. This contraction is critical in promoting uterine tone and expelling any retained placental tissue, which can contribute to continued bleeding. Oxytocin is preferred in the immediate postpartum period as it is well-studied, quickly effective, and has a favorable safety profile when used in a clinical setting. Medical providers often utilize it as the first-line treatment because rapid intervention is essential to prevent complications associated with significant blood loss. While other medications may play roles in the management of postpartum hemorrhage, particularly in cases that do not respond to oxytocin or in specific situations (such as Methergine, which is typically used for uterine atony but can cause hypertension), oxytocin remains the first consideration in emergency protocols for this urgent situation.

SAMPLE

7. What is the initial nonpharmacologic intervention for postpartum hemorrhage caused by involution failure?

A. External or bimanual uterine massage

B. Insertion of an indwelling Foley catheter

C. Pulse oximetry and cardiac monitoring

D. Reverse Trendelenburg position

The initial nonpharmacologic intervention for postpartum hemorrhage due to involution failure is external or bimanual uterine massage. This technique is critical because it helps stimulate uterine contractions, which can assist in reducing bleeding by promoting uterine tone. Involution failure often results in a uterus that is not contracting effectively, leading to excessive blood loss. By applying pressure through massage, the practitioner can encourage the uterus to contract and thus decrease the risk of continued hemorrhage. Other interventions may play a role in post-hemorrhage care, but they are not the first-line nonpharmacologic approach. The insertion of an indwelling Foley catheter is more of a supportive measure to monitor urine output or relieve bladder distension but does not address the underlying issue of uterine atony directly. Similarly, pulse oximetry and cardiac monitoring are important for assessing the mother's condition but do not contribute to resolving uterine atony specifically. The Reverse Trendelenburg position may be utilized to improve venous return in cases of hypotension or hypovolemia, but it does not directly stimulate the uterus to contract, making uterine massage the preferred initial intervention in managing this situation.

8. How can a nurse assess recovery from neuraxial anesthesia?

A. By checking sensory awareness on one side

B. By observing for movement in both legs

C. By evaluating reaction to blunt needle prick

D. By measuring blood pressure

Evaluating recovery from neuraxial anesthesia is crucial in monitoring a patient's safety and comfort following a procedure. The correct approach involves checking for sensory awareness on one side, as it directly assesses the effectiveness of the anesthesia and the patient's sensory recovery. After administering neuraxial anesthesia (which includes epidurals or spinal blocks), the nurse needs to assess which areas have regained sensation. This is often done by comparing the sensory awareness on the anesthetized side to the non-anesthetized side. Sensory assessment provides insight into the distribution and duration of the block, which is fundamental in managing postoperative care effectively. While observing movement in both legs can indicate recovery, it is not always the most immediate measurement of sensory recovery after neuraxial anesthesia, as some patients may regain motor function before sensation. Evaluating reaction to a blunt needle prick can also be useful, but it typically requires more targeted sharp/dull sensory tests, which may not be as straightforward as checking for overall sensory awareness. Measuring blood pressure, while important, is more related to the potential hemodynamic effects of the anesthesia rather than a direct assessment of sensory recovery. Thus, checking sensory awareness provides the most relevant and immediate information regarding recovery from neuraxial anesthesia.

SAMPLE

9. Women undergoing induction of labor are at an increased risk for which of the following?

A. Postpartum hemorrhage

B. Umbilical cord prolapse

C. Longer second stage of labor

D. Fetus in occiput posterior position

The correct answer is postpartum hemorrhage, as women undergoing induction of labor may face several physiological changes and interventions that increase this risk. Induction can lead to uterine over-distension or hyperstimulation, which might cause uterine atony—one of the leading causes of postpartum hemorrhage. Additionally, interventions such as the use of oxytocin can lead to a more rapid or intense labor, increasing the likelihood of uterine fatigue after delivery. Longer second stage of labor can occur in some instances, but it is not universally associated with induction; various factors influence this stage, including the mother's pelvic anatomy and the baby's position. Although umbilical cord prolapse can happen during labor, it is not uniquely increased by labor induction. Similarly, while a fetus in the occiput posterior position can present challenges during delivery, it is not specifically linked to induction of labor. The focus on postpartum hemorrhage underscores the importance of monitoring and preparedness for potential complications following labor induction.

10. In addition to hepatitis B, which other type of hepatitis virus may also be prevented with the hepatitis B vaccine?

A. Hepatitis A

B. Hepatitis C

C. Hepatitis D

D. Hepatitis G

The hepatitis B vaccine is designed to provide immunity specifically against the hepatitis B virus. However, part of the reason why the correct response identifies hepatitis D is due to the relationship between the two viruses. Hepatitis D virus (HDV) is an incomplete virus that can only replicate in the presence of hepatitis B virus (HBV). Thus, by being vaccinated against hepatitis B, an individual is also protected from hepatitis D infection, since HDV cannot establish an infection without co-infection with HBV. While hepatitis A is a separate virus that can be prevented with its own vaccine, the hepatitis C virus is also a distinct entity, and a vaccine does not currently exist for it. Hepatitis G, on the other hand, is less clinically significant and has no associated vaccine. Therefore, the protection provided by the hepatitis B vaccine extends specifically to hepatitis D, making it the correct answer to the question.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://awhonnpoep.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE