BASIC LIFE SUPPORT (BLS) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://basiclifesupport.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is the first step in managing a patient with a suspected cardiac arrest?**

   A. Begin CPR immediately

   B. Call emergency services

   C. Check for responsiveness

   D. Start using an AED

2. **In the case of a cardiac arrest due to an opioid overdose, what should be prioritized?**

   A. Administer naloxone first

   B. Notify family members

   C. Ensure high-quality CPR over naloxone administration

   D. Transport the patient to the hospital quickly

3. **What is the depth of chest compressions required for an adult patient?**

   A. At least 1 inch

   B. At least 2 inches

   C. At least 3 inches

   D. At least 4 inches

4. **Which situation is appropriate for using an AED on a patient with an implanted device?**

   A. AED pads can be placed directly over the device

   B. The device should be removed before use

   C. AED pads must not be placed directly over the device

   D. Only use AED if the device is malfunctioning

5. **Which condition is the primary reason for prioritizing CPR in opioid overdose cases?**

   A. Loss of consciousness

   B. Respiratory failure

   C. Circulatory collapse

   D. Complete airway obstruction

SAMPLE

6. During CPR for a child or infant with an advanced airway, how often should one provider deliver ventilations?

A. Every 30 seconds

B. Every 2 to 3 seconds

C. Continuous ventilations

D. Once every minute

7. If you suspect an opioid overdose in a person who is unresponsive but has a strong pulse, what action should you take?

A. Start CPR immediately

B. Provide rescue breathing

C. Use an AED

D. Administer naloxone

8. What should you attempt next if an adult patient becomes unresponsive while clearing their obstructed airway and no object is visible in their mouth?

A. Give 2 ventilations

B. Perform a finger sweep

C. Start chest compressions again

D. Check for responsiveness

9. What is the recommended depth for performing chest compressions on an adult?

A. 1 inch

B. 1.5 inches

C. 2 inches

D. 2.5 inches

10. If a child with an obstructed airway becomes unresponsive, what should you do first?

A. Immediately start providing rescue breaths

B. Call for emergency medical assistance

C. Gently lower the child to the ground and begin CPR

D. Attempt back blows and abdominal thrusts

SAMPLE

Answers

SAMPLE

1. C
2. C
3. B
4. C
5. B
6. B
7. B
8. A
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. What is the first step in managing a patient with a suspected cardiac arrest?

A. Begin CPR immediately

B. Call emergency services

C. Check for responsiveness

D. Start using an AED

The first step in managing a patient with a suspected cardiac arrest is to check for responsiveness. This is essential because it helps you assess the patient's condition immediately. By checking for responsiveness, you determine whether the individual is conscious or not, which further guides your next actions. If the person is unresponsive, it indicates that they may require emergency assistance and immediate intervention. Following this initial assessment, if the person is indeed unresponsive, the next appropriate steps would typically include calling emergency services for help and then starting CPR if the patient shows no signs of circulation, such as breathing or movement. This systematic approach ensures that the provider prioritizes assessing the patient's state before moving to procedures like CPR or defibrillation. Overall, checking for responsiveness is a crucial first step in effectively managing cardiac arrests or other emergencies.

2. In the case of a cardiac arrest due to an opioid overdose, what should be prioritized?

A. Administer naloxone first

B. Notify family members

C. Ensure high-quality CPR over naloxone administration

D. Transport the patient to the hospital quickly

In the case of a cardiac arrest due to an opioid overdose, the priority is to ensure high-quality CPR over naloxone administration. The reason for this is that CPR is crucial for maintaining blood flow and oxygen delivery to vital organs during cardiac arrest. When a patient suffers a cardiac arrest, time is of the essence; the brain and other organs become increasingly susceptible to damage the longer they go without oxygen. While administering naloxone is an effective treatment for reversing opioid overdoses, it should not delay the initiation of CPR. In fact, if a patient is unresponsive or not breathing normally due to an opioid overdose, immediate chest compressions can help buy time until naloxone can be provided. High-quality CPR involves proper technique, adequate depth and rate of compressions, and minimizing interruptions, which are essential for improving the chances of survival. Other options like notifying family members or transporting the patient to the hospital are important but secondary actions that should follow the initiation of high-quality CPR.

3. What is the depth of chest compressions required for an adult patient?

A. At least 1 inch

B. At least 2 inches

C. At least 3 inches

D. At least 4 inches

The required depth of chest compressions for an adult patient during cardiopulmonary resuscitation (CPR) is at least 2 inches. This depth is crucial because compressions that are too shallow may not effectively circulate blood and provide adequate perfusion to vital organs, such as the heart and brain. When performing compressions, aiming for a depth of 2 to 2.4 inches helps ensure that the heart is compressed sufficiently to generate blood flow. Research and guidelines from organizations such as the American Heart Association emphasize the importance of this depth to optimize the chances of survival in cases of cardiac arrest. By adhering to this standard, responders can greatly improve the effectiveness of their CPR efforts and increase the likelihood of a positive outcome for the patient.

SAMPLE

4. Which situation is appropriate for using an AED on a patient with an implanted device?

A. AED pads can be placed directly over the device

B. The device should be removed before use

C. AED pads must not be placed directly over the device

D. Only use AED if the device is malfunctioning

Using an AED on a patient with an implanted device such as a pacemaker or implantable cardioverter-defibrillator (ICD) requires careful consideration to ensure both the effectiveness of the defibrillation and the safety of the patient. It is essential that the AED pads are not placed directly over the implanted device because this can impede the delivery of the shock to the heart and may also cause damage to the device itself. When using the AED, the pads should be placed at least a few inches away from the implanted device to ensure the electrical discharge can effectively reach the heart muscle and is not obstructed. Placing the pads correctly helps maximize the chance of a successful shock while minimizing the risk of causing harm to the patient or the implanted device. This approach ensures compliance with safety protocols in emergency care when dealing with patients with implanted devices.

5. Which condition is the primary reason for prioritizing CPR in opioid overdose cases?

A. Loss of consciousness

B. Respiratory failure

C. Circulatory collapse

D. Complete airway obstruction

In cases of opioid overdose, prioritizing CPR is primarily due to respiratory failure. Opioids depress the central nervous system, leading to decreased respiratory drive, which can result in inadequate or absent breathing. When breathing stops or is significantly compromised, it deprives the brain and other vital organs of oxygen, leading to potential brain death and organ failure in a matter of minutes. Performing CPR helps restore circulation and provides some degree of ventilation, which is crucial in delivering oxygen to the body until advanced medical help can be rendered. While loss of consciousness may occur as a result of the respiratory failure, it is the lack of adequate breathing that directly threatens life by causing oxygen deprivation. Circulatory collapse can occur as a later consequence of prolonged respiratory failure and inadequate oxygen delivery. Complete airway obstruction is a critical condition, but in the context of opioid overdose specifically, the primary issue is the respiratory failure caused by the drug's effects. This makes addressing the breathing irregularities an immediate focus for those providing aid.

SAMPLE

6. During CPR for a child or infant with an advanced airway, how often should one provider deliver ventilations?

A. Every 30 seconds

B. Every 2 to 3 seconds

C. Continuous ventilations

D. Once every minute

In the context of providing CPR for a child or infant with an advanced airway, the correct practice involves delivering ventilations every 2 to 3 seconds. This recommendation is based on the need to ensure that effective breaths are being provided to the child or infant without compromising the quality of chest compressions. When an advanced airway is utilized, continuous chest compressions can be administered, and the timing of ventilations is adjusted to this protocol. Delivering a breath every 2 to 3 seconds aligns with the goal of maintaining a proper ratio of ventilation to circulation, ensuring that the child receives adequate oxygen while also prioritizing effective chest compressions. Other timing options, such as every 30 seconds or once every minute, would not support the physiological needs of the child during CPR. Additionally, continuous ventilations would not provide effective support in a scenario where chest compressions are critical for maintaining circulation. Therefore, establishing a rhythm of ventilations every 2 to 3 seconds is essential for optimal resuscitation outcomes in this age group.

7. If you suspect an opioid overdose in a person who is unresponsive but has a strong pulse, what action should you take?

A. Start CPR immediately

B. Provide rescue breathing

C. Use an AED

D. Administer naloxone

In the context of an unresponsive person with a strong pulse who is suspected of an opioid overdose, the appropriate action to take is to provide rescue breathing. When a person is unresponsive but still has a pulse, it indicates that the heart is functioning, but the individual may not be breathing adequately or at all due to the effects of opioids. Rescue breathing helps to ensure that oxygen continues to flow into the lungs and gets delivered to the brain and other vital organs. This action is critically important to prevent brain damage and maintain tissue viability while waiting for further medical help or intervention. Administering naloxone is typically a highly effective action in cases of suspected opioid overdose, but it should ideally be done in conjunction with providing rescue breathing, especially if the individual is breathing inadequately. In this scenario, it's also essential to call for emergency medical help to ensure comprehensive care is provided as soon as possible. While CPR and the use of an AED are important in cases of cardiac arrest, they are not the immediate appropriate actions when there is a strong pulse. Rescue breaths address the primary issue of insufficient respiration due to the overdose.

SAMPLE

8. What should you attempt next if an adult patient becomes unresponsive while clearing their obstructed airway and no object is visible in their mouth?

A. Give 2 ventilations

B. Perform a finger sweep

C. Start chest compressions again

D. Check for responsiveness

If an adult patient becomes unresponsive while you are addressing an obstructed airway and no object is visible in their mouth, the appropriate next step is to provide 2 ventilations. This action is crucial because it is essential to attempt to deliver breath to the lungs; a lack of responsiveness indicates that breathing is compromised. The goal in this situation is to re-establish oxygenation in the body. Following the delivery of ventilations, if the airway remains obstructed and the patient does not respond to rescue breaths, you would then need to reassess the situation, which might include resuming chest compressions and repositioning the airway as needed. However, when the primary issue is unresponsiveness with a clear inability to breathe, delivering 2 ventilations is a logical and necessary step as it directly addresses the issue of insufficient air supply. Therefore, in the context of Basic Life Support, the focus on securing the airway and trying to breathe for the patient is the most immediate and appropriate response.

9. What is the recommended depth for performing chest compressions on an adult?

A. 1 inch

B. 1.5 inches

C. 2 inches

D. 2.5 inches

The recommended depth for performing chest compressions on an adult is 2 inches. This depth is essential for effectively generating adequate blood flow and ensuring that chest compressions are both effective and capable of promoting circulation to vital organs during cardiopulmonary resuscitation (CPR). Compression depth is critical because if compressions are too shallow, it may not adequately compress the heart to circulate blood effectively, which can significantly decrease the chances of survival following a cardiac arrest. The guideline of 2 inches is based on evidence that this depth allows for the best chance of achieving enough blood pressure in the coronary and cerebral arteries to sustain life until advanced medical help can arrive. Moreover, using a depth greater than 2 inches can potentially cause injury to the patient's ribs or other thoracic structures, which is why the emphasis is placed specifically on achieving the correct depth rather than exceeding it.

SAMPLE

10. If a child with an obstructed airway becomes unresponsive, what should you do first?

A. Immediately start providing rescue breaths

B. Call for emergency medical assistance

C. Gently lower the child to the ground and begin CPR

D. Attempt back blows and abdominal thrusts

In the event that a child with an obstructed airway becomes unresponsive, the priority is to ensure that the child receives immediate care to restore their breathing and circulation. Gently lowering the child to the ground and beginning CPR is the correct first action in this scenario. Specifically, when a child is unresponsive, the likelihood of cardiac arrest increases, especially if the airway has been obstructed long enough to cause a lack of oxygen. Starting CPR immediately can help maintain blood flow to vital organs and can be critical in preventing further complications. It's important to note that while rescue breaths are essential for a child in cardiac arrest, they should only be administered after ensuring circulation is established through chest compressions. Calling for emergency medical assistance is vital, but initiating CPR takes precedence because it directly addresses the immediate threat to the child's life. Attempting back blows and abdominal thrusts is appropriate when the child is conscious and responsive, as these methods help clear an obstruction in the airway; however, when the child is unresponsive, initiating CPR is the most crucial step.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://basiclifesupport.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE