BASIC LIFE SUPPORT INSTRUCTOR PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://basiclifesupportinstructor.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. How long should you deliver a breath during CPR?

A. 0.5 seconds

B. 1 second

C. 1.5 seconds

D. 2 seconds

2. What is the first step a rescuer should take in pediatric BLS before starting compressions?

A. Check for pulse

B. Verify scene safety

C. Activate emergency response

D. Start rescue breaths immediately

3. What is the first action to take if you witness a cardiac arrest?

A. Check for breathing

B. Start chest compressions

C. Call emergency services

D. Use an AED

4. If a patient in an opioid emergency has a pulse but is not breathing normally, what should you do?

A. Start chest compressions

B. Provide breaths and give naloxone

C. Call for immediate medical help

D. Administer oxygen only

5. What should be done immediately after calling 911?

A. Start CPR if the person is unresponsive and not breathing

B. Wait for emergency services to arrive

C. Check social media

D. Provide a snack to bystanders

SAMPLE

6. How often should the compressor rotate during BLS resuscitation?

A. Every minute

B. Every 5 minutes

C. Every 2 minutes

D. Every 3 minutes

7. What characterizes pulseless ventricular tachycardia (pVT)?

A. The heart beats too slowly

B. The ventricles contract at an extremely rapid rate

C. The heart exhibits a chaotic electrical activity

D. The patient appears responsive

8. Which is NOT a requirement to pass the BLS course?

A. Participate and pass hands-on skills

B. Pass the adult CPR and AED skills

C. Pass the infant CPR skills test

D. Complete a written exam with a score of 70%

9. If a pediatric patient has a heart rate less than 60 and signs of poor perfusion, what is the next step?

A. Continue rescue breaths

B. Monitor until help arrives

C. Start CPR immediately

D. Check pulse every two minutes

10. What technique is preferred for performing infant compressions when using two rescuers?

A. 2 finger technique

B. 2 thumb encircling hands

C. Single hand technique

D. Heel of hand technique

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. B
5. A
6. C
7. B
8. D
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. How long should you deliver a breath during CPR?

A. 0.5 seconds

B. 1 second

C. 1.5 seconds

D. 2 seconds

During CPR, delivering a breath should take approximately 1 second. This duration is recommended because it allows sufficient time for air to enter the lungs of the victim without causing excessive pressure, which can lead to complications such as gastric inflation. When providing rescue breaths, it's important to ensure that each breath is effective, which is assessed by observing the rise and fall of the chest. Taking about 1 second for each breath offers a balance that is neither too short, which may not adequately fill the lungs, nor too long, which could disrupt the flow of chest compressions and impede blood circulation to vital organs. Thus, the 1-second guideline helps maintain the rhythm of CPR while ensuring ventilation is effective.

2. What is the first step a rescuer should take in pediatric BLS before starting compressions?

A. Check for pulse

B. Verify scene safety

C. Activate emergency response

D. Start rescue breaths immediately

The correct answer is verifying scene safety, as this is a fundamental step in any emergency response situation, including pediatric Basic Life Support (BLS). Before a rescuer approaches and begins any lifesaving measures, it is crucial to ensure that the environment is safe for both the rescuer and the victim. This means checking for potential hazards such as traffic, fire, toxic substances, or any other dangers that could further complicate the situation or lead to additional injuries. Scene safety helps set a secure environment that allows effective intervention. Once the scene is confirmed to be safe, the rescuer can then proceed with assessing the patient, checking for responsiveness, activating emergency response, and delivering compressions and breaths if necessary. Understanding the importance of verifying scene safety is vital for effective emergency response, as it prioritizes the well-being of both the rescuer and the person in need of assistance.

3. What is the first action to take if you witness a cardiac arrest?

A. Check for breathing

B. Start chest compressions

C. Call emergency services

D. Use an AED

When you witness a cardiac arrest, the first action to take is to ensure that emergency services are notified immediately. This is crucial because the chances of survival decrease significantly with each passing minute during a cardiac arrest event. Calling emergency services alerts ambulance personnel to the situation, allowing them to mobilize and arrive to provide advanced care as soon as possible. While checking for breathing, starting chest compressions, and using an AED are vital components of the response to cardiac arrest, they should be performed promptly after ensuring that help is on the way. Once emergency services are called, the responder can then proceed to check for breathing and begin chest compressions or use an AED if available. This prompt alerting of emergency services lays the groundwork for comprehensive care and optimal patient outcomes.

SAMPLE

4. If a patient in an opioid emergency has a pulse but is not breathing normally, what should you do?

A. Start chest compressions

B. Provide breaths and give naloxone

C. Call for immediate medical help

D. Administer oxygen only

In the scenario where a patient is in an opioid emergency and has a pulse but is not breathing normally, the primary course of action is to provide breaths and administer naloxone. This approach addresses both the immediate need for ventilation and the underlying issue with opioid overdose. When someone is experiencing an opioid overdose, their respiratory system is often severely depressed, leading to inadequate or absent breathing. While the patient may still have a pulse, their lack of effective respiration poses a significant risk of hypoxia and potential brain damage. Providing rescue breaths helps to ensure that oxygen is being delivered to the lungs and subsequently the bloodstream, which is critical in preventing further complications. Additionally, administering naloxone is crucial because it is an opioid antagonist that can rapidly reverse the effects of opioid overdose, restoring normal breathing and consciousness. It is important to act quickly, as the combination of providing breaths and the administration of naloxone can significantly improve the chance of a positive outcome for the patient. Calling for immediate medical help is also a necessary part of the response, but it should be done in tandem with providing breaths and naloxone, rather than solely relying on calling for help. Administering oxygen alone does not address the critical need for immediate respiratory assistance and the action of nal

5. What should be done immediately after calling 911?

A. Start CPR if the person is unresponsive and not breathing

B. Wait for emergency services to arrive

C. Check social media

D. Provide a snack to bystanders

Immediately after calling 911, starting CPR is a crucial step if the person is unresponsive and not breathing. This action is essential because CPR can help maintain blood circulation and oxygenation to the vital organs until professional help arrives. Time is a critical factor in cardiac arrest situations; the sooner CPR is initiated, the higher the chances of survival and a better neurological outcome for the victim. By beginning CPR right after notifying emergency services, you are actively contributing to the victim's chance of survival. It demonstrates the urgency of the situation. Actions like waiting for emergency services, checking social media, or providing snacks to bystanders distract from the immediate need for medical assistance and neglect the foundational purpose of calling for help in the first place. Therefore, starting CPR aligns with the best practices for life-saving interventions in emergencies.

SAMPLE

6. How often should the compressor rotate during BLS resuscitation?

A. Every minute

B. Every 5 minutes

C. Every 2 minutes

D. Every 3 minutes

During Basic Life Support (BLS) resuscitation, the compressor should rotate every 2 minutes to optimize the effectiveness of chest compressions. This rotation allows for the fatigue factor to be addressed, as high-quality, continuous chest compressions are essential for maintaining blood flow to the heart and brain. When rescuers switch every 2 minutes, it helps to ensure that compressions remain effective and consistent in depth and rate, which is crucial for the victim's chances of survival. Fatigue can lead to a decrease in compression quality, including insufficient depth or pauses in compressions, which can be detrimental during a cardiac arrest scenario. Regular rotation every 2 minutes aligns with the recommended guidelines set forth by resuscitation authorities and promotes teamwork, allowing for a more efficient response in emergency situations.

7. What characterizes pulseless ventricular tachycardia (pVT)?

A. The heart beats too slowly

B. The ventricles contract at an extremely rapid rate

C. The heart exhibits a chaotic electrical activity

D. The patient appears responsive

Pulseless ventricular tachycardia (pVT) is characterized by the ventricles contracting at an extremely rapid rate. In this condition, the heart's electrical signals cause the ventricles to beat quickly and ineffectively, typically at rates greater than 100 beats per minute. This rapid contraction does not allow for adequate filling of the ventricles or effective pumping of blood, leading to a lack of pulse and, ultimately, insufficient blood flow to the body's organs. The high rate of ventricular contractions in pVT can lead to severe hemodynamic instability and is a life-threatening condition that requires immediate intervention. The recognition of this rhythm is crucial for the initiation of appropriate treatment, such as cardiopulmonary resuscitation (CPR) and possible defibrillation, to restore a normal heart rhythm and improve chances of survival. While other choices reference different conditions, they do not accurately describe the specific characteristics of pVT. The heart does not beat too slowly in this case, nor does it exhibit chaotic electrical activity, which is more characteristic of conditions like ventricular fibrillation. Additionally, a patient in pVT is typically unresponsive due to the lack of effective circulation, making it impossible for them to appear responsive.

8. Which is NOT a requirement to pass the BLS course?

A. Participate and pass hands-on skills

B. Pass the adult CPR and AED skills

C. Pass the infant CPR skills test

D. Complete a written exam with a score of 70%

To successfully complete the Basic Life Support (BLS) course, a participant typically must demonstrate proficiency in essential skills and understand the theoretical knowledge necessary for effective response to cardiac emergencies. While it is important to pass hands-on skills, particularly those related to adult CPR and AED, as well as infant CPR, these practical components are crucial for real-life application. Although a written exam is commonly part of many certification programs, it's important to note that the focus in this case is on demonstrating practical skills and knowledge relevant to life-saving techniques. Thus, while passing a written exam may be beneficial in confirming understanding of the material, the actual requirements for passing the course center more on hands-on performance. As such, successfully completing the theoretical part of the course with a score of 70% is often not a strict requirement for passing the BLS course itself, making this aspect less critical than the demonstrated skills.

SAMPLE

9. If a pediatric patient has a heart rate less than 60 and signs of poor perfusion, what is the next step?

A. Continue rescue breaths

B. Monitor until help arrives

C. Start CPR immediately

D. Check pulse every two minutes

When a pediatric patient presents with a heart rate of less than 60 beats per minute accompanied by signs of poor perfusion, it indicates a critical situation that requires immediate intervention. In children, a heart rate below this threshold, especially in the presence of poor perfusion, suggests that the patient may be experiencing severe bradycardia or potentially cardiac arrest, both of which necessitate urgent action to restore adequate blood flow and oxygenation. Starting CPR immediately is crucial because it helps maintain circulation to vital organs, increasing the chances of survival and recovery. CPR provides artificial respiration and circulation, which is vital when spontaneous heartbeats are inadequate to sustain life. In this scenario, other actions such as monitoring until help arrives or continuing rescue breaths may not be sufficient and could delay necessary resuscitation efforts. Checking the pulse every two minutes would also not provide timely intervention needed in such an emergent context. Therefore, initiating CPR right away is the most appropriate step to take in order to improve the chances of a positive outcome for the patient.

10. What technique is preferred for performing infant compressions when using two rescuers?

A. 2 finger technique

B. 2 thumb encircling hands

C. Single hand technique

D. Heel of hand technique

The preferred technique for performing infant compressions with two rescuers is the two thumb encircling hands method. This technique provides optimal support and allows for better coordination between the rescuers. In this approach, one rescuer uses their thumbs to perform compressions on the infant's sternum while their fingers encircle the infant's back, providing stability and allowing the rescuer to maintain control of the infant's position. Using the two thumb encircling hands technique not only enhances the effectiveness of the compressions by delivering more force due to the use of both thumbs but also reduces the risk of injury to the infant's chest compared to other methods. This technique also allows for easy transition to rescue breaths, facilitating a smoother flow of care during resuscitation. Other methods of performing compressions, such as the 2 finger technique, while useful in some situations, do not offer the same level of effectiveness and safety as the two thumb encircling hands approach when two rescuers are involved. The single hand technique and heel of hand technique are not recommended for infants, as they may not provide sufficient force for effective compressions on such a small and delicate body. Therefore, the two thumb encircling hands method stands out as the most effective and safest

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://basiclifesupportinstructor.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE