BONENT CONTINUING DIALYSIS EDUCATION (CDC) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://bonentcdc.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which of the following is NOT a key step in the research process?

A. Analyzing data

B. Developing the study design

C. Eliminating unusual variation

D. Preparing the final report before the analysis

2. What contributes to the total pressure in the blood circuit during dialysis?

A. Resistance to flow and blood pump control

B. Patient heart rate and diet

C. Machine calibration and technician skill

D. Temperature and humidity of the environment

3. Which of the following is a common complication of untreated high blood pressure in dialysis patients?

A. Cognitive impairment

B. Severe anemia

C. Bone fractures

D. Cardiovascular complications

4. Why are patients on dialysis often prescribed vitamin D?

A. To reduce cholesterol levels

B. To manage mineral and bone disorders due to reduced kidney function

C. To boost immune system strength

D. To prevent hair loss

5. Which electrolyte imbalance is commonly monitored in dialysis patients?

A. Calcium levels

B. Sodium levels

C. Magnesium levels

D. Potassium levels

SAMPLE

6. What are the recommended management strategies for pericarditis?

A. Increased frequency of dialysis, use of anti-inflammatory drugs, controlled heparinization

B. Decreased fluid intake, increased physical activity, use of corticosteroids

C. Immediate surgery, complete bed rest, avoidance of infections

D. Daily monitoring of vital signs, use of iron supplements, routine x-rays

7. What happens when the hemodialysis machine detects a blood leak?

A. The machine continues operating normally

B. An alarm sounds and the blood pump will stop

C. The treatment is paused for patient assessment

D. The machine notifies the technician only in silent mode

8. What is the purpose of pre-treatment assessments in dialysis?

A. To evaluate patient health status and tailor dialysis treatment plans

B. To decrease treatment time

C. To limit the number of staff required

D. To determine dietary preferences

9. What can cause hemolysis of red blood cells during dialysis?

A. Hyperthermic fluid delivered to the dialyzer

B. Cold saline solution used in treatment

C. High levels of oxygen in the dialysate

D. Rapid infusion of medication

10. What is the importance of monitoring ultrafiltration rates during hemodialysis?

A. To ensure adequate medication administration

B. To assess the risk of blood clots

C. To ensure adequate fluid removal and avoid complications

D. To monitor electrolyte levels

SAMPLE

Answers

SAMPLE

1. D
2. A
3. D
4. B
5. D
6. A
7. B
8. A
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. Which of the following is NOT a key step in the research process?

A. Analyzing data

B. Developing the study design

C. Eliminating unusual variation

D. Preparing the final report before the analysis

In the research process, preparation of the final report is typically done after data analysis has been completed. This sequence is crucial because the report needs to reflect the findings and insights derived from interpreting the analyzed data. Preparing the report before analyzing the data would not allow the researcher to include accurate conclusions or recommendations based on the actual results of the study. Additionally, the research process is structured to include essential steps such as developing a study design to define the methodology and objectives clearly and analyzing data to draw meaningful conclusions. Eliminating unusual variation may also be a critical step to ensure the reliability and validity of research findings. However, the preparation of a final report must follow the data analysis to accurately represent the study's outcomes.

2. What contributes to the total pressure in the blood circuit during dialysis?

A. Resistance to flow and blood pump control

B. Patient heart rate and diet

C. Machine calibration and technician skill

D. Temperature and humidity of the environment

The total pressure in the blood circuit during dialysis is influenced primarily by factors that affect fluid dynamics, including the resistance to flow and the control of the blood pump. Resistance to flow is generated by various components of the system, such as the bloodlines, filters, and any narrowing in the tubing or connections. When resistance increases, the blood pump must work harder to maintain adequate flow, which in turn affects the overall pressure in the circuit. Additionally, the blood pump control plays a crucial role in regulating the flow rate of blood through the dialysis machine. If the pump is set to a higher flow rate, it will exert greater pressure within the circuit to overcome any resistance encountered. Thus, the interplay between the resistance to flow and the pump settings directly determines the total pressure experienced within the blood circuit during a dialysis session. The other options focus on factors that have little impact on the pressure within the circuit itself during dialysis. Patient heart rate and diet, although important for overall health and treatment effectiveness, do not directly influence the pressure in the dialysis machine's blood circuit. Machine calibration and technician skill are more concerned with ensuring the machine operates effectively and safely rather than directly contributing to pressure levels. Lastly, environmental conditions like temperature and humidity can affect the operation of a

SAMPLE

3. Which of the following is a common complication of untreated high blood pressure in dialysis patients?

A. Cognitive impairment

B. Severe anemia

C. Bone fractures

D. Cardiovascular complications

In dialysis patients, untreated high blood pressure often leads to significant cardiovascular complications. Chronic hypertension can cause increased workload on the heart, leading to hypertrophy of the heart muscle, heart failure, and an elevated risk of both heart attacks and strokes. In the context of dialysis, where patients may already have underlying cardiovascular risks due to kidney failure, the impact of high blood pressure can be even more pronounced. Managing blood pressure effectively is critical for reducing the risk of these serious outcomes. Cognitive impairment may occur as a long-term consequence of cardiovascular issues or could be associated with other factors like uremia. Severe anemia is more closely related to the kidneys' inability to produce adequate erythropoietin, rather than directly from high blood pressure. Bone fractures can occur in dialysis patients due to mineral and bone disorders associated with chronic kidney disease, but they are not a direct consequence of untreated hypertension. Thus, cardiovascular complications stand out as the direct and immediate consequence of poorly managed high blood pressure in these patients.

4. Why are patients on dialysis often prescribed vitamin D?

A. To reduce cholesterol levels

B. To manage mineral and bone disorders due to reduced kidney function

C. To boost immune system strength

D. To prevent hair loss

Patients on dialysis are often prescribed vitamin D primarily to manage mineral and bone disorders that arise due to reduced kidney function. When kidneys are not functioning properly, they struggle to convert vitamin D into its active form, which is crucial for the absorption of calcium and phosphate in the body. This disruption leads to imbalances in calcium and phosphate levels, which can result in conditions such as osteodystrophy—a common issue among patients with kidney disease. When vitamin D is supplemented, it helps regulate these minerals effectively, promotes bone health, and mitigates the risk of fractures. Additionally, it can help in maintaining appropriate parathyroid hormone levels, reducing the risk of secondary hyperparathyroidism, a potential complication in patients with chronic kidney disease. Thus, the administration of vitamin D is significant for maintaining bone mineral density and preventing the myriad of bone-related issues associated with renal failure and dialysis treatment.

SAMPLE

5. Which electrolyte imbalance is commonly monitored in dialysis patients?

A. Calcium levels

B. Sodium levels

C. Magnesium levels

D. Potassium levels

In dialysis patients, monitoring potassium levels is particularly critical due to the risk of hyperkalemia, which can lead to serious cardiac complications. The kidneys typically help regulate potassium balance by excreting excess levels in urine. However, patients with kidney failure are unable to effectively eliminate potassium, leading to accumulations in the bloodstream. This can result in potentially unsafe levels, causing life-threatening conditions such as arrhythmias or cardiac arrest. During dialysis treatments, potassium levels can fluctuate significantly, so healthcare providers must closely monitor these levels to ensure they remain within a safe range. The manipulation of dialysate composition can also be used to manage potassium levels during treatment, making it a vital aspect of care for dialysis patients. While calcium, sodium, and magnesium levels are important in the management of dialysis patients, the immediate and significant threat posed by elevated potassium makes it a priority in monitoring during dialysis treatment.

6. What are the recommended management strategies for pericarditis?

A. Increased frequency of dialysis, use of anti-inflammatory drugs, controlled heparinization

B. Decreased fluid intake, increased physical activity, use of corticosteroids

C. Immediate surgery, complete bed rest, avoidance of infections

D. Daily monitoring of vital signs, use of iron supplements, routine x-rays

The recommended management strategies for pericarditis primarily focus on addressing inflammation, fluid management, and preventing complications. Increasing the frequency of dialysis is beneficial in this context because patients with pericarditis often have underlying conditions such as chronic kidney disease, where fluid overload is a concern. Dialysis helps to remove excess fluid that can accumulate and exacerbate pericardial effusion, thereby reducing pressure on the heart. Using anti-inflammatory drugs is a cornerstone of treating pericarditis. Nonsteroidal anti-inflammatory drugs (NSAIDs) or sometimes colchicine are commonly administered to reduce inflammation in the pericardium and alleviate symptoms. Controlled heparinization may be part of the management plan, especially if the pericarditis is related to anticoagulation therapy, but it must be done judiciously to avoid complicating the patient's condition. In contrast, the other strategies listed do not directly address the specific needs associated with pericarditis management. Options involving decreased fluid intake or increased physical activity may not recognize the potential for fluid overload and strain on the heart. Immediate surgery is typically reserved for cases involving severe complications like tampons or constrictive pericarditis. Routine monitoring and using iron supplements or x-rays do not directly

SAMPLE

7. What happens when the hemodialysis machine detects a blood leak?

A. The machine continues operating normally

B. An alarm sounds and the blood pump will stop

C. The treatment is paused for patient assessment

D. The machine notifies the technician only in silent mode

When the hemodialysis machine detects a blood leak, it is crucial for patient safety and effective treatment that the machine responds appropriately. The correct response involves the activation of alarms to alert staff to the situation. Specifically, when a blood leak is detected, the machine will sound an alarm and automatically stop the blood pump. This immediate action is designed to prevent any potential blood loss and protect the patient from further complications related to a blood leak. Ensuring that the blood pump stops is critical because it prevents the patient's blood from entering any area of the machine that may have been compromised by the leak, thus safeguarding against infection and ensuring the integrity of the treatment process. The alarm serves as an important signal for healthcare providers to take appropriate action and assess the situation, making it a crucial part of maintaining safe dialysis procedures. Other options do not correctly reflect the machine's designed response to a blood leak. Continuing operations normally or notifying the technician only in silent mode would pose significant risks to patient safety. While patient assessment is necessary, it occurs in response to the alarm and stop actions, not as an initial step without the machine's alerts.

8. What is the purpose of pre-treatment assessments in dialysis?

A. To evaluate patient health status and tailor dialysis treatment plans

B. To decrease treatment time

C. To limit the number of staff required

D. To determine dietary preferences

The purpose of pre-treatment assessments in dialysis is fundamentally to evaluate the patient's health status and tailor dialysis treatment plans accordingly. This process ensures that healthcare providers can gather comprehensive information about the patient's medical history, current health, and specific needs. By thoroughly understanding the patient's condition, practitioners can make informed decisions on the best course of treatment, which may include customizing the frequency and type of dialysis, managing medications, and addressing any concurrent health issues. The focus on a patient-centered approach in these assessments allows for better clinical outcomes and can enhance the overall quality of life for the patient. This step is crucial in identifying potential complications, ensuring appropriate interventions, and ultimately providing safer and more effective dialysis therapy. Other options, such as decreasing treatment time, limiting staff requirements, or determining dietary preferences, do not encompass the primary goal of pre-treatment assessments. While these factors may have relevance in the broader context of dialysis management, they do not inherently capture the essential purpose of establishing a thorough understanding of the patient's health status to enhance treatment effectiveness.

SAMPLE

9. What can cause hemolysis of red blood cells during dialysis?

A. Hyperthermic fluid delivered to the dialyzer

B. Cold saline solution used in treatment

C. High levels of oxygen in the dialysate

D. Rapid infusion of medication

Hemolysis of red blood cells during dialysis can indeed occur when hyperthermic (excessively hot) fluid is delivered to the dialyzer. This situation can lead to an increase in the temperature of the blood, which can disrupt the membrane integrity of red blood cells, ultimately causing them to rupture or break down. The sensitivity of red blood cells to temperature changes is an important factor in maintaining their viability. High temperatures can impair the ability of the cells to maintain their shape and functionality, leading to hemolysis. In the context of the other options, the use of cold saline solutions typically does not promote hemolysis; in fact, lower temperatures can help protect red blood cells from damage. High levels of oxygen in the dialysate may pose other risks, but they are not directly correlated with hemolysis in the same way as temperature. Rapid infusion of medications can introduce various complications in dialysis treatment, but it is not specifically linked to the direct hemolysis of red blood cells as hyperthermic fluid is. Thus, understanding the impact of temperature on red blood cell integrity is crucial in preventing hemolysis during dialysis.

10. What is the importance of monitoring ultrafiltration rates during hemodialysis?

A. To ensure adequate medication administration

B. To assess the risk of blood clots

C. To ensure adequate fluid removal and avoid complications

D. To monitor electrolyte levels

Monitoring ultrafiltration rates during hemodialysis is crucial to ensure adequate fluid removal and to avoid complications related to fluid overload. During the dialysis session, ultrafiltration refers to the process of removing excess fluid from the patient's bloodstream. If the ultrafiltration rate is too low, patients may not have enough fluid removed, which can lead to complications such as hypertension, pulmonary edema, or heart failure. Conversely, if the rate is too high, it can cause dehydration, cramping, or hypotension, particularly if large volumes of fluid are removed too quickly. Maintaining the appropriate ultrafiltration rate is essential for optimizing patient safety and comfort during treatment, as well as ensuring effective management of fluid balance. This is why it is considered one of the key components to monitor throughout the hemodialysis process. Being aware of and adjusting the ultrafiltration rate helps clinicians prevent potential complications that would negatively affect patient health outcomes.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://bonentcdc.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE