

BOVINE UPPER MUSCULOSKELETAL APPARATUS AND SMALL RUMINANT DISEASE PRACTICE TEST (SAMPLE)

STUDY GUIDE



**SAMPLE STUDY GUIDE
WITH LIMITED QUESTIONS**

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://bovineuppermskapparatus.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Rectal examination findings may aid in identifying which condition?**
 - A. Spondylosis
 - B. Intervertebral disc disease
 - C. Meningioma
 - D. Fractured ilium
- 2. Which of the following is NOT among the four major spine conditions of cattle?**
 - A. Physitis of vertebral bodies
 - B. Congenital (hemivertebrae; congenital arthrogryposis)
 - C. Luxations
 - D. Fractured tibia
- 3. OC/OCD usually affects the fastest growing animals in a group and usually is in individuals, not a herd.**
 - A. True
 - B. False
 - C. True, but only in herd outbreaks
 - D. False; affects groups equally
- 4. Which splint has controversial use in cruciate ligament tears and immobilizes for 6-8 weeks followed by an additional 4 months sexual rest, but risks additional atrophy of the limb?**
 - A. Robert Jones splint
 - B. Ehmer sling
 - C. Walker splint
 - D. Cast
- 5. Regarding spastic syndrome in cattle, which statement is correct?**
 - A. It has onset at 3-7 years, inheritance is unclear, and there are no postmortem lesions.
 - B. It begins at birth and has a defined autosomal dominant inheritance.
 - C. It presents with fever and rapid decline.
 - D. It is caused by a bacterial infection leading to lameness.

- 6. Which statement about contagious ovine digital dermatitis (CODD) is true?**
- A. CODD is noncontagious and usually localized to a single foot.
 - B. CODD occurs in cattle.
 - C. CODD always resolves without treatment.
 - D. CODD often involves chronic hoof horn separation starting at the coronary band and can be highly contagious.
- 7. Describe the signs and initial management of olecranon bursitis in cattle.**
- A. Elbow swelling that is always warm and painful with fever; management includes immediate amputation
 - B. Elbow swelling that is non-painful and cold; management with warm compresses
 - C. Elbow swelling that may be sterile (non-painful) or septic (warm, painful, fever); management includes aspiration, lavage, systemic antibiotics if septic, and rest
 - D. Elbow swelling with no impact on limb function; no treatment needed
- 8. Describe the ballottement test in forelimb assessment and what a positive finding indicates.**
- A. Gentle joint compression to assess for palpable displacement or fluctuation; a positive test suggests joint effusion or instability.
 - B. Palpation of digital pulses
 - C. Testing flexion/extension range of motion only
 - D. Ultrasound assessment of joint depth
- 9. In CAE, which joint involvement is typical and what type of synovitis is seen?**
- A. One or both carpi with proliferative synovitis
 - B. Hocks with fibrous synovitis
 - C. Stifle with degenerative synovitis
 - D. Carpi and hocks with proliferative synovitis
- 10. Which factor is NOT associated with OC/OCD?**
- A. Regular, balanced exercise and nutrition
 - B. Rapid growth; Grain-rich diet; High P:Ca ratio; Abnormal biomechanics
 - C. Vitamin deficiencies as sole cause
 - D. Infectious causes

Answers

SAMPLE

1. A
2. D
3. A
4. C
5. A
6. D
7. C
8. A
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. Rectal examination findings may aid in identifying which condition?

- A. Spondylosis**
- B. Intervertebral disc disease
- C. Meningioma
- D. Fractured ilium

Rectal examination can reveal vertebral changes along the spine when you assess the ventral midline of the lumbar region. In spondylosis, there are osteophyte formations and bony proliferations along the vertebral bodies that can feel like irregular, firm ridges during per-rectal palpation. This makes rectal findings a clue to a degenerative spinal process rather than to pelvic or intracranial conditions. Intervertebral disc disease usually presents with neurologic deficits and pain that are not specifically diagnosed by rectal palpation. Meningioma is a tumor of the meninges that would show signs of neural involvement but not a characteristic per-rectal vertebral change. Fractured ilium is a pelvic injury with pain and instability in the pelvic region, which rectal palpation does not characteristically reveal as the vertebral changes do.

2. Which of the following is NOT among the four major spine conditions of cattle?

- A. Physitis of vertebral bodies
- B. Congenital (hemivertebrae; congenital arthrogryposis)
- C. Luxations
- D. Fractured tibia**

Understanding which problems involve the vertebral column versus other bones is key here. In cattle, spine-related conditions you'd expect include physitis of the vertebral bodies (growth-plate inflammation within the vertebrae), congenital vertebral anomalies such as hemivertebrae and related arthrogryposis, and luxations of spinal joints where the vertebrae become dislocated. These affect the spine and can cause back pain, deformity, or neurological signs. A fracture of the tibia, however, is a fracture of a long bone in the leg and does not involve the spine, so it is not among the spine conditions.

3. OC/OCD usually affects the fastest growing animals in a group and usually is in individuals, not a herd.

- A. True**
- B. False
- C. True, but only in herd outbreaks
- D. False; affects groups equally

OC/OCD is a developmental orthopedic issue tied to rapid growth. In young cattle and small ruminants, the most pronounced lesions arise when animals are growing fastest because the process of endochondral ossification in the developing joints is stressed and can falter, leading to cartilage flaps or osteochondral defects in weight-bearing joints such as the shoulder, elbow, knee (stifle), or fetlock. Because growth rate varies among individuals within the same group, affected animals tend to be the faster growers rather than every animal in the herd being affected. It isn't a contagious or herd-wide disease; it's a developmental problem that shows up more in those individuals experiencing the most rapid growth, with nutrition and genetics influencing the risk. So the statement aligns with the typical pattern of OC/OCD: fast growers are more at risk, and the condition is usually seen in individuals rather than uniformly across the herd.

4. Which splint has controversial use in cruciate ligament tears and immobilizes for 6-8 weeks followed by an additional 4 months sexual rest, but risks additional atrophy of the limb?

- A. Robert Jones splint
- B. Ehmer sling
- C. Walker splint**
- D. Cast

Immobilizing a cruciate ligament tear with a splint hinges on a difficult balance: the goal is to protect the healing tissues by limiting joint motion, but the cranial cruciate ligament has limited ability to heal reliably in the intra-articular environment, and long immobilization often leads to muscle atrophy and joint stiffness. The Walker splint is designed to immobilize the hind limb at the knee (stifle) to restrict movement around the cruciate area. In the scenario described, it's used to hold the stifle in a way that minimizes stress on the ligament for about 6–8 weeks, followed by an extended period of rest (around several months) to protect the healing tissues. The downside is that such prolonged disuse markedly increases the risk of disuse atrophy of the quadriceps and other limb muscles, which can complicate recovery and function even if the ligament heals. Other splints serve different purposes: the Ehmer sling is used primarily for hip luxation, not cruciate injuries; the Robert Jones splint is generally for distal limb fractures and soft-tissue injuries of the distal limb; and a cast (in a broader sense) is a general immobilization method but is not the specialized, knee-focused approach described here.

5. Regarding spastic syndrome in cattle, which statement is correct?

- A. It has onset at 3-7 years, inheritance is unclear, and there are no postmortem lesions.**
- B. It begins at birth and has a defined autosomal dominant inheritance.
- C. It presents with fever and rapid decline.
- D. It is caused by a bacterial infection leading to lameness.

Spastic syndrome in cattle is best described as a late-onset, likely hereditary neuromuscular condition. The key features are that it tends to start in adult animals, often around the 3–7 year range, and the inheritance pattern is not clearly defined or consistently reproducible. Postmortem examinations typically do not reveal defining tissue lesions, which supports a functional or neural-muscular origin rather than an inflammatory or infectious disease. This helps distinguish it from other scenarios. If it began at birth and followed a definite autosomal dominant pattern, you'd expect a clearer, codified hereditary transmission. If fever and rapid decline were present, that would suggest an acute infectious or systemic illness. If a bacterial infection caused lameness, you'd expect localized lesions or inflammatory changes, not a non-lesion-based spastic episode.

6. Which statement about contagious ovine digital dermatitis (CDD) is true?

- A. CDD is noncontagious and usually localized to a single foot.
- B. CDD occurs in cattle.
- C. CDD always resolves without treatment.
- D. CDD often involves chronic hoof horn separation starting at the coronary band and can be highly contagious.**

CDD is a contagious infectious hoof disease of sheep that presents with chronic hoof horn separation beginning at the coronary band. This starting point at the coronary band is a hallmark of CDD, and the condition tends to progress, causing progressive separation of the horn and lameness that can become chronic if not treated properly. The infection is highly transmissible within flocks and can spread between animals and feet through contaminated equipment or environment, making it much more than a single-foot problem. It is an ovine-specific disease, not something that primarily affects cattle. And it does not resolve on its own; management typically requires thorough foot care, hygiene to limit spread, and appropriate antimicrobial or topical treatments.

7. Describe the signs and initial management of olecranon bursitis in cattle.

- A. Elbow swelling that is always warm and painful with fever; management includes immediate amputation
- B. Elbow swelling that is non-painful and cold; management with warm compresses
- C. Elbow swelling that may be sterile (non-painful) or septic (warm, painful, fever); management includes aspiration, lavage, systemic antibiotics if septic, and rest**
- D. Elbow swelling with no impact on limb function; no treatment needed

Olecranon bursitis in cattle presents as swelling over the elbow that can be either sterile (non-painful) or septic (warm, painful with possible fever). Because the swelling can be due to non-infected inflammation or infection, the initial management must both relieve the accumulated fluid and address infection if present. Aspiration empties the bursa, providing relief and a sample to assess for infection. Lavage then cleans out inflammatory material and debris, reducing the bacterial load and irritation. If septic bursitis is suspected or confirmed, systemic antibiotics are needed to treat the infection beyond the local bursa. Resting the limb minimizes further irritation and supports healing, while good housing and analgesia improve comfort and recovery. This approach covers both sterile and septic scenarios and targets drainage, infection control when needed, and limb rest, which explains why it's the best choice.

8. Describe the ballottement test in forelimb assessment and what a positive finding indicates.

A. Gentle joint compression to assess for palpable displacement or fluctuation; a positive test suggests joint effusion or instability.

B. Palpation of digital pulses

C. Testing flexion/extension range of motion only

D. Ultrasound assessment of joint depth

Ballottement in the forelimb is a manual test for joint effusion and stability. The idea is to apply gentle pressure to the joint so the joint surfaces can "ballot" against each other. If there is excess fluid inside the joint or the ligaments are lax, the fluid and the moving surfaces will displace and then rebound when you release, producing a palpable fluctuation or a soft, fluid-filled wave. A normal joint resists this movement and feels relatively firm, so a distinct fluctuation indicates effusion or instability. This isn't about feeling pulses, testing how far the joint can bend, or imaging the joint depth. Those other actions assess different things (circulation, range of motion alone, or visualize anatomy), whereas ballottement specifically checks for intra-articular fluid and joint laxity.

9. In CAE, which joint involvement is typical and what type of synovitis is seen?

A. One or both carpi with proliferative synovitis

B. Hocks with fibrous synovitis

C. Stifle with degenerative synovitis

D. Carpi and hocks with proliferative synovitis

Caprine Arthritis Encephalitis most often causes chronic arthritis in the carpal joints, with one or both carpi commonly involved. The hallmark lesion is proliferative synovitis, meaning the synovial membrane becomes thickened with villous projections due to ongoing inflammatory stimulation from the lentiviral infection. While other joints can be affected, the carpi are the classic site and the proliferative pattern distinguishes CAE arthritis from other forms of synovitis, such as fibrous or degenerative types.

10. Which factor is NOT associated with OC/OCD?

A. Regular, balanced exercise and nutrition

B. Rapid growth; Grain-rich diet; High P:Ca ratio; Abnormal biomechanics

C. Vitamin deficiencies as sole cause

D. Infectious causes

OC/OCD arises when rapid skeletal growth in young cattle and small ruminants interacts with factors that load and stress the developing joints. The classic drivers are growth-promoting dietary and metabolic conditions, such as grain-rich diets that accelerate growth and create an imbalance in mineral supply (notably a high phosphorus-to-calcium ratio), along with abnormal biomechanics that place unusual stress on joints. Regular, balanced exercise and nutrition, on the other hand, does not contribute to these disturbances. In fact, it supports proper bone and cartilage development and is not a factor that predisposes to OC/OCD. While vitamin deficiencies or infectious causes can affect joint health in other contexts, they are not established primary causes of OC/OCD, which remains multifactorial but is best explained by accelerated growth and biomechanical/loading factors rather than a routine, well-balanced regimen.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://bovineuppermskapparat.us.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE