CAMLPR TRANSFUSION PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://camlprtransfusion.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Enchinococcus cysts, salmon roe, and pigeon egg whites neutralize which antibody?**

   A. Anti-Sda

   B. Anti-P1

   C. Anti-Lewis

   D. Anti-Chido/Rogers

2. **Which statement best describes the Indirect Antiglobulin Test (IAT)?**

   A. To detect in vitro sensitization of RBCs by antibodies

   B. To detect in vivo sensitization of RBCs

   C. To quantify bilirubin

   D. To measure hematocrit

3. **What does the genotype rr indicate in terms of C antigen status?**

   A. Homozygous for presence of C antigen

   B. Homozygous for absence of C antigen

   C. Heterozygous for C antigen

   D. No C antigen or D antigen

4. **What is the common dose of packed red blood cells per kilogram in pediatric patients?**

   A. 10-15 mL/kg per unit

   B. 5-7 mL/kg per unit

   C. 20-25 mL/kg per unit

   D. 1-2 mL/kg per unit

5. **Why is phenotyping performed for a patient who will receive daratumumab?**

   A. Daratumumab causes panagglutination in IAT testing, so phenotyping determines the antigen present on the RBCs to prevent sensitization and production of antibodies.

   B. To determine ABO group

   C. To determine the Rh type only

   D. To measure iron stores

SAMPLE

6. In antibody screen panel interpretation, what is the Rule of 3?

A. There must be 3 antigen-positive cells that react for the suspected antibody and 3 antigen-negative cells that don't react

B. There must be 3 positive controls

C. There must be 3 washes

D. There must be 3 negative screens

7. If RhC is negative, which antigen is absent from the red cells?

A. D antigen

B. C antigen

C. E antigen

D. c antigen

8. What are the benefits of leukoreduction in blood products?

A. Reduces febrile nonhemolytic transfusion reactions, decreases alloimmunization to HLA, and lowers risk of transmitting leukocyte-associated infections.

B. Increases the risk of infections.

C. Has no impact on transfusion reactions.

D. Only reduces bacterial contamination risk.

9. Based on saliva inhibition test results, what is the secretor status? Saliva + Anti-A + A cells: +; Saliva + Anti-B + B cells: +; Saliva + Anti-H + H cells: +.

A. Secretor

B. Non-secretor

C. Weak secretor

D. Bombay phenotype

10. What best describes a minor crossmatch, and when is it performed?

A. IS crossmatch when there is a positive antibody screen

B. Minor crossmatch uses enhanced AHG

C. The same as electronic crossmatch

D. IS crossmatch is used when there is a negative antibody screen and confirmation of two ABO groupings

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. A
5. B
6. A
7. B
8. A
9. B
10. D

SAMPLE

Explanations

SAMPLE

1. Enchinococcus cysts, salmon roe, and pigeon egg whites neutralize which antibody?

A. Anti-Sda

B. Anti-P1

C. Anti-Lewis

D. Anti-Chido/Rogers

Neutralization of antibodies by soluble antigen is being tested here, specifically the anti-P1 antibody. Anti-P1 is a naturally occurring antibody in the P blood group system that targets the P1 antigen on red cells. Soluble P1 antigen is present in echinococcus cyst fluid, salmon roe, and pigeon egg whites. When serum containing anti-P1 is exposed to these substances, the antibodies bind the soluble P1 antigen and become blocked from reacting with P1 on red cells. This absorption or neutralization confirms the antibody's identity because the antibody activity disappears after exposure to P1-containing material. The other antibodies have different targets and aren't predictably neutralized by these particular substances, so this pattern specifically points to anti-P1.

2. Which statement best describes the Indirect Antiglobulin Test (IAT)?

A. To detect in vitro sensitization of RBCs by antibodies

B. To detect in vivo sensitization of RBCs

C. To quantify bilirubin

D. To measure hematocrit

The main idea is that the indirect antiglobulin test detects antibodies in a patient's serum by causing them to bind to red cell antigens in vitro. In this test, patient plasma is incubated with test red cells so that any antibodies present can attach to those antigens. After washing away unbound stuff, anti-human globulin is added to reveal whether sensitization occurred; agglutination shows that antibodies capable of reacting with red cell antigens are present in the serum. This is an in vitro process, used for screening and identifying unexpected antibodies before transfusion. This differs from the direct antiglobulin test, which looks for antibodies already bound to red cells inside the patient (in vivo). Bilirubin quantification and hematocrit measurement are unrelated to detecting red cell antibodies.

3. What does the genotype rr indicate in terms of C antigen status?

A. Homozygous for presence of C antigen

B. Homozygous for absence of C antigen

C. Heterozygous for C antigen

D. No C antigen or D antigen

In the C/c gene system, whether the C antigen is present on red cells depends on the combination of alleles you inherit. A single dominant C allele is enough to express the C antigen, so CC or Cc individuals show the C antigen. Only when both alleles are recessive and there is no C allele at all do you fail to express C. Therefore, having two recessive alleles (rr) means no C antigen is produced on the red cells. This is what "homozygous for absence of C antigen" describes. The result is a C antigen–negative phenotype. (The D antigen status is a separate Rh factor and isn't determined by this same C/c locus.)

SAMPLE

4. What is the common dose of packed red blood cells per kilogram in pediatric patients?

A. 10-15 mL/kg per unit

B. 5-7 mL/kg per unit

C. 20-25 mL/kg per unit

D. 1-2 mL/kg per unit

In pediatric transfusion practice, packed red blood cells are given by weight to achieve a reliable improvement in oxygen-carrying capacity while avoiding fluid overload. The standard single-unit dose for children is about 10 to 15 mL per kilogram of body weight. This means a child's dose scales with size, so smaller children receive less total volume and larger children receive more, keeping the transfusion effective but not excessive. One unit of PRBC is typically around a few hundred milliliters, so a 20 kg child would receive roughly 200–300 mL when a full unit is given, aligned with that per-kilogram target. If the dose were much lower (e.g., 5–7 mL/kg), the transfusion might not raise hemoglobin enough to improve tissue oxygen delivery. If it were much higher (e.g., 20–25 mL/kg), the child could experience fluid overload or heart-lung strain. The 10–15 mL/kg range reflects a balance between achieving a meaningful hematologic benefit and maintaining hemodynamic safety.

5. Why is phenotyping performed for a patient who will receive daratumumab?

A. Daratumumab causes panagglutination in IAT testing, so phenotyping determines the antigen present on the RBCs to prevent sensitization and production of antibodies.

B. To determine ABO group

C. To determine the Rh type only

D. To measure iron stores

Daratumumab binds CD38 on red cells and can cause panagglutination in antibody screening tests, making it hard to see any underlying antibodies. Because the safety of transfusion hinges on knowing the patient's blood group, phenotype testing is performed to determine the ABO group so compatible blood can be chosen even when serologic testing is compromised. In other words, establishing the ABO phenotype before transfusion ensures you provide an ABO-compatible unit despite the interference from daratumumab.

6. In antibody screen panel interpretation, what is the Rule of 3?

A. There must be 3 antigen-positive cells that react for the suspected antibody and 3 antigen-negative cells that don't react

B. There must be 3 positive controls

C. There must be 3 washes

D. There must be 3 negative screens

In antibody screen panel interpretation, you confirm a specific antibody by showing a distinguishable pattern of reactions across multiple panel cells. The Rule of 3 means you should observe at least three panel cells that express the target antigen and react with the patient's serum, and at least three panel cells that lack that antigen and do not react. This combination provides stronger evidence that the antibody is truly present and specifically targeted, rather than a random or nonspecific reaction. Having three reactive cells helps establish that the antibody is consistently detecting the antigen, while three non-reactive cells help prove specificity by showing those without the antigen don't react. If you don't meet these thresholds, the result is not definitive and further testing or a larger panel may be needed. The other options describe procedural elements or different controls that aren't part of this interpretation rule.

SAMPLE

7. If RhC is negative, which antigen is absent from the red cells?

A. D antigen

B. C antigen

C. E antigen

D. c antigen

The RhC status tells you about the C antigen on the red cell surface. The Rh system includes D and the C/c and E/e antigen pairs; the term RhC specifically refers to the C antigen on the RhCE glycoprotein. If RhC is negative, the C antigen is not present on the red cells, while D and other Rh antigens can be present or absent independently. So the absent antigen is C. In practice, a RhC negative individual would require C-negative red cells to avoid alloimmunization.

8. What are the benefits of leukoreduction in blood products?

A. Reduces febrile nonhemolytic transfusion reactions, decreases alloimmunization to HLA, and lowers risk of transmitting leukocyte-associated infections.

B. Increases the risk of infections.

C. Has no impact on transfusion reactions.

D. Only reduces bacterial contamination risk.

Leukoreduction works by removing most donor white blood cells from blood products before transfusion, which lowers several reaction and infection risks that can follow a transfusion. First, febrile nonhemolytic transfusion reactions are largely driven by cytokines released by donor leukocytes and by recipient reactions to those leukocytes. By removing those white cells, the amount of triggering material is reduced, so fevers and related symptoms are less likely to occur. Second, exposure to donor leukocyte antigens can lead to alloimmunization against HLA antigens. Fewer donor leukocytes mean fewer opportunities for the recipient's immune system to encounter those HLA antigens, decreasing the chance of sensitization that can complicate future transfusions or organ or stem cell transplants. Third, leukocytes can carry certain infections, most notably CMV, and leukoreduction lowers the risk of transmitting these leukocyte-associated infections to recipients, especially those who are immunocompromised or CMV-seronegative. The other statements contradict what leukoreduction actually achieves: it does not increase infection risk, it does impact transfusion reactions, and its benefit is not limited to reducing bacterial contamination alone.

9. Based on saliva inhibition test results, what is the secretor status? Saliva + Anti-A + A cells: +; Saliva + Anti-B + B cells: +; Saliva + Anti-H + H cells: +.

A. Secretor

B. Non-secretor

C. Weak secretor

D. Bombay phenotype

Secretor status is decided by whether ABH substances are present in saliva. In a saliva inhibition test, these soluble ABH substances in secretions will block the binding of anti-A, anti-B, and anti-H antibodies to their corresponding red cells, leading to no clumping (inhibition) when the saliva is mixed with the antisera and the specific cells. Here, there is clumping with anti-A and A cells, with anti-B and B cells, and with anti-H and H cells. That lack of inhibition means the saliva did not contain ABH substances to neutralize the antibodies. So the individual does not secrete ABH antigens in saliva, which defines non-secretor status.

SAMPLE

10. What best describes a minor crossmatch, and when is it performed?

A. IS crossmatch when there is a positive antibody screen

B. Minor crossmatch uses enhanced AHG

C. The same as electronic crossmatch

D. IS crossmatch is used when there is a negative antibody screen and confirmation of two ABO groupings

A minor crossmatch checks donor plasma for antibodies that could react with the recipient's red cells. It is used when the recipient has a negative antibody screen (no detected alloantibodies) and both donor and recipient ABO groupings have been confirmed. Under these conditions, an immediate spin crossmatch can be performed to quickly verify that the donor's plasma antibodies will not target the recipient's red cells. This helps prevent a hemolytic reaction from donor antibodies in plasma-containing products. If there were antibodies detected in the recipient, or if the ABOs weren't confirmed, a different crossmatch approach would be required.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://camlprtransfusion.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE