CANADIAN RED CROSS STANDARD FIRST AID PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://canadianredcrossstdfirstaid.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What should you do first if someone is struggling to breathe due to asthma and their medication is not working?**

   A. Contact a family member

   B. Call EMS and get an AED

   C. Encourage deep breathing

   D. Wait for symptoms to improve

2. **Which of the following is NOT a sign or symptom of a mental concussion?**

   A. Clouded mindset

   B. Drowsiness

   C. Increased energy

   D. Slowed reaction times

3. **What describes a stroke?**

   A. Temporary loss of consciousness

   B. When blood flow to a muscle is repeated

   C. When blood flow to the brain is interrupted

   D. When the heart stops beating

4. **Which of the following describes 'complete choking'?**

   A. Can still breathe and talk

   B. Airway is completely blocked

   C. Person experiences anxiety and confusion

   D. Individual can cough forcefully

5. **Which of the following should NOT be done during a seizure?**

   A. Move nearby objects away

   B. Place a soft object under their head

   C. Try to hold the person down

   D. Roll the person onto their side after the seizure

6. **Which process is involved when air moves in and out of the lungs?**

   A. Catabolic reactions

   B. Neuromuscular transmission

   C. Contraction and relaxation of the diaphragm

   D. Cellular respiration

SAMPLE

7. What should you do if you are tired while performing CPR?

A. Stop immediately and rest

B. Continue until EMS arrives or signs of life appear

C. Switch responsibilities only if a second person arrives

D. Take a break every few minutes

8. Which symptom could indicate severe internal bleeding?

A. Fever

B. Soft tissue tenderness

C. Headache

D. Fatigue

9. What should be done if there is an impaled object in the eye?

A. Remove the object carefully with fingers

B. Call emergency medical services immediately

C. Wash the eye with soap and water

D. Apply ice to the eye

10. What should be done if someone is unresponsive and not breathing normally?

A. Continue monitoring until help arrives

B. Start CPR and administer naloxone as soon as possible

C. Keep them awake and talking

D. Call a friend for help

SAMPLE

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. C
6. C
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What should you do first if someone is struggling to breathe due to asthma and their medication is not working?

A. Contact a family member

B. Call EMS and get an AED

C. Encourage deep breathing

D. Wait for symptoms to improve

When someone is struggling to breathe due to an asthma attack and their medication is not providing relief, it is crucial to prioritize safety and ensure professional medical assistance is on the way. Calling Emergency Medical Services (EMS) is essential because they can provide advanced care and resources that may be beyond what a bystander can offer. An Automated External Defibrillator (AED) may also be helpful if the person's condition deteriorates to the point of requiring cardiac assistance. Although it might be tempting to take other actions like encouraging deep breathing or waiting for symptoms to improve, these options can delay critical intervention. Asthma can escalate quickly, leading to severe respiratory distress or a life-threatening situation, so immediate action is necessary. In such cases, seeking emergency help ensures that the affected person receives the best possible care as promptly as possible.

2. Which of the following is NOT a sign or symptom of a mental concussion?

A. Clouded mindset

B. Drowsiness

C. Increased energy

D. Slowed reaction times

In the context of a mental concussion, the signs and symptoms typically reflect a negative impact on cognitive function and alertness. A clouded mindset, drowsiness, and slowed reaction times are all indicative of the effects that a concussion can have on an individual's mental state. These symptoms are associated with cognitive impairment and decreased responsiveness, which are common after sustaining a concussion. In contrast, increased energy does not align with these typical symptoms. Rather than reflecting a state of cognitive impairment, increased energy suggests heightened alertness and function, which contradicts the expected signs of a concussion. Thus, identifying increased energy as not being a sign or symptom of a mental concussion is accurate, as it indicates the opposite of the cognitive difficulties that concussed individuals usually experience.

3. What describes a stroke?

A. Temporary loss of consciousness

B. When blood flow to a muscle is repeated

C. When blood flow to the brain is interrupted

D. When the heart stops beating

A stroke is defined as a medical condition that occurs when blood flow to the brain is interrupted. This interruption can happen due to a blockage in a blood vessel (ischemic stroke) or due to a rupture of a blood vessel, leading to bleeding in the brain (hemorrhagic stroke). When blood flow is compromised, brain cells begin to die, which can result in loss of functions controlled by the affected area of the brain, such as movement or speech. Recognizing the signs of a stroke and understanding its nature is crucial for prompt medical intervention, which can significantly improve outcomes for the affected individual. Other options refer to different medical situations. Temporary loss of consciousness relates to conditions such as fainting or seizures, and blood flow to a muscle would pertain more to issues like angina or muscle cramps rather than a brain-related condition. The stopping of the heart directly refers to cardiac arrest, which is a separate and life-threatening situation, distinct from the cerebral events that characterize a stroke.

EXAMPLE

4. Which of the following describes 'complete choking'?

A. Can still breathe and talk

B. Airway is completely blocked

C. Person experiences anxiety and confusion

D. Individual can cough forcefully

The description of 'complete choking' is characterized by a situation in which the airway is completely blocked. This condition prevents any air from flowing into or out of the lungs, making it impossible for the person to breathe, speak, or make any sound. In a state of complete choking, the individual typically displays signs of severe distress, including a panic response due to the inability to breathe. Recognizing complete choking is crucial because it requires immediate intervention, such as the Heimlich maneuver or calling for emergency medical assistance. When the airway is fully obstructed, the urgency of the situation escalates, as the lack of oxygen can lead to unconsciousness and potential fatality within a matter of minutes. In less severe cases, such as when a person can breathe, talk, or cough forcefully, the airway is only partially obstructed, and they might not need immediate emergency action. This distinction is essential for appropriate first aid response and interventions to be applied effectively.

5. Which of the following should NOT be done during a seizure?

A. Move nearby objects away

B. Place a soft object under their head

C. Try to hold the person down

D. Roll the person onto their side after the seizure

During a seizure, it is crucial to prioritize the safety of the individual experiencing the episode. Trying to hold the person down is not advisable, as it can lead to increased injury to both the person having the seizure and the individual attempting to provide assistance. The contractions and movements that occur during a seizure are involuntary, and forcing someone down can lead to muscle strains or broken bones, as well as potentially increase the risk of airway obstruction. On the other hand, moving nearby objects away can create a safer environment, and placing a soft object under the person's head helps to prevent head injuries from potential impacts with the ground. After the seizure has ended, rolling the person onto their side can be beneficial in keeping the airway clear and allowing any fluids to drain, reducing the risk of choking. Each of these actions supports the well-being of the person experiencing the seizure and follows best practices in first aid response.

SAMPLE

6. Which process is involved when air moves in and out of the lungs?

A. Catabolic reactions

B. Neuromuscular transmission

C. Contraction and relaxation of the diaphragm

D. Cellular respiration

The process that is involved when air moves in and out of the lungs is the contraction and relaxation of the diaphragm. During inhalation, the diaphragm contracts and moves downward, creating a vacuum that allows air to be drawn into the lungs. This increase in thoracic cavity volume decreases the pressure inside the lungs compared to the outside atmosphere, causing air to flow in. Conversely, during exhalation, the diaphragm relaxes and moves back up, pushing air out of the lungs as the thoracic cavity volume decreases and the pressure rises. Other processes mentioned do not directly describe the mechanism of air movement in and out of the lungs. For instance, catabolic reactions refer to metabolic processes that break down molecules for energy but do not pertain to pulmonary function. Neuromuscular transmission relates to the communication between nerves and muscles, which is indeed involved in the process but not the primary action affecting air movement itself. Cellular respiration is the process by which cells utilize oxygen to produce energy and is a metabolic process that occurs within cells after respiration has taken place, but it does not explain the mechanics of breathing.

7. What should you do if you are tired while performing CPR?

A. Stop immediately and rest

B. Continue until EMS arrives or signs of life appear

C. Switch responsibilities only if a second person arrives

D. Take a break every few minutes

When performing CPR, it is crucial to continue the process until Emergency Medical Services (EMS) arrives or the person shows signs of life. This is because the primary goal of CPR is to maintain blood circulation and oxygenation to vital organs, particularly the brain and heart, until professional help can take over or the individual shows recovery signs. If you stop CPR prematurely due to fatigue, it can result in irreversible damage or death, as the heart may remain in a state of arrhythmia or lack of rhythm during that critical time. Continuing CPR ensures that you are doing everything possible to save a life. While switching responsibilities can be beneficial when another trained individual arrives, the emphasis should remain on uninterrupted CPR until professional responders are on the scene or until the person regains consciousness and begins to breathe normally. Taking unnecessary breaks or stopping for rest directly contradicts the life-saving intent of CPR.

SAMPLE

8. Which symptom could indicate severe internal bleeding?

A. Fever

B. Soft tissue tenderness

C. Headache

D. Fatigue

Soft tissue tenderness can indeed indicate severe internal bleeding, particularly if it is accompanied by other symptoms. When severe internal bleeding occurs, it often leads to the accumulation of blood within the body, which can cause swelling and tenderness in the affected area. This tenderness results from the pressure that builds up from the blood pooling inside tissue cavities. In the context of severe internal bleeding, soft tissue tenderness is a critical symptom because it can help identify where the bleeding might be occurring. Other symptoms often seen with severe internal bleeding include rapid pulse, pale skin, confusion, or signs of shock, but soft tissue tenderness is particularly relevant in suggesting localized internal issues. While the other symptoms listed—fever, headache, and fatigue—can indicate various medical issues, they are not specific indicators of severe internal bleeding. Fever is commonly associated with infections, headache can stem from numerous issues like stress or dehydration, and fatigue could be a result of various conditions, not necessarily tied to internal bleeding. Therefore, recognizing soft tissue tenderness is essential in assessing the potential for severe internal injuries, warranting immediate medical attention.

9. What should be done if there is an impaled object in the eye?

A. Remove the object carefully with fingers

B. Call emergency medical services immediately

C. Wash the eye with soap and water

D. Apply ice to the eye

If there is an impaled object in the eye, the recommended action is to call emergency medical services immediately. This is crucial because the eye is an extremely delicate and sensitive organ, and improper handling of an impaled object could lead to further injury, increased bleeding, or permanent vision loss. Medical professionals are trained to handle such emergencies and can provide the necessary care in a controlled environment, ensuring the individual's safety and optimizing outcomes. Taking immediate action by contacting emergency services helps facilitate quick and appropriate medical intervention. It is also important to keep the individual calm and immobile, avoiding any attempts to remove the object or wash the eye, as these actions can exacerbate the injury and complicate treatment.

SAMPLE

10. What should be done if someone is unresponsive and not breathing normally?

A. Continue monitoring until help arrives

B. Start CPR and administer naloxone as soon as possible

C. Keep them awake and talking

D. Call a friend for help

When encountering someone who is unresponsive and not breathing normally, the appropriate action is to start CPR and, if available, administer naloxone as soon as possible. Immediate CPR is crucial because it helps maintain blood flow and oxygen to vital organs, especially the brain, until professional medical help can take over. Administering naloxone is particularly important in cases of an opioid overdose, as it can quickly reverse the effects of opioids and restore normal breathing. The combination of CPR and naloxone significantly increases the chance of survival and reduces the risk of long-term brain damage. Time is of the essence in these situations, and initiating these actions promptly is vital. Monitoring the individual without taking action does not address the life-threatening emergency at hand. Encouraging someone who is unresponsive to remain awake and talking is not appropriate, as their condition requires immediate medical attention, not distraction or stimulation. Simply calling a friend for help, while potentially helpful for additional support, does not provide the immediate intervention that is responsible for addressing unresponsiveness and abnormal breathing. Therefore, starting CPR and administering naloxone when indicated stands out as the critical, life-saving response in this scenario.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://canadianredcrossstdfirstaid.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE