CARDIAC-VASCULAR NURSING (CVRN) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cvrn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Why is troponin 1 preferred over other biomarkers for the detection of acute myocardial infarction (AMI)?

A. Is more specific to myocardial ischemic events

B. Remains elevated for 21 days

C. Rises immediately within the onset of ischemia

D. Is more predictive of transmural infarction

2. What should a cardiac-vascular nurse advise a patient who wishes to refuse an experimental cardiac drug?

A. The medication dose has been scheduled for administration.

B. The primary investigator will speak with the patient about his or her decision.

C. The patient has a right to refuse the medication and to withdraw from the study.

D. Participation in the study is critical to improving the quality of patient outcomes.

3. What characterizes right-sided heart failure?

A. Elevated pulmonary artery pressure

B. Reduced hepatic perfusion pressure

C. Increased systemic venous pressure

D. Decreased systemic arterial pressure

4. What is the most common type of cardiomyopathy?

A. Hypertrophic cardiomyopathy

B. Restrictive cardiomyopathy

C. Dilated cardiomyopathy

D. Arrhythmogenic right ventricular cardiomyopathy

5. In the case of a patient with negative troponins experiencing recurring chest pain, what is the cardiac-vascular nurse's next best action?

A. Activate the cardiac catheterization team

B. Administer analgesics

C. Obtain a 12-lead electrocardiogram

D. Promote relaxation and monitor the response

SAMPLE

6. Describe how peripheral vascular disease affects limb circulation.

A. It results in increased blood flow to the extremities.

B. It leads to reduced blood flow to the extremities, causing pain and potential tissue damage.

C. It has no significant impact on limb circulation.

D. It causes varicose veins exclusively.

7. Which lipid level is typically raised with lipid-lowering therapy?

A. Lipoprotein (a)

B. Triglycerides

C. High density lipoprotein cholesterol

D. Low density lipoprotein cholesterol

8. What indicates effective understanding of discharge teaching for a patient after coronary artery bypass surgery?

A. Asking whether smoking one cigarette per day is acceptable

B. Clarifying when cardiac rehabilitation will begin

C. Describing plans to relax in a hot tub during recovery

D. Planning to resume driving his or her spouse to work next week

9. For a patient with an abdominal aortic aneurysm undergoing medical treatment, what should be the nurse's priority?

A. Preparing for dialysis to correct renal failure

B. Restricting activities of daily living and low-level exercise

C. Monitoring the aneurysm size monthly with an endocardiogram

D. Maintaining mean blood pressure at a normal level

10. What benefit can a patient recovering from a myocardial infarction gain from meditation?

A. Increases sympathetic nervous system activity.

B. Decreases sympathetic nervous system activity.

C. Increases the release of catecholamines.

D. Decreases vasodilation.

SAMPLE

Answers

SAMPLE

1. A
2. C
3. C
4. C
5. C
6. B
7. C
8. B
9. D
10. B

SAMPLE

Explanations

SAMPLE

1. Why is troponin 1 preferred over other biomarkers for the detection of acute myocardial infarction (AMI)?

A. Is more specific to myocardial ischemic events

B. Remains elevated for 21 days

C. Rises immediately within the onset of ischemia

D. Is more predictive of transmural infarction

Troponin I is preferred for the detection of acute myocardial infarction because it is highly specific for myocardial tissue. Unlike other cardiac biomarkers, such as creatine kinase MB (CK-MB) or myoglobin, which can be elevated due to a variety of conditions affecting skeletal muscle or other tissues, troponin I is specifically found in cardiac muscle. This specificity allows clinicians to differentiate between cardiac and non-cardiac causes of elevated enzyme levels, making it a more reliable indicator of myocardial injury. Furthermore, troponin I levels increase significantly within a few hours of myocardial injury, making it a crucial tool in the early diagnosis of AMI. Its elevated levels can indicate damage sustained during the ischemic event, and it can persist in the bloodstream for a period of time, assisting in the assessment of myocardial injury over time, although it is especially useful in the initial stages of diagnosis. The other options, while presenting some facts about the biomarker's behavior, do not capture the primary reason for troponin I's preferred status in clinical practice—its high specificity to heart muscle injury.

2. What should a cardiac-vascular nurse advise a patient who wishes to refuse an experimental cardiac drug?

A. The medication dose has been scheduled for administration.

B. The primary investigator will speak with the patient about his or her decision.

C. The patient has a right to refuse the medication and to withdraw from the study.

D. Participation in the study is critical to improving the quality of patient outcomes.

The patient has the fundamental right to refuse any medical treatment, including participation in clinical trials for experimental drugs. This principle is rooted in the ethical guidelines surrounding informed consent, which emphasize that individuals should have the autonomy to make choices about their own healthcare. The right to withdraw from a study at any point, without facing penalties or losing access to standard care, is a critical aspect of clinical research ethics. By acknowledging the patient's right to refuse the medication and to withdraw from the study, the cardiac-vascular nurse ensures that the patient feels empowered and respected in their decision-making process regarding their treatment options. This response also reinforces the ethical responsibility of healthcare providers to prioritize patient autonomy while providing necessary information about the potential implications of their choice.

SAMPLE

3. What characterizes right-sided heart failure?

A. Elevated pulmonary artery pressure

B. Reduced hepatic perfusion pressure

C. Increased systemic venous pressure

D. Decreased systemic arterial pressure

Right-sided heart failure is characterized by an inability of the right ventricle to effectively pump blood to the lungs, which leads to a series of compensatory changes in the cardiovascular system. One of the hallmark features of this condition is increased systemic venous pressure. As the right side of the heart struggles to handle the volume of blood returning from the body, blood begins to back up in the systemic venous circulation. This elevation in pressure can lead to symptoms such as peripheral edema, ascites, and jugular venous distension, which are common manifestations of right-sided heart failure. Increased systemic venous pressure reflects the heart's inability to effectively manage the return of blood, leading to congestion in the venous system. This is a critical aspect of right-sided heart failure and is essential for understanding the clinical signs and symptoms associated with the condition.

4. What is the most common type of cardiomyopathy?

A. Hypertrophic cardiomyopathy

B. Restrictive cardiomyopathy

C. Dilated cardiomyopathy

D. Arrhythmogenic right ventricular cardiomyopathy

Dilated cardiomyopathy is recognized as the most common type of cardiomyopathy. It is characterized by the dilation of the heart chambers, particularly the left ventricle, which leads to impaired systolic function and reduced cardiac output. This condition can result from various factors, including ischemic heart disease, hypertension, genetic predispositions, and exposure to toxins or infections. The significance of recognizing dilated cardiomyopathy lies in its potential complications, such as heart failure and arrhythmias. Its prevalence makes it a critical condition for healthcare professionals, particularly in the context of assessing and managing patients with heart disease. Understanding dilated cardiomyopathy can guide effective diagnostic approaches and treatment strategies, improving patient outcomes.

5. In the case of a patient with negative troponins experiencing recurring chest pain, what is the cardiac-vascular nurse's next best action?

A. Activate the cardiac catheterization team

B. Administer analgesics

C. Obtain a 12-lead electrocardiogram

D. Promote relaxation and monitor the response

When a patient with negative troponins continues to experience recurring chest pain, the next best action for the cardiac-vascular nurse is to obtain a 12-lead electrocardiogram (ECG). This is crucial because the ECG can provide immediate information about the electrical activity of the heart. The ECG results can reveal changes such as ischemia, infarction, or arrhythmias that might not be indicated by troponin levels. Given that troponins are often markers for myocardial injury but not definitive for all causes of chest pain, conducting an ECG is necessary to assess the cardiac condition more thoroughly. Analyzing the ECG can lead to prompt identification of potentially serious issues that require intervention and can guide further management or diagnostic steps. This proactive approach ensures that any possible acute coronary syndrome or other cardiac anomalies are promptly addressed, thereby improving patient outcomes.

SAMPLE

6. Describe how peripheral vascular disease affects limb circulation.

A. It results in increased blood flow to the extremities.

B. It leads to reduced blood flow to the extremities, causing pain and potential tissue damage.

C. It has no significant impact on limb circulation.

D. It causes varicose veins exclusively.

Peripheral vascular disease (PVD) significantly impacts limb circulation by causing reduced blood flow to the extremities. This condition often results from atherosclerosis, which is the buildup of fatty deposits within the arteries. As these arteries become narrowed or blocked, the volume of blood that can be delivered to the limbs decreases. This reduced blood flow can lead to a range of symptoms, including intermittent claudication, which is characterized by pain or cramping in the legs during physical activity. If the blood supply is severely compromised, it can result in ischemia, leading to tissue damage and, in severe cases, limb loss. Consequently, individuals with PVD often experience symptoms at rest and may develop chronic non-healing wounds due to inadequate perfusion, reinforcing the potential for significant complications associated with this condition. The focus on reduced blood flow and its consequences, such as pain and potential tissue damage, elucidates why this aspect of PVD is critical for understanding its impact on limb health.

7. Which lipid level is typically raised with lipid-lowering therapy?

A. Lipoprotein (a)

B. Triglycerides

C. High density lipoprotein cholesterol

D. Low density lipoprotein cholesterol

High-density lipoprotein cholesterol often referred to as HDL cholesterol, is the lipid level that is typically raised with lipid-lowering therapy. This is particularly significant because HDL is traditionally known as "good" cholesterol. It plays a critical role in removing low-density lipoprotein (LDL) cholesterol from the bloodstream and transporting it to the liver for excretion or reuse. Many lipid-lowering therapies, especially those involving statins, aim not only to lower harmful LDL cholesterol levels but can also have the beneficial side effect of increasing HDL levels. The increase in HDL can contribute to an improved lipid profile and a lower risk of cardiovascular events. In contrast, while triglycerides are commonly targeted and can be decreased with certain therapies, they do not typically increase. Similarly, LDL cholesterol is often the primary target of lipid-lowering medications and is usually reduced rather than raised. Lipoprotein (a) levels are less influenced by standard lipid-lowering treatments. Thus, the increase in HDL cholesterol is a key indicator of successful lipid management in patients undergoing therapy.

SAMPLE

8. What indicates effective understanding of discharge teaching for a patient after coronary artery bypass surgery?

A. Asking whether smoking one cigarette per day is acceptable

B. Clarifying when cardiac rehabilitation will begin

C. Describing plans to relax in a hot tub during recovery

D. Planning to resume driving his or her spouse to work next week

The indication of effective understanding of discharge teaching for a patient after coronary artery bypass surgery is reflected in the clarification of when cardiac rehabilitation will begin. Cardiac rehabilitation is a critical component of recovery, as it helps patients regain their strength, improve cardiovascular health, and reduce the risk of future cardiac events. An informed patient who inquires about the start of cardiac rehabilitation is demonstrating awareness of the importance of a structured recovery plan and the need for guided exercise and education regarding lifestyle changes. Engaging in cardiac rehabilitation signifies recognition of the need for ongoing support after surgery, which plays a vital role in the long-term success of the procedure and overall cardiac health. Understanding when and how to participate in rehabilitation can significantly impact the recovery process and the patient's ability to make safe and effective lifestyle adjustments going forward.

9. For a patient with an abdominal aortic aneurysm undergoing medical treatment, what should be the nurse's priority?

A. Preparing for dialysis to correct renal failure

B. Restricting activities of daily living and low-level exercise

C. Monitoring the aneurysm size monthly with an endocardiogram

D. Maintaining mean blood pressure at a normal level

Maintaining mean blood pressure at a normal level is particularly crucial for a patient with an abdominal aortic aneurysm undergoing medical treatment. Elevated blood pressure can increase the stress and pressure on the aneurysm wall, which can lead to the risk of rupture. By ensuring that the blood pressure remains within a normal range, the nurse plays a critical role in minimizing the risk of complications associated with the aneurysm. Controlling blood pressure helps to manage the overall cardiovascular health of the patient, which is particularly important given that aneurysms are often associated with other vascular conditions. This approach supports the goal of reducing the likelihood of surgical intervention and allows for safe management of the patient's condition through non-invasive treatments. In contrast, other options may not effectively address the immediate health risks associated with an abdominal aortic aneurysm. For instance, preparing for dialysis may apply to patients with renal failure, which is not a direct focus in managing aneurysms. Similarly, while restricting activities is relevant in minimizing stress on the body, it does not directly correlate with managing blood pressure or preventing complications from the aneurysm itself. Monitoring the aneurysm size is important, but it does not take precedence over managing blood pressure, which can have immediate implications for the patient's safety and stability

SAMPLE

10. What benefit can a patient recovering from a myocardial infarction gain from meditation?

A. Increases sympathetic nervous system activity.

B. Decreases sympathetic nervous system activity.

C. Increases the release of catecholamines.

D. Decreases vasodilation.

Meditation can significantly benefit a patient recovering from a myocardial infarction by decreasing sympathetic nervous system activity. This reduction leads to various physiological advantages, especially in the context of heart health. When the sympathetic nervous system is activated, it can lead to increased heart rate, elevated blood pressure, and higher levels of stress hormones like catecholamines. All of these can pose risks for a patient in recovery from a heart attack. Conversely, meditation fosters a state of relaxation and mindfulness, which directly counteracts the stress response associated with sympathetic activation. As the sympathetic activity decreases, the patient may experience improved heart rate variability, lower blood pressure, and a calming effect that can aid in their overall recovery. This shift can also promote better mental health and emotional well-being, which are essential components of cardiac rehabilitation. In summary, the therapeutic practice of meditation intertwines with heart health by mitigating stress responses, supporting more favorable cardiovascular outcomes, and contributing to the overall recovery journey post-myocardial infarction.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cvrn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE