

CARDIOPULMONARY EXAM SKILLS PRACTICE TEST (SAMPLE)

STUDY GUIDE



**SAMPLE STUDY GUIDE
WITH LIMITED QUESTIONS**

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://cardiopulmonaryskills.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Which abnormal BP response describes a BP pattern of a lower than expected increase during exercise, often due to beta-blocker medications?**
 - A. Chronotropic Incompetence
 - B. Hypotensive
 - C. Hypertensive
 - D. Blunted
- 2. Tension pneumothorax commonly presents with which features?**
 - A. Hypotension and tracheal shift
 - B. Hypertension with bradycardia
 - C. Normal vital signs
 - D. Fever and productive cough
- 3. For optimal JVP assessment, at what patient position should you evaluate?**
 - A. 0 degrees (flat)
 - B. 15–20 degrees
 - C. 30–45 degrees
 - D. 60–70 degrees
- 4. Which auscultation area corresponds to the mitral valve?**
 - A. Aortic area (right upper sternal border)
 - B. Pulmonic area (left upper sternal border)
 - C. Erb's point (left sternal border, 3rd/4th intercostal)
 - D. Mitral area (apex at left fifth intercostal space midclavicular line)
- 5. Which breathing pattern uses nearly closed lips during expiration to maintain positive pressure in the bronchioles and prevent premature collapse of weakened airways?**
 - A. Diaphragmatic breathing
 - B. Hyperpnea
 - C. Orthopnea
 - D. Pursed-lip breathing

- 6. What term denotes a heart rate response where the heart rate does not increase proportionally with increasing exercise intensity?**
- A. Blunted
 - B. Chronotropic Incompetence
 - C. Post-exercise Hypotension
 - D. Hypertensive
- 7. Which BP response during exercise testing is most indicative of poor LV function when SBP rises poorly?**
- A. Hypotensive
 - B. Hypertensive
 - C. Blunted
 - D. Post-exercise Hypotension
- 8. Which symptom is most characteristic of a pneumothorax?**
- A. Bilateral crackles
 - B. JVD with hypotension
 - C. Gradual onset chest pain
 - D. Sudden onset chest pain and dyspnea
- 9. Which anatomical landmark is specifically listed as a chest wall motion measurement site along with the sternal angle and xiphoid process?**
- A. Midpoint between xiphoid process and umbilicus
 - B. Umbilicus
 - C. Xiphoid process
 - D. Sternal notch
- 10. Palpation focusing on the lower lobes targets which lung regions?**
- A. Upper lobes
 - B. Lower lobes
 - C. Middle lobes
 - D. Apex

Answers

SAMPLE

1. D
2. A
3. C
4. D
5. D
6. B
7. A
8. D
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. Which abnormal BP response describes a BP pattern of a lower than expected increase during exercise, often due to beta-blocker medications?

- A. Chronotropic Incompetence
- B. Hypotensive
- C. Hypertensive
- D. Blunted**

The main idea here is how blood pressure should respond as you start exercising. Normally, systolic blood pressure rises with increasing work because the heart pumps more blood to meet muscle demands. A blunted BP response means the systolic pressure doesn't rise as much as expected for the level of exercise. This pattern is commonly seen with beta-blocker medications, which blunt the heart's rate and contractility, limiting the rise in cardiac output during activity and thus damping the BP increase. This differs from a chronotropic incompetence, which concerns the heart rate not increasing adequately, not the blood pressure rise itself. It also differs from a hypotensive response, where BP fails to rise or falls, and a hypertensive response, where BP climbs excessively. So the description best fits a blunted BP response.

2. Tension pneumothorax commonly presents with which features?

- A. Hypotension and tracheal shift**
- B. Hypertension with bradycardia
- C. Normal vital signs
- D. Fever and productive cough

The key idea is that tension pneumothorax creates an increasingly pressured pleural space that pushes the mediastinal structures, including the trachea, away from the affected side. This pressure also compresses the great vessels, reducing venous return to the heart and causing a drop in cardiac output, which leads to hypotension (often with tachycardia). The trachea shifts away from the side of the pneumothorax as a classic sign of this mediastinal displacement. Clinically, you typically see rapid breathing and distress, hypoxemia, distended neck veins, and decreased or absent breath sounds on the affected side. This pattern contrasts with fever and productive cough (suggesting infection), normal vital signs (not consistent with the hemodynamic impact), or hypertension with bradycardia (not characteristic of this emergency).

3. For optimal JVP assessment, at what patient position should you evaluate?

- A. 0 degrees (flat)
- B. 15–20 degrees
- C. 30–45 degrees**
- D. 60–70 degrees

Jugular venous pressure is estimated by how high the jugular venous pulse sits above the sternum, which reflects right atrial pressure. The best position to visualize this pulsation clearly and obtain a reliable measurement is semi-recumbent with the head of the bed elevated about 30 to 45 degrees. At this angle, the venous column is visible without being flattened by lying completely flat, and the pulsation remains prominent enough to measure the vertical height from the sternal angle to the top of the pulse. If you're flat, the pulsation can be difficult to distinguish and may be confused with the carotid pulse; if you're too upright, the pulsation can fade and become unreliable. Hence, 30–45 degrees is the optimal position.

4. Which auscultation area corresponds to the mitral valve?

- A. Aortic area (right upper sternal border)
- B. Pulmonic area (left upper sternal border)
- C. Erb's point (left sternal border, 3rd/4th intercostal)
- D. Mitral area (apex at left fifth intercostal space midclavicular line)**

The main idea is where each heart valve's sounds are best heard. The mitral valve sits at the apex of the heart, which is located along the left midclavicular line. That's why auscultation at the apex—often with the patient in the left lateral decubitus position—is where mitral sounds and murmurs are most clearly detected. The aortic valve is best heard at the right upper sternal border, the pulmonic valve at the left upper sternal border, and Erb's point along the left sternal border around the third or fourth intercostal space is a spot where many sounds can be heard together. So, the mitral area corresponds to the apex at the left midclavicular line.

5. Which breathing pattern uses nearly closed lips during expiration to maintain positive pressure in the bronchioles and prevent premature collapse of weakened airways?

- A. Diaphragmatic breathing
- B. Hyperpnea
- C. Orthopnea
- D. Pursed-lip breathing**

Pursed-lip breathing is the pattern that uses nearly closed lips during expiration to create a small amount of back pressure in the airways, helping to keep the bronchioles open as air exits. In obstructive conditions where airways tend to collapse during expiration, this added expiratory pressure slows down and lengthens the exhale, reducing dynamic airway compression and air trapping. The result is easier, more efficient breathing and better gas exchange. This technique is done by inhaling normally through the nose and then exhaling slowly through pursed lips, like blowing out through a straw, so the exhalation is longer than the inhalation. The other patterns don't serve this purpose: diaphragmatic breathing emphasizes abdominal movement and relaxation rather than maintaining airway pressure; hyperpnea involves rapid, deep breathing during exertion; orthopnea refers to difficulty breathing when lying flat, not a technique to keep airways open during expiration.

6. What term denotes a heart rate response where the heart rate does not increase proportionally with increasing exercise intensity?

- A. Blunted
- B. Chronotropic Incompetence**
- C. Post-exercise Hypotension
- D. Hypertensive

During exercise, heart rate should rise in tandem with workload to boost cardiac output and meet the body's increasing oxygen needs. When the heart rate does not increase proportionally as exercise intensity grows, this is chronotropic incompetence. It reflects an impaired ability of the sinus node or autonomic system to elevate rate and can result from aging, cardiovascular disease, or medications such as beta-blockers. It's commonly assessed during graded exercise testing by comparing the peak heart rate to the predicted maximum (often roughly 220 minus age) or by using a chronotropic index. This pattern limits exercise capacity and can signal underlying cardiac dysfunction. The other terms don't describe this specific heart rate response: post-exercise hypotension involves a drop in blood pressure after activity, hypertensive refers to high blood pressure, and "blunted" is a vague descriptor without the precise clinical meaning.

7. Which BP response during exercise testing is most indicative of poor LV function when SBP rises poorly?

- A. Hypotensive**
- B. Hypertensive
- C. Blunted
- D. Post-exercise Hypotension

The key idea is how the heart handles increasing workload during exercise. As exercise intensity rises, a healthy left ventricle can boost cardiac output, and systolic blood pressure should rise accordingly. If the systolic BP does not rise as expected, it signals that the heart's ability to increase stroke volume and maintain perfusion is impaired. A hypotensive response during exercise—where systolic BP actually falls or drops significantly as effort increases—most strongly indicates poor LV function. This happens because the ventricle can't augment output to meet the higher metabolic demand, so blood pressure falls, reflecting limited cardiac reserve and potential significant LV dysfunction or severe ischemia. A blunted rise can occur with various noncardiac factors like deconditioning or medications, and while it's abnormal, it isn't as specific or alarming for LV dysfunction as a clear drop in SBP during exertion. A hypertensive response isn't the problem here, since an excessive BP rise doesn't point to poor LV function. Post-exercise hypotension is a different phenomenon that occurs after the test and doesn't reflect the LV response during peak exercise.

8. Which symptom is most characteristic of a pneumothorax?

- A. Bilateral crackles
- B. JVD with hypotension
- C. Gradual onset chest pain
- D. Sudden onset chest pain and dyspnea**

Pneumothorax typically shows an abrupt change: air entering the pleural space causes the lung to collapse and provokes sudden chest pain with shortness of breath. This combination—sudden, pleuritic chest pain plus dyspnea—best captures the classic presentation, reflecting the rapid loss of lung volume on the affected side. Bilateral crackles are more aligned with fluid in the lungs, such as heart failure, not a collapsed lung. JVD with hypotension can occur in tension pneumothorax, an emergency, but it's not the typical presentation of a non-tension pneumothorax and the symptom that people most commonly notice first is the sudden chest pain with dyspnea. Gradual onset chest pain suggests other processes (like angina or musculoskeletal pain) rather than the abrupt event of a pneumothorax. So the hallmark symptom is the sudden onset chest pain with dyspnea.

9. Which anatomical landmark is specifically listed as a chest wall motion measurement site along with the sternal angle and xiphoid process?

A. Midpoint between xiphoid process and umbilicus

B. Umbilicus

C. Xiphoid process

D. Sternal notch

Chest wall motion assessment uses fixed midline landmarks to quantify how the chest expands during breathing. Along with the sternal angle and the xiphoid process, a commonly used site is the midpoint between the xiphoid process and the umbilicus. This location provides a reliable, easily palpable point on the lower chest wall that reflects overall thoracoabdominal excursion during deep inspiration. Placing measurements at these three levels lets you track expansion from upper to lower chest and detect asymmetry or limited motion that might not be evident at a single site. The midpoint between the xiphoid and the umbilicus is favored over other candidates like the umbilicus itself (which can vary with body habitus) because it stays on the chest wall midline and correlates well with true chest wall movement.

10. Palpation focusing on the lower lobes targets which lung regions?

A. Upper lobes

B. Lower lobes

C. Middle lobes

D. Apex

Palpation focusing on the lower lobes targets the diaphragmatic bases of the lungs—the posterior and basal regions of the chest where the lower lobes extend toward the diaphragm. These areas are best accessed by palpating the lower portion of the chest, especially on the posterior side, where the bases sit closest to the chest wall. The apex sits at the very top and the upper and middle regions occupy higher portions of the chest, so they aren't the focus when you're targeting the lower lobes.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://cardiopulmonaryskills.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE