CARDIOPULMONARY RESUSCITATION (CPR) PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cpr.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. When should you use an AED during a cardiac arrest situation?

A. Only if the patient is conscious

B. As soon as it is available

C. After performing CPR for 10 minutes

D. Only when trained personnel arrive

2. What should you do if you are uncertain about providing CPR?

A. Wait for someone else to arrive

B. Immediately begin CPR regardless of the situation

C. Call for emergency services and seek their guidance

D. Take a few moments to assess the scene

3. In adult CPR, what is the correct sequence of actions?

A. Check responsiveness, call for help, give compressions

B. Call for help, check responsiveness, give rescue breaths

C. Give compressions, check pulse, call for help

D. Give rescue breaths, call for help, check responsiveness

4. What does red color coding indicate in a medical emergency?

A. Open chest CPR

B. DNR

C. Close chest CPR

D. Shock protocols

5. What does the term "agonal breathing" refer to?

A. Normal breathing that suggests the victim is stable

B. Gasping or irregular breathing that may occur in cardiac arrest

C. Rapid and deep breathing during exertion

D. Breathing that indicates the victim is recovering

6. What should you ensure about the victim before using an AED?

A. The victim is sitting upright

B. The victim is lying flat on their back

C. The victim is on an elevated surface

D. The victim has a pulse

SAMPLE

7. What is the correct hand position for CPR on a child?

A. Both hands on the upper chest

B. One hand on the lower half of the sternum

C. Both hands placed over the heart

D. One hand on the abdomen

8. How frequently should intratracheal drugs be administered during CPR?

A. Every 10-15 minutes

B. Every 3-5 minutes

C. Every cycle only once

D. Continuously without breaks

9. What is a key indicator that ET tube placement is successful?

A. Increased heart rate

B. High blood pressure

C. Stable ETCO2 readings

D. Improved level of consciousness

10. How long should you continue CPR?

A. Until emergency personnel arrive, the person starts to show signs of life, or you are too exhausted to continue

B. For exactly 10 minutes

C. Until the person is revived

D. Until a doctor arrives to take over

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. B
5. B
6. B
7. B
8. B
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. When should you use an AED during a cardiac arrest situation?

A. Only if the patient is conscious

B. As soon as it is available

C. After performing CPR for 10 minutes

D. Only when trained personnel arrive

Using an AED during a cardiac arrest situation should occur as soon as it is available. The primary goal in a cardiac arrest scenario is to restore a viable heart rhythm and revive the individual as quickly as possible. An automated external defibrillator (AED) is designed to assess the heart's rhythm and provide a shock if necessary. Immediate use of an AED can significantly improve the chances of survival, particularly within the first few minutes after cardiac arrest occurs. Delaying its use, such as waiting for trained personnel or after performing CPR for several minutes, may diminish the likelihood of a successful outcome. Quick access to and application of the AED, combined with high-quality CPR, maximizes the chances of restoring a normal heart rhythm and sustaining life. In contrast, using an AED on a conscious patient is inappropriate, as the device is specifically designed for situations where the heart is not effectively pumping blood. Hence, the emphasis is on early application of the AED as part of the emergency response protocol in cardiac emergencies.

2. What should you do if you are uncertain about providing CPR?

A. Wait for someone else to arrive

B. Immediately begin CPR regardless of the situation

C. Call for emergency services and seek their guidance

D. Take a few moments to assess the scene

When you are uncertain about providing CPR, it is essential to call for emergency services and seek their guidance. This action is critical as it ensures that you are following the most current medical protocols and getting professional assistance on the way. Emergency service operators are trained to provide instruction in emergency situations, including how to assess the victim and what immediate steps to take. They can help determine whether CPR is necessary, provide guidance on how to proceed, and reassure you while support is en route. This approach combines the urgency of the situation with the necessity of proper assessment and professional input, ensuring that the victim receives appropriate care as soon as possible. Emergency services can also dispatch help while you receive direction, which is a vital part of the rescue process.

3. In adult CPR, what is the correct sequence of actions?

A. Check responsiveness, call for help, give compressions

B. Call for help, check responsiveness, give rescue breaths

C. Give compressions, check pulse, call for help

D. Give rescue breaths, call for help, check responsiveness

The correct sequence of actions in adult CPR begins with checking for responsiveness. This initial step is crucial as it allows you to determine whether the person is conscious or unresponsive and helps guide your subsequent actions. If the person is unresponsive, calling for help is the next critical step to ensure that emergency medical services are alerted as soon as possible. After calling for help, the next action is to begin chest compressions. This prioritization of compressions is based on current CPR guidelines that emphasize the importance of early chest compressions in providing blood flow to vital organs in an unresponsive individual, particularly when a cardiac arrest occurs. This sequence fosters an efficient and effective response that can significantly increase the chance of survival and minimize the potential for irreversible damage until professional help arrives.

SAMPLE

4. What does red color coding indicate in a medical emergency?

A. Open chest CPR

B. DNR

C. Close chest CPR

D. Shock protocols

In a medical emergency, red color coding typically indicates a Do Not Resuscitate (DNR) order. This is an important directive concerning the wishes of a patient regarding resuscitative efforts. When a patient has a DNR in place, it signifies that they have requested not to undergo CPR or other life-saving interventions in the event of cardiac arrest or respiratory failure. Recognizing the color coding for DNR is critical for medical personnel, as it helps ensure compliance with the patient's wishes and guides the approach to their care. In scenarios where a DNR is indicated, medical providers will not initiate resuscitation efforts such as chest compressions or advanced airway management. This helps to honor the patient's autonomy and the decisions they have made regarding their medical treatment. Understanding this color coding is vital for effective communication in emergencies and for providing care that aligns with patient preferences.

5. What does the term "agonal breathing" refer to?

A. Normal breathing that suggests the victim is stable

B. Gasping or irregular breathing that may occur in cardiac arrest

C. Rapid and deep breathing during exertion

D. Breathing that indicates the victim is recovering

The term "agonal breathing" refers to a specific type of breathing pattern characterized by gasping or irregular breaths that can occur in the setting of cardiac arrest. This phenomenon often occurs when the brain is not receiving adequate oxygen due to a lack of effective heart function. Agonal breathing may be mistaken for normal breathing by bystanders, but it is typically shallow, labored, and ineffective in providing adequate oxygenation. Recognizing this type of breathing is crucial, as it indicates that the person is in a life-threatening situation requiring immediate medical intervention, specifically cardiopulmonary resuscitation (CPR) and defibrillation if necessary. Identifying agonal breathing can prompt bystanders to act swiftly to help the victim, potentially saving their life.

6. What should you ensure about the victim before using an AED?

A. The victim is sitting upright

B. The victim is lying flat on their back

C. The victim is on an elevated surface

D. The victim has a pulse

Before using an Automated External Defibrillator (AED), it is essential that the victim is lying flat on their back. This position allows for proper placement of the AED pads on the chest, which is crucial for effective defibrillation. The pads must be applied without interference, and the flat position helps ensure that they are in contact with the skin and that the heart can be effectively assessed and treated. Additionally, positioning the victim flat on their back increases safety during the defibrillation process. If the victim is sitting upright or on an elevated surface, it can complicate pad placement and may lead to ineffective shock delivery. Ensuring the victim has a pulse is not necessary before using the AED, as the device is meant to treat potential cardiac arrest situations, where a pulse is not present.

SAMPLE

7. What is the correct hand position for CPR on a child?

A. Both hands on the upper chest

B. One hand on the lower half of the sternum

C. Both hands placed over the heart

D. One hand on the abdomen

The correct hand position for CPR on a child involves placing one hand on the lower half of the sternum. This position allows for effective chest compressions needed to circulate blood during cardiac arrest. The lower half of the sternum is the ideal location because it is in line with the center of the chest, where compressions can generate the necessary pressure to pump blood effectively, while also reducing the risk of causing injury to the ribs. Using one hand is appropriate for a child, as it provides enough force while being mindful of the child's smaller size compared to an adult. It is essential to ensure that the compressions are deep enough (approximately 1.5 to 2 inches) and at a rate of 100 to 120 compressions per minute for maximum effectiveness. The focus should always be on delivering compressions that are both effective and appropriate for the anatomical differences found in children.

8. How frequently should intratracheal drugs be administered during CPR?

A. Every 10-15 minutes

B. Every 3-5 minutes

C. Every cycle only once

D. Continuously without breaks

Administering intratracheal drugs during CPR is done to enhance the effectiveness of resuscitation efforts, particularly when intravenous access is not possible. The correct frequency is every 3-5 minutes, which aligns with the need for timely drug administration to support cardiac function and overall resuscitation efforts. The rationale for this frequency is tied to the pharmacokinetics of the drugs used; administering them at 3-5 minute intervals allows for timely peaks in drug concentration, promoting better outcomes during the critical moments of cardiac arrest. This interval also corresponds to the cycles of CPR that allow for safety and reassessment of the patient's condition. Inappropriate frequencies, such as continuous administration or too infrequent intervals, could lead to ineffective resuscitation. Continuous administration may not allow for adequate evaluation of the patient's response, while administering drugs every 10-15 minutes might delay action during a critical time when rapid intervention is necessary. The method of drug administration, the clinical setting, and the patient's response to therapy all play crucial roles in effective CPR management.

SAMPLE

9. What is a key indicator that ET tube placement is successful?

A. Increased heart rate

B. High blood pressure

C. Stable ETCO2 readings

D. Improved level of consciousness

A stable ETCO2 (End-Tidal Carbon Dioxide) reading is a key indicator of successful endotracheal (ET) tube placement. When the ET tube is correctly positioned in the trachea, it allows for proper ventilation of the lungs, leading to effective gas exchange. Elevated levels of ETCO2 in the exhaled air indicate that carbon dioxide is being adequately removed from the bloodstream, reflecting good perfusion and ventilation. Monitored ETCO2 levels should show a consistent reading within the expected range, typically between 35 and 45 mmHg, which suggests that the ventilated air is reaching the lungs appropriately. If the ETCO2 readings are significantly lower or absent, it may indicate that the tube is not placed correctly in the trachea, possibly in the esophagus, and that the patient is not effectively ventilated. In contrast, indicators such as increased heart rate could arise from patient stress or other underlying conditions and are not definitively tied to the placement of an ET tube. Similarly, high blood pressure and improved levels of consciousness may reflect a variety of health factors unrelated to respiratory support. Therefore, although they might be observed during treatment, they do not specifically validate the adequacy of ET tube placement as reliably

10. How long should you continue CPR?

A. Until emergency personnel arrive, the person starts to show signs of life, or you are too exhausted to continue

B. For exactly 10 minutes

C. Until the person is revived

D. Until a doctor arrives to take over

Continuing CPR is crucial for maximizing the chances of survival in a cardiac arrest situation. The correct choice highlights three key circumstances under which CPR should be continued: until emergency personnel arrive, until the person shows signs of life, or until you become too exhausted to keep performing the procedure. Emergency personnel are trained to take over advanced cardiac care once they arrive, making it essential to keep performing CPR until they do. Signs of life, such as normal breathing or movement, indicate that the heart may have started to function again, signaling that CPR is no longer needed. It is also important to acknowledge the physical limits of the rescuer; exhaustion can impair the effectiveness of CPR, thereby necessitating a change in approach. The other options suggest fixed time limits or conditions that do not align with the flexible, situational nature of CPR protocols. Resuscitation cannot be curtailed after a set duration without evaluating the rescuer's condition and the patient's status, which emphasizes the need for situational awareness during these critical moments.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cpr.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE