CASUALTY ACTUARIAL SOCIETY MAS-1 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://casualactuarialsocmas1.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Ordinary least squares estimators are inherently unbiased.

A. True

B. False

C. Sometimes

D. Never

2. What is the MVUE of a Poisson distribution?

A. sqrt(x_bar)

B. x_bar

C. 1/x_bar

D. x_bar^2

3. In the given model, E[X] equals E[θ].

A. True

B. False

C. Only if Var(θ)=0

D. Not necessarily

4. In Poisson regression, which statement about the variance-mean relationship is true?

A. Variance is constant

B. Variance equals mean squared

C. Variance equals mean

D. Variance is inverse of mean

5. Ridge regression uses an L2 penalty on coefficients.

A. True

B. False

C. Uses an L1 penalty

D. Uses no penalty

SAMPLE

6. As lambda increases towards infinity, the ridge penalty term has no effect and the estimates become unconstrained.

A. True

B. It depends on the data

C. False

D. Ridge becomes identical to OLS

7. What is the canonical link for the inverse Gaussian distribution?

A. Identity

B. Log

C. Inverse

D. Inverse squared

8. Is the density f(y; θ) = θ y for y > θ a member of the exponential family?

A. Yes, it is a standard member of the exponential family.

B. No, the domain cannot depend on θ and thus it is not in the exponential family.

C. Yes, if θ is treated as a natural parameter.

D. No, only when θ is fixed.

9. Is boosting an example of supervised or unsupervised learning?

A. Unsupervised

B. Reinforcement

C. Semi-supervised

D. Supervised

10. The MVUE of the scale parameter beta in Gamma(shape=alpha, scale=beta) is which expression?

A. alpha / x_bar

B. x_bar / alpha

C. x_bar * alpha

D. x_bar

SAMPLE

Answers

SAMPLE

1. A
2. B
3. D
4. C
5. A
6. C
7. D
8. B
9. D
10. B

SAMPLE

Explanations

SAMPLE

1. Ordinary least squares estimators are inherently unbiased.

A. True

B. False

C. Sometimes

D. Never

The key idea is that OLS is unbiased when the regression errors have zero mean and are uncorrelated with the regressors, i.e., E[ε|X] = 0. Under this exogeneity condition, the OLS coefficients satisfy E[β_hat|X] = β, because β_hat = (X'X)^{-1}X'Y = β + (X'X)^{-1}X'ε, and E[X'ε|X] = X'E[ε|X] = 0. Taking expectations gives E[β_hat] = β. So, on average across repeated samples, the estimator hits the true parameter values. If that zero-mean exogeneity fails—due to endogeneity, omitted variables, measurement error, or simultaneity—OLS can be biased. So the statement is true within the standard assumptions, but not guaranteed in all situations.

2. What is the MVUE of a Poisson distribution?

A. sqrt(x_bar)

B. x_bar

C. 1/x_bar

D. x_bar^2

For Poisson data, the natural statistic to use for estimating the rate λ is the sum of observations, which is complete and sufficient for λ. The sample mean $\bar{X} = (X1 + ... + Xn)/n$ is a function of that complete sufficient statistic, and it is unbiased since $E[\bar{X}] = E[(1/n)\sum Xi] = \lambda$. By the Lehmann–Scheffé theorem, the unbiased estimator that is a function of the complete sufficient statistic has the smallest possible variance among all unbiased estimators of λ. Since $\bar{X}$ is unbiased for λ and is derived from the complete sufficient statistic, it is the MVUE for λ. That's why the best estimator here is the sample mean. The other transformations of $\bar{X}$ (like $\sqrt{\bar{X}}$, $1/\bar{X}$, or $\bar{X}^2$) do not have E[·] equal to λ in general, so they are biased and cannot be the MVUE.

3. In the given model, E[X] equals E[θ].

A. True

B. False

C. Only if Var(θ)=0

D. Not necessarily

The ability to link E[X] and E[θ] hinges on how X behaves given θ. By the law of total expectation, E[X] = E[E[X|θ]]. For E[X] to equal E[θ], the conditional mean of X given θ must be θ itself (that is, E[X|θ] = θ for almost every θ). When that holds, E[X] = E[θ]. But in a general model, X could depend on θ in other ways. If E[X|θ] is some function of θ other than θ, or if there's a bias in the conditional mean, then E[X] becomes E[E[X|θ]] = E[g(θ)] with g(θ) ≠ θ, which need not equal E[θ]. For example, if X|θ = 2θ, then E[X] = 2 E[θ], not E[θ]. If X|θ = θ^2, then E[X] = E[θ^2], again not generally equal to E[θ]. Thus the statement is not guaranteed; it's not necessarily true unless the model specifies E[X|θ] = θ (or an equivalent condition that preserves the expectation).

SAMPLE

4. In Poisson regression, which statement about the variance-mean relationship is true?

A. Variance is constant

B. Variance equals mean squared

C. Variance equals mean

D. Variance is inverse of mean

In Poisson regression, the outcome follows a Poisson distribution with mean mu, and a defining property is that the variance equals the mean: Var(Y) = mu. Since mu is modeled as a function of predictors (often with a log link, mu = exp(Xβ)), the variance naturally grows with the mean rather than staying constant. So the variance-mean relationship highlighted here is Var(Y) = E(Y) = mu. This differs from a constant-variance assumption and from a variance that scales with the square of the mean. If overdispersion occurs (Var(Y) > mu), you might use a quasi-Poisson or negative binomial approach to allow greater variance.

5. Ridge regression uses an L2 penalty on coefficients.

A. True

B. False

C. Uses an L1 penalty

D. Uses no penalty

*Ridge regression adds a penalty that is proportional to the sum of squares of the coefficients. This L2 penalty discourages large coefficient values, shrinking them toward zero to stabilize estimates when predictors are correlated. The objective becomes the usual residual sum of squares plus lambda times the sum of squared coefficients: RSS + lambda * sum(beta_j^2). As lambda grows, all coefficients are pulled toward zero, but they are typically not driven exactly to zero, keeping the model dense. This behavior helps reduce variance from multicollinearity, unlike doing nothing (no penalty) or using an L1 penalty (which can drive some coefficients to zero, producing a sparse model).*

6. As lambda increases towards infinity, the ridge penalty term has no effect and the estimates become unconstrained.

A. True

B. It depends on the data

C. False

D. Ridge becomes identical to OLS

Ridge regression adds a penalty on coefficient magnitudes to the ordinary least squares objective. The objective is the residual sum of squares plus lambda times the sum of squared coefficients. As lambda grows, the penalty term becomes more influential, pulling the coefficients toward zero to reduce complexity and variance. If lambda goes to infinity, the penalty dominates the objective, and the estimates collapse toward zero (ignoring the intercept). They become more constrained, not unconstrained. Since OLS corresponds to lambda = 0, ridge does not become identical to OLS at large lambda. So the statement is false.

SAMPLE

7. What is the canonical link for the inverse Gaussian distribution?

A. Identity

B. Log

C. Inverse

D. Inverse squared

The concept being tested is how the link function connects the mean to the natural parameter in the exponential-family form of a distribution. For the inverse Gaussian distribution with mean μ, writing its density in exponential-family form shows the natural parameter depends on μ in a way that makes the appropriate link be g(μ) = μ^(-2). In other words, the canonical link is the inverse-squared link because it aligns the linear predictor with the distribution's natural parameter. The other options do not match that exponential-family relationship for the inverse Gaussian, so they're not the canonical choice.

8. Is the density f(y; θ) = θ y for y > θ a member of the exponential family?

A. Yes, it is a standard member of the exponential family.

B. No, the domain cannot depend on θ and thus it is not in the exponential family.

C. Yes, if θ is treated as a natural parameter.

D. No, only when θ is fixed.

The key idea is that a density in the standard exponential family must have a fixed support that does not depend on the parameter. Here the density is defined only for y > θ, so the range of y changes with θ. That dependence on θ in the domain violates the fixed-support requirement of the standard exponential family. Moreover, evaluating the normalization shows a fundamental problem: ∫ from θ to ∞ of θ y dy diverges (it becomes infinite for θ > 0, and isn't a valid nonnegative density for θ ≤ 0). So this cannot be a proper density for any θ, which further confirms it cannot be a member of the standard exponential family. If the domain were fixed independently of θ, it could be possible to express a density in exponential-family form, but as given, the domain dependence rules it out.

9. Is boosting an example of supervised or unsupervised learning?

A. Unsupervised

B. Reinforcement

C. Semi-supervised

D. Supervised

Boosting is a supervised learning approach. It trains a sequence of simple models, or weak learners, on labeled data, using the mistakes of earlier models to focus the next one on harder examples. Because each training example comes with a known target label, the learning signal comes from comparing predictions to those true labels. This is different from unsupervised learning, which finds structure without labels, and from reinforcement learning, which learns through interaction with an environment and rewards. While boosting can be extended to semi-supervised settings in some cases, the standard, core boosting methods rely on labeled data, making supervised the best description.

SAMPLE

10. The MVUE of the scale parameter beta in Gamma(shape=alpha, scale=beta) is which expression?

A. alpha / x_bar

B. x_bar / alpha

C. x_bar * alpha

D. x_bar

*Estimating the scale parameter beta from a Gamma distribution with known shape alpha relies on how the mean relates to beta. The sample mean X has expectation E[X] = alpha * beta, so dividing by alpha gives an unbiased estimator for beta: X /alpha has E[X /alpha] = beta. This estimator can be written as S/(n alpha), where S = sum X_i and n is the sample size; S follows Gamma(n alpha, beta), and S is a complete sufficient statistic for beta. Since the estimator is a function of S, Lehmann–Scheffé guarantees it is the MVUE. Therefore, the best estimator for beta is X divided by alpha.*

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://casualactuarialsocmas1.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE