CENTRAL NERVOUS SYSTEM PHARMACOLOGY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cennervoussyspharmacology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What are common side effects of methylphenidate?

A. Fatigue and weight gain

B. Agitation and insomnia

C. Diarrhea and abdominal pain

D. Confusion and tremors

2. In addition to ADHD, Atomoxetine is sometimes investigated for its efficacy in treating?

A. Sleep Disorders

B. Obesity

C. Impulse Control Disorders

D. Social Anxiety Disorder

3. Which treatment is specifically used for alcohol overdose?

A. Fomepizole

B. Disulfiram

C. Naltrexone

D. Acetylcysteine

4. Which of the following is a common side effect of Atomoxetine?

A. Insomnia

B. Euphoria

C. Weight gain

D. Increased appetite

5. How do stimulant medications function in the treatment of narcolepsy?

A. By inducing sleep

B. By reducing anxiety

C. By promoting wakefulness

D. By enhancing mood

6. Which drug is a common competitive nicotinic antagonist?

A. Propofol

B. Midazolam

C. Atracurium

D. Fentanyl

SAMPLE

7. What is a primary characteristic of ketamine in anesthesia?

A. It does not provide analgesia

B. It induces dissociative anesthesia

C. It is only used for pediatric patients

D. It is only an intravenous agent

8. What condition may occur with sudden discontinuation of major tranquilizers?

A. Serotonin syndrome

B. Neuroleptic malignant syndrome (NMS)

C. Withdrawal syndrome

D. Extrapyramidal symptoms

9. What is a common side effect of antipsychotic medication in older adults?

A. Increased risk of falls and fractures due to sedation

B. Improved cognitive function

C. Decreased appetite

D. Reduction in mobility

10. Which type of antipsychotic medication is less likely to cause extrapyramidal symptoms?

A. Typical antipsychotics

B. Atypical antipsychotics

C. Older antipsychotics

D. High-potency antipsychotics

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. A
5. C
6. C
7. B
8. B
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What are common side effects of methylphenidate?

A. Fatigue and weight gain

B. Agitation and insomnia

C. Diarrhea and abdominal pain

D. Confusion and tremors

Methylphenidate, a common stimulant medication used primarily to treat Attention Deficit Hyperactivity Disorder (ADHD) and narcolepsy, often leads to side effects related to its stimulant nature. Agitation and insomnia are particularly notable among these side effects. Agitation may occur due to the increased neurotransmitter activity, primarily dopamine and norepinephrine, that methylphenidate promotes. This heightened neurotransmitter action can lead to increased arousal and anxiety levels, which may manifest as restlessness or agitation. Insomnia is another frequently reported side effect. The stimulating effect of methylphenidate can interfere with the normal sleep-wake cycle, making it more difficult for individuals to fall asleep or stay asleep. This is particularly common when the medication is taken later in the day, as the stimulating effects can linger. In contrast, choices involving fatigue and weight gain, diarrhea and abdominal pain, or confusion and tremors do not commonly reflect the expected side effects of methylphenidate. Instead, these side effects might be more aligned with other medications or conditions rather than the stimulant properties of methylphenidate. Understanding the common side effects of methylphenidate is crucial for proper patient management and education regarding medication use.

2. In addition to ADHD, Atomoxetine is sometimes investigated for its efficacy in treating?

A. Sleep Disorders

B. Obesity

C. Impulse Control Disorders

D. Social Anxiety Disorder

Atomoxetine is primarily known as a selective norepinephrine reuptake inhibitor used for treating attention-deficit/hyperactivity disorder (ADHD). Its mechanism of action involves increasing norepinephrine levels in the brain, which can help improve attention and reduce impulsivity and hyperactivity. This pharmacological profile makes it a candidate for investigating treatments for disorders characterized by impulsivity and issues with self-control, such as impulse control disorders. Impulse control disorders include conditions such as intermittent explosive disorder and kleptomania, where individuals struggle with resisting the temptation to engage in harmful or socially unacceptable behaviors. Atomoxetine's ability to modulate norepinephrine levels may help in managing symptoms associated with these disorders, potentially allowing individuals to have better control over their impulses. While Atomoxetine has been explored for other potential uses, it is especially relevant in the context of impulse control due to its targeted effects on cognitive processes associated with attention and impulse.

SAMPLE

3. Which treatment is specifically used for alcohol overdose?

A. Fomepizole

B. Disulfiram

C. Naltrexone

D. Acetylcysteine

Fomepizole is specifically used for alcohol overdose because it acts as an inhibitor of the enzyme alcohol dehydrogenase. This enzyme is responsible for the metabolism of alcohol to acetaldehyde, which is the toxic metabolite associated with alcohol poisoning. By blocking this enzyme, fomepizole effectively slows down the metabolism of ethanol, thereby reducing the accumulation of toxic metabolites and giving the patient more time to be treated and recover. It is particularly used in cases of methanol or ethylene glycol poisoning as well, where similar metabolic pathways are involved. Other treatments listed, such as disulfiram, naltrexone, and acetylcysteine, serve different purposes in clinical practice. Disulfiram is used for alcohol dependence, as it creates an unpleasant reaction when alcohol is consumed, thus discouraging drinking. Naltrexone is also used for alcohol use disorder to reduce cravings and the rewarding effects of alcohol. Acetylcysteine is an antidote for acetaminophen overdose, helping to replenish glutathione and protect the liver. Each of these medications has a distinct role and is not aimed specifically at managing alcohol overdose.

4. Which of the following is a common side effect of Atomoxetine?

A. Insomnia

B. Euphoria

C. Weight gain

D. Increased appetite

Atomoxetine is a selective norepinephrine reuptake inhibitor commonly used in the treatment of attention-deficit hyperactivity disorder (ADHD). One of the well-documented side effects of Atomoxetine is insomnia. This occurs due to the medication's stimulation of norepinephrine pathways, which can interfere with sleep patterns. While euphoria, weight gain, and increased appetite can occur with some medications, these are not typical side effects associated with Atomoxetine. In fact, Atomoxetine is generally more associated with decreased appetite and potential weight loss, rather than weight gain or increased appetite. Understanding these pharmacological effects is crucial for recognizing how Atomoxetine may influence mood and physical health, particularly in pediatric populations where monitoring for insomnia is essential.

5. How do stimulant medications function in the treatment of narcolepsy?

A. By inducing sleep

B. By reducing anxiety

C. By promoting wakefulness

D. By enhancing mood

Stimulant medications function in the treatment of narcolepsy primarily by promoting wakefulness. Narcolepsy is a sleep disorder characterized by excessive daytime sleepiness, and the goal of stimulant therapy is to counteract this symptom. These medications work by increasing the levels of neurotransmitters such as dopamine and norepinephrine in the brain, which enhances alertness and reduces the tendency to fall asleep during the day. This increase in wakefulness can help individuals with narcolepsy maintain a more normal level of functioning throughout their daily activities. While other aspects like mood and anxiety can be indirectly influenced by wakefulness and alertness, the primary mechanism of stimulant medications in the context of narcolepsy treatment is to promote wakefulness. Inducing sleep or enhancing mood is not the primary function of these agents for this specific condition, as their core purpose is to help maintain a state of alertness.

SAMPLE

6. Which drug is a common competitive nicotinic antagonist?

A. Propofol

B. Midazolam

C. Atracurium

D. Fentanyl

Atracurium is a common competitive nicotinic antagonist that is primarily used as a neuromuscular blocker during anesthesia. By binding to nicotinic acetylcholine receptors at the neuromuscular junction, atracurium prevents acetylcholine from interacting with these receptors. This action leads to muscle relaxation and paralysis, which is particularly useful during surgical procedures to facilitate intubation and provide optimal operating conditions without muscle movement. In contrast, propofol and midazolam are both central nervous system depressants but act through different mechanisms. Propofol is primarily a GABA receptor agonist that induces sedation and anesthesia. Midazolam also enhances GABAergic activity, providing sedation, anxiolysis, and amnesia. Fentanyl, on the other hand, is an opioid analgesic that primarily acts on mu-opioid receptors to provide pain relief rather than functioning as a neuromuscular blocker. Thus, the distinct mechanism of action of atracurium as a competitive antagonist at nicotinic receptors sets it apart from the other drugs listed, confirming its role as a common competitive nicotinic antagonist.

7. What is a primary characteristic of ketamine in anesthesia?

A. It does not provide analgesia

B. It induces dissociative anesthesia

C. It is only used for pediatric patients

D. It is only an intravenous agent

Ketamine is primarily characterized by its ability to induce dissociative anesthesia, which is a unique state of anesthesia where the patient feels detached from their environment and body. This dissociation allows for surgical procedures to be performed without the patient experiencing pain or awareness, making it highly effective in various settings, including emergency and surgical procedures. One of the key aspects of dissociative anesthesia is that it creates a trance-like state while preserving certain reflexes and cardiovascular stability, distinguishing it from other forms of anesthesia. This characteristic makes ketamine invaluable, especially in situations where traditional anesthetics might be unsuitable. While options regarding analgesic effects and limitations to specific patient populations may hold some truths, they do not encompass the primary unique action of ketamine in anesthesia, which is its capacity to induce dissociative states. Additionally, though ketamine is often administered intravenously, it can also be used via other routes such as intramuscularly or intranasally, thereby broadening its applications beyond just intravenous use.

SAMPLE

8. What condition may occur with sudden discontinuation of major tranquilizers?

A. Serotonin syndrome

B. Neuroleptic malignant syndrome (NMS)

C. Withdrawal syndrome

D. Extrapyramidal symptoms

Neuroleptic malignant syndrome (NMS) is a severe reaction that can occur with the withdrawal of major tranquilizers, primarily antipsychotic medications. This life-threatening condition is characterized by symptoms such as hyperthermia, muscle rigidity, autonomic dysregulation, and altered mental status. When major tranquilizers are suddenly discontinued, the abrupt change in medication levels can lead to a rebound effect, causing the body to react in an exaggerated manner, which embodies the characteristics of NMS. In the context of antipsychotic treatment, NMS can manifest when withdrawal happens too rapidly or when there is an interplay of medications that destabilizes the dopaminergic system. This syndrome illustrates the importance of careful management and gradual tapering of antipsychotic medications to prevent serious complications like NMS. The other conditions, while relevant in the context of CNS pharmacology, do not specifically relate to the sudden discontinuation of major tranquilizers in the same manner.

9. What is a common side effect of antipsychotic medication in older adults?

A. Increased risk of falls and fractures due to sedation

B. Improved cognitive function

C. Decreased appetite

D. Reduction in mobility

The side effect of increased risk of falls and fractures due to sedation is particularly relevant in older adults taking antipsychotic medications. Antipsychotics, especially atypical ones, can cause sedation, dizziness, and orthostatic hypotension, which is a drop in blood pressure upon standing. These effects can impair balance and coordination, significantly elevating the risk of falls. In older individuals, these dangers are compounded since they may already have conditions that affect mobility or balance, and their bodies may not tolerate medication side effects as well as younger adults. The potential for sedation to lead to falls makes this a crucial consideration in prescribing and managing antipsychotic treatment in the elderly population. The other options provided do not align with common side effects observed with antipsychotic medications. Improved cognitive function is not a known effect of these drugs; in fact, they can sometimes have negative impacts on cognitive abilities. Decreased appetite may occur in some patients but is not as universally acknowledged or impactful as the risks associated with sedation. Reduction in mobility could be a result of side effects, but it is frequently tied more directly to sedation and its effects on falls, making the risk of falls and fractures the more prominent concern in this context.

SAMPLE

10. Which type of antipsychotic medication is less likely to cause extrapyramidal symptoms?

A. Typical antipsychotics

B. Atypical antipsychotics

C. Older antipsychotics

D. High-potency antipsychotics

Atypical antipsychotics are considered less likely to cause extrapyramidal symptoms (EPS) compared to typical antipsychotics. The reason for this difference lies in the pharmacological profiles of these drug classes. Typical antipsychotics primarily block dopamine D2 receptors in the brain, which is effective for managing psychotic symptoms but also leads to a higher incidence of EPS due to the disruption of normal dopaminergic activity in areas of the brain that control motor function. These symptoms can include tremors, rigidity, and bradykinesia. In contrast, atypical antipsychotics not only block D2 receptors but also affect other neurotransmitter systems, particularly serotonin receptors (5-HT2A). This broader mechanism of action appears to mitigate the risk of EPS, making atypical antipsychotics generally more tolerable in terms of movement disorders. This unique action helps to balance the dopaminergic pathways in a way that reduces the likelihood of movement-related side effects. Therefore, the characteristic differences in receptor activity and side effect profiles between typical and atypical antipsychotics clarify why atypicals are preferred in many cases, particularly in individuals who are at higher risk or already experiencing EPS.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cennervoussyspharmacology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE