CERTIFIED INDUSTRIAL HYGIENIST PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://industrialhygeinist.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which condition is often associated with phosphorous compounds?

A. Adult respiratory distress syndrome (ARDS)

B. Chronic lung disease

C. Cancer and dermatitis

D. Neurological disorders

2. What is the half-life of carboxyhemoglobin in carbon monoxide poisoning?

A. 2 hours

B. 4 hours

C. 6 hours

D. 8 hours

3. Which of the following bacteria is associated with "rabbit fever"?

A. Tuberculosis

B. Lyme disease

C. Tularemia

D. Plague

4. Which of the following symptoms is associated with carbon monoxide exposure?

A. Visual disturbances

B. Headaches

C. Euphoria

D. Impaired balance

5. Which of the following is a known irritation caused by sulfur dioxide exposure?

A. Skin and mucous membrane irritation

B. Hearing loss

C. Nervous system disorders

D. Cardiovascular disease

SAMPLE

6. What does Boyle's Law express mathematically?

A. P1T1 = P2T2

B. P1V1 = P2V2

C. P1R1 = P2R2

D. P1V2 = P2V1

7. Which range of wavelengths is associated with corneal damage from optical radiation?

A. 600-800 nm

B. 200-400 nm

C. 400-700 nm

D. Below 400 nm

8. What health effect is particularly associated with exposure to microwave radiation?

A. Enhanced vision

B. Increased heart rate

C. Decreased metabolic rate

D. Improved digestion

9. Which of the following diseases does vinyl chloride NOT cause?

A. Raynaud's syndrome

B. Lung cancer

C. Diabetes

D. Angiosarcoma of liver

10. What is the maximum body temperature to prevent impairment from heat stress?

A. Below 37 C or 98.6 F

B. Greater than 40 C or 104 F

C. Within + 1 C of 37 C or 98.6 F

D. No lower than 36 C or 96.8 F

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. A
6. B
7. D
8. B
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. Which condition is often associated with phosphorous compounds?

A. Adult respiratory distress syndrome (ARDS)

B. Chronic lung disease

C. Cancer and dermatitis

D. Neurological disorders

Phosphorus compounds, particularly those that are organophosphates, have been associated with a range of health issues primarily due to their effects on the respiratory system. Inhalation exposure to certain phosphorus compounds can cause inflammation and damage to lung tissues, leading to conditions such as adult respiratory distress syndrome (ARDS). ARDS is characterized by rapid onset of inflammation in the lungs, leading to severe respiratory failure, which can be exacerbated by exposure to harmful chemicals. In the context of the other options, while chronic lung disease can also result from various environmental exposures, it tends to be more closely associated with pollutants and long-term injury rather than acute exposure to phosphorus compounds. Cancer and dermatitis are also health concerns, but they are typically linked with prolonged exposure or different classes of toxicants, rather than being specifically noted with phosphorus. Neurological disorders are primarily associated with different neurotoxic agents, and while some phosphorus compounds can affect the nervous system, they are not as commonly directly linked to acute respiratory health issues as ARDS is. Therefore, the association between phosphorus compounds and the development of ARDS underlines the significance of monitoring exposure to these compounds in occupational settings, where the risk of respiratory distress can be critical.

2. What is the half-life of carboxyhemoglobin in carbon monoxide poisoning?

A. 2 hours

B. 4 hours

C. 6 hours

D. 8 hours

The half-life of carboxyhemoglobin, which is the compound formed when carbon monoxide binds to hemoglobin, is approximately four hours in cases of carbon monoxide poisoning when the person is breathing room air. Understanding this half-life is crucial in clinical situations as it impacts the duration of treatment and the assessment of the patient's recovery. If a patient is exposed to carbon monoxide, the body starts to eliminate carboxyhemoglobin over time. The four-hour half-life means that after this period, half of the carboxyhemoglobin present would be eliminated, assuming normal conditions without interventions like hyperbaric oxygen therapy, which can significantly reduce the half-life. This timeframe informs medical professionals about the expected recovery rate and guides the need for further therapeutic interventions, including providing oxygen or considering hyperbaric therapy for severe cases. Understanding this aspect of carbon monoxide poisoning is vital for effective treatment and care management.

SAMPLE

3. Which of the following bacteria is associated with "rabbit fever"?

A. Tuberculosis

B. Lyme disease

C. Tularemia

D. Plague

The bacteria associated with "rabbit fever" is indeed tularemia, which is caused by the bacterium Francisella tularensis. This disease is often linked to contact with infected rabbits or rodents and can also be transmitted through bites from infected insects or ingestion of contaminated food and water. Tularemia is characterized by a variety of symptoms, including fever, swollen lymph nodes, skin ulcers, and respiratory issues, depending on the method of transmission and the location of the infection within the body. The nickname "rabbit fever" stems from its historical association with rabbits as a common reservoir for the bacterium, which can cause outbreaks in rural areas where hunting and handling of these animals occur. Understanding the specific characteristics and transmission pathways of tularemia is essential for comprehensive industrial hygiene practices, particularly in environments where there's potential exposure to wildlife. This reinforces the importance of appropriate safety measures and monitoring in controlling infections related to zoonotic diseases. The other diseases listed, such as tuberculosis, Lyme disease, and plague, are caused by different pathogens and are associated with unique transmission routes and symptoms, hence differentiating them from tularemia.

4. Which of the following symptoms is associated with carbon monoxide exposure?

A. Visual disturbances

B. Headaches

C. Euphoria

D. Impaired balance

Headaches are commonly associated with carbon monoxide exposure due to its binding affinity to hemoglobin in the blood. When carbon monoxide is inhaled, it competes with oxygen for binding sites on hemoglobin, leading to decreased oxygen transport to tissues. This reduction in oxygen delivery can result in various symptoms, with headaches being one of the most prevalent. As the brain is particularly sensitive to changes in oxygen levels, the vascular changes and irritation in the brain caused by hypoxia can lead to the development of headaches. Other symptoms of carbon monoxide exposure can include dizziness, confusion, and even loss of consciousness at higher levels, but headaches are often one of the first signs reported during exposure. Understanding the presentation of symptoms such as headaches helps in assessing exposure risk and implementing appropriate safety measures.

SAMPLE

5. Which of the following is a known irritation caused by sulfur dioxide exposure?

A. Skin and mucous membrane irritation

B. Hearing loss

C. Nervous system disorders

D. Cardiovascular disease

The known irritation caused by sulfur dioxide exposure is related to its ability to affect the respiratory system and the mucous membranes. When sulfur dioxide is inhaled, it can lead to irritation of the eyes, nose, throat, and lungs. This results from its corrosive nature; as a gas, sulfur dioxide can easily react with moisture in the respiratory tract and mucous membranes to form sulfurous acid, causing inflammation, pain, and irritation. The symptoms commonly associated with exposure include coughing, mucus secretion, difficulty breathing, and a burning sensation in the throat or chest. Individuals with pre-existing respiratory conditions, such as asthma, are particularly vulnerable to these irritating effects, which can exacerbate their symptoms. Other options, while they may be serious health issues, are not directly linked to sulfur dioxide exposure. Hearing loss and nervous system disorders, for example, can be caused by other environmental toxins or conditions, but sulfur dioxide is primarily recognized for its irritant effects rather than these more systemic effects. Cardiovascular disease can be influenced by a multitude of factors, including air pollution, but sulfur dioxide itself is not a primary cause associated with that condition.

6. What does Boyle's Law express mathematically?

A. P1T1 = P2T2

B. P1V1 = P2V2

C. P1R1 = P2R2

D. P1V2 = P2V1

Boyle's Law, which is foundational in the study of gases in the field of physics and chemistry, states that, at constant temperature, the pressure of a given quantity of gas is inversely proportional to its volume. This relationship is beautifully expressed mathematically as the product of pressure and volume being a constant value for a given amount of gas. When you see the equation represented as P1V1 = P2V2, it indicates that if the pressure (P) increases, the volume (V) will decrease, and vice versa, provided that the temperature remains unchanged. This direct relationship allows for the calculation of how changes in pressure can impact the volume of a gas, which is a crucial concept in fields like industrial hygiene, where understanding the behavior of gases is important for safety and health assessments. The other equations listed do not correctly capture the essence of Boyle's Law. They might represent other physical laws or principles but are not applicable to the relationship governed by Boyle's Law concerning pressure and volume at a constant temperature.

SAMPLE

7. Which range of wavelengths is associated with corneal damage from optical radiation?

A. 600-800 nm

B. 200-400 nm

C. 400-700 nm

D. Below 400 nm

Corneal damage from optical radiation primarily occurs in the range of wavelengths below 400 nm, which includes ultraviolet (UV) radiation. This UV radiation can have harmful effects on various parts of the eye, especially the cornea, leading to conditions such as photokeratitis, cataracts, and other ocular injuries. The UV spectrum is typically categorized into three regions: UVA (320-400 nm), UVB (280-320 nm), and UVC (100-280 nm). It is primarily the UVB and UVC wavelengths that are concentrated in causing damage due to their higher energy levels and capacity to penetrate biological tissues. The intensity of radiation in this range is often sufficient to cause cellular damage, leading to inflammation and contributing to long-term eye health issues. In contrast, the other ranges of wavelengths presented do not pose the same level of risk for corneal damage. For instance, the ranges from 600-800 nm and 400-700 nm mainly fall into the visible light spectrum and are much less likely to cause direct damage to the cornea. While they can still affect vision and lead to discomfort, they do not carry the same risk of physical injury as the wavelengths below 400 nm. Therefore, identifying that wavelengths below 400 nm

8. What health effect is particularly associated with exposure to microwave radiation?

A. Enhanced vision

B. Increased heart rate

C. Decreased metabolic rate

D. Improved digestion

Exposure to microwave radiation is primarily associated with increased heart rate due to the way microwaves interact with biological tissues. Microwaves can generate heat within the body, which can lead to physiological responses, including variations in heart rate as the body reacts to the warmth. When microwave energy is absorbed, especially at certain power densities, it can lead to thermal effects that reflect the body's response to excess heat, thereby causing an increase in heart rate as the body attempts to manage the elevated temperatures and maintain homeostasis. Understanding this relationship is crucial for assessing potential health risks related to microwave exposure, particularly in occupational settings where exposure levels might be higher. The other options do not correlate with the known effects of microwave radiation exposure. For instance, enhanced vision, decreased metabolic rate, and improved digestion are not established health outcomes related to microwave radiation.

SAMPLE

9. Which of the following diseases does vinyl chloride NOT cause?

A. Raynaud's syndrome

B. Lung cancer

C. Diabetes

D. Angiosarcoma of liver

Vinyl chloride is a well-documented occupational and environmental hazard primarily known for its association with certain serious health conditions. Among those conditions are lung cancer and angiosarcoma of the liver, which have been directly linked to prolonged exposure to this chemical. Similarly, Raynaud's syndrome, which involves episodes of reduced blood flow to the fingers and toes, has also been observed in workers exposed to vinyl chloride, though the mechanism is less clearly defined compared to the carcinogenic effects. In contrast, diabetes is not associated with vinyl chloride exposure. While various chemicals and environmental factors can influence the risk of developing diabetes, vinyl chloride is not recognized as a risk factor for this condition. This distinction is important in the realm of industrial hygiene and health risk assessment, as it underlines the specific toxicological effects of substances and helps in crafting safety regulations and health guidelines for workers potentially exposed to harmful agents. Understanding these associations aids practitioners in identifying risk factors and implementing appropriate preventive measures.

10. What is the maximum body temperature to prevent impairment from heat stress?

A. Below 37 C or 98.6 F

B. Greater than 40 C or 104 F

C. Within + 1 C of 37 C or 98.6 F

D. No lower than 36 C or 96.8 F

The correct response emphasizes the importance of maintaining body temperature within a specific range to prevent heat stress impairment. Typically, a core body temperature exceeding 38-39 C (approximately 100.4-102.2 F) can lead to physiological changes that impair function, performance, and potentially threaten health. Maintaining a body temperature within 1 C of the average temperature of 37 C (or 98.6 F) helps ensure that the body's thermoregulatory mechanisms are functioning effectively, preventing conditions such as heat exhaustion or heat stroke. This range is particularly relevant in occupational settings where workers may be exposed to high temperatures, emphasizing the need for strict monitoring and control measures to ensure safety. Establishing guidelines around this temperature helps in risk assessment and management strategies aimed at protecting individuals from the hazards of heat stress, which can impair cognitive and physical performance. Other options fail to provide a solid protective threshold or reference point for prevention of heat-related illnesses, either suggesting temperatures that are too low or too high, which would not effectively mitigate the risks associated with heat stress.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://industrialhygeinist.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE