CERTIFIED NEPHROLOGY NURSE CERTIFICATION (CNN) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cnn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Name the two main types of dialysis used for renal replacement therapy.

A. Hemodialysis and peritoneal dialysis

B. Cycler dialysis and nocturnal dialysis

C. Continuous renal replacement therapy and intermittent dialysis

D. Standard dialysis and continuous dialysis

2. Name one common symptom of nephritic syndrome.

A. Proteinuria

B. Hematuria

C. Prauritus

D. Edema

3. What is the recommended protein intake per day for adults with chronic kidney disease?

A. 1.2g/kg/day

B. 0.6g/kg/day

C. 0.8g/kg/day

D. 1.0g/kg/day

4. Which process describes the movement of particles from high concentration to low concentration?

A. Osmosis

B. Diffusion

C. Ultrafiltration

D. Contraction

5. In diabetic nephropathy, what is an early sign of kidney damage?

A. Hematuria

B. Proteinuria

C. Microalbuminuria

D. Oliguria

SAMPLE

6. What is the primary function of the vasa recta in the kidney?

A. Filtration of blood

B. Concentration of urine

C. Regulation of blood pressure

D. Excretion of waste products

7. What symptom characterizes azotemia?

A. Increased urination

B. Decreased urination

C. High blood urea nitrogen levels

D. Fluid retention

8. What dietary modification might be recommended for a patient with advanced CKD?

A. Increase sodium intake

B. High protein diet

C. Limit phosphorus intake

D. Decrease fluid restriction

9. What is the most common cause of end-stage renal disease (ESRD) in the United States?

A. Hypertension

B. Polycystic kidney disease

C. Diabetes mellitus

D. Chronic glomerulonephritis

10. What effect does NSAIDs have on kidney function?

A. Enhances renal perfusion

B. Promotes urinary retention

C. Can cause interstitial nephritis

D. Reduces glomerular filtration rate

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. C
6. B
7. C
8. C
9. C
10. D

SAMPLE

Explanations

SAMPLE

1. Name the two main types of dialysis used for renal replacement therapy.

A. Hemodialysis and peritoneal dialysis

B. Cycler dialysis and nocturnal dialysis

C. Continuous renal replacement therapy and intermittent dialysis

D. Standard dialysis and continuous dialysis

The two main types of dialysis used for renal replacement therapy are hemodialysis and peritoneal dialysis. Hemodialysis involves the use of a machine and a dialyzer, which acts as an artificial kidney to filter waste products and excess fluids from the blood. This process typically occurs in a dialysis center but can also be done at home with appropriate training. Peritoneal dialysis, on the other hand, uses the lining of the abdominal cavity (the peritoneum) as the filter. A special dialysis solution is introduced into the abdomen, where it absorbs waste products and excess fluids. After a designated time, the solution is drained away, taking the wastes with it. This method can be done at home and offers patients more flexibility and independence. Other options mentioned, such as cycler dialysis and nocturnal dialysis, refer to specific methods or schedules for delivering dialysis and do not represent the fundamental types of dialysis. Continuous renal replacement therapy is a specialized form generally applied in acute settings rather than the routine maintenance types of hemodialysis and peritoneal dialysis. Thus, the distinction clearly rests with the two primary types: hemodialysis and peritoneal dialysis.

2. Name one common symptom of nephritic syndrome.

A. Proteinuria

B. Hematuria

C. Prauritus

D. Edema

Hematuria, which is the presence of blood in the urine, is indeed a common symptom of nephritic syndrome. This condition is characterized by inflammation of the glomeruli in the kidneys, which can lead to leakage of red blood cells into the urine. This symptom, along with others such as hypertension and renal impairment, highlights the underlying glomerular damage and inflammation that occurs in nephritic syndrome. In contrast, while proteinuria, or excess protein in the urine, can be present, it is more prominent in nephrotic syndrome. Pruritus, which refers to itching, is not a defining characteristic of nephritic syndrome and rather relates to other conditions or complications like uremia. Edema, although it can occur, is less common specifically in nephritic syndrome compared to nephrotic syndrome, where significant fluid retention is typically more evident. Thus, hematuria distinctly indicates the type of kidney dysfunction characteristic of nephritic syndrome.

SAMPLE

3. What is the recommended protein intake per day for adults with chronic kidney disease?

A. 1.2g/kg/day

B. 0.6g/kg/day

C. 0.8g/kg/day

D. 1.0g/kg/day

For adults with chronic kidney disease (CKD), the recommended protein intake is generally around 0.8 grams per kilogram of body weight per day. This amount is considered appropriate for maintaining nutritional status without exacerbating kidney damage. In CKD, managing protein intake is crucial because excessive protein can lead to a buildup of nitrogenous waste products in the blood, thereby increasing the workload on the kidneys. The 0.8g/kg/day recommendation strikes a balance, allowing individuals to receive enough protein to meet their nutritional needs while minimizing the risk of further harm to renal function. This is particularly important in the context of CKD, where maintaining muscle mass and overall health is critical for patient outcomes. As kidney function declines, protein needs may further decrease, which is why healthcare providers often closely monitor dietary recommendations tailored to the individual's overall health status, stage of kidney disease, and other coexisting medical conditions.

4. Which process describes the movement of particles from high concentration to low concentration?

A. Osmosis

B. Diffusion

C. Ultrafiltration

D. Contraction

The movement of particles from an area of high concentration to an area of low concentration is known as diffusion. This process occurs naturally as particles tend to move towards a state of equilibrium, where concentrations are balanced across a space. It is a fundamental principle in biology and chemistry, relevant to many physiological processes. For example, in the kidneys, diffusion plays a crucial role in the regulation of substances such as electrolytes and urea, allowing for the proper balance of these materials in the blood and urine. This process is passive and does not require energy, which distinguishes it from active transport mechanisms that move substances against their concentration gradient. In contrast, osmosis specifically refers to the movement of water across a semipermeable membrane, while ultrafiltration involves the movement of water and small solutes through a filter, influenced by pressure. Contraction does not relate to particle movement in the context of concentration gradients. Understanding these distinctions is essential in nephrology and overall physiological processes.

SAMPLE

5. In diabetic nephropathy, what is an early sign of kidney damage?

A. Hematuria

B. Proteinuria

C. Microalbuminuria

D. Oliguria

Microalbuminuria is considered an early sign of kidney damage in diabetic nephropathy because it reflects the initial leakage of albumin into the urine. In a healthy individual, the kidneys filter blood and prevent significant amounts of protein, such as albumin, from passing into the urine. However, in diabetic nephropathy, even small amounts of albumin can start to appear as the kidney's filtration mechanisms begin to fail. Detecting microalbuminuria, which is defined as the presence of 30 to 300 mg of albumin per gram of creatinine in the urine, allows for early identification of kidney impairment. This is crucial because early intervention can help mitigate further kidney damage through improved glycemic control and the use of medications such as ACE inhibitors. In contrast, hematuria typically suggests a different type of kidney pathology unrelated to early diabetic changes. Oliguria, which refers to significantly reduced urine output, indicates more advanced kidney dysfunction rather than early damage. Recognizing microalbuminuria allows healthcare providers to take preventive measures before the disease progresses to more severe forms of kidney damage, such as established proteinuria or chronic renal failure.

6. What is the primary function of the vasa recta in the kidney?

A. Filtration of blood

B. Concentration of urine

C. Regulation of blood pressure

D. Excretion of waste products

The vasa recta play a crucial role in the concentration of urine. These are specialized capillaries that run alongside the nephron loops (specifically the loops of Henle) in the renal medulla. Their primary function is to maintain the osmotic gradient established by the loop of Henle, which is essential for the kidney's ability to concentrate urine. As blood flows through the vasa recta, it participates in the countercurrent exchange mechanism. This allows for the reabsorption of water and solutes as it descends and ascends through the medulla. The osmotic gradient created by the longer loop of Henle and maintained by the vasa recta enables the kidney to concentrate urine efficiently, which is vital for water conservation and proper fluid balance in the body. In summary, the primary function of the vasa recta in the kidney is to facilitate the concentration of urine through their involvement in the countercurrent exchange system, supporting the kidney's ability to concentrate waste and conserve water.

SAMPLE

7. What symptom characterizes azotemia?

A. Increased urination

B. Decreased urination

C. High blood urea nitrogen levels

D. Fluid retention

Azotemia is primarily characterized by high levels of urea nitrogen in the blood, specifically indicated by elevated blood urea nitrogen (BUN) levels. This condition often arises from kidney dysfunction, where the kidneys are unable to effectively filter waste products from the blood, leading to an accumulation of nitrogenous waste. An increase in the concentration of these substances in the bloodstream is a hallmark of impaired renal function and is central to diagnosing azotemia. While fluid retention and changes in urination may be associated with kidney issues, they are not defining symptoms of azotemia itself. Increased or decreased urination can vary based on the specific causes of kidney impairment, such as acute or chronic kidney disease, but they do not specifically characterize azotemia. Thus, recognizing elevated blood urea nitrogen as the central feature clarifies the understanding of azotemia in clinical practice.

8. What dietary modification might be recommended for a patient with advanced CKD?

A. Increase sodium intake

B. High protein diet

C. Limit phosphorus intake

D. Decrease fluid restriction

In the management of advanced chronic kidney disease (CKD), it is crucial to address dietary modifications to help manage the disease's progression and mitigate complications. Limiting phosphorus intake is a recommended dietary intervention for patients with advanced CKD due to the kidneys' decreased ability to excrete phosphorus. High levels of phosphorus can lead to complications such as bone disease and cardiovascular issues, as the body can begin to extract calcium from the bones to maintain normal serum levels. Phosphorus is prevalent in many foods, particularly in dairy products, processed foods, and some meats. Patients are often advised to avoid foods high in phosphorus and limit their overall phosphorus intake to prevent the buildup of phosphorus in the blood, which can lead to hyperphosphatemia. This dietary change can help preserve bone health and reduce the risk of cardiovascular problems associated with high phosphorus levels. The other dietary options mentioned would not be appropriate for a patient with advanced CKD. For instance, increasing sodium intake can lead to fluid retention and hypertension, both of which are already concerns in CKD. A high protein diet can increase the burden on the kidneys and accelerate their decline, while decreasing fluid restriction could exacerbate fluid overload, especially in patients approaching end-stage renal disease. Thus, limiting phosphorus intake stands

SAMPLE

9. What is the most common cause of end-stage renal disease (ESRD) in the United States?

A. Hypertension

B. Polycystic kidney disease

C. Diabetes mellitus

D. Chronic glomerulonephritis

Diabetes mellitus is the most common cause of end-stage renal disease (ESRD) in the United States, accounting for approximately 44% of all cases. This condition leads to diabetic nephropathy, characterized by damage to the kidneys due to prolonged high blood glucose levels. The resultant microvascular changes, including glomerular hypertension and hyperfiltration, eventually lead to progressive kidney damage and loss of function. The association between diabetes and renal impairment is significant, especially as the prevalence of diabetes has increased markedly over the years. Proper management of blood sugar levels can slow the progression of kidney disease; however, once significant damage has occurred, the risk of evolving to ESRD is greatly heightened. This underscores the importance of careful monitoring and management of patients with diabetes to prevent complications that can lead to renal failure. Understanding the implications of diabetes on kidney function is crucial for nephrology nurses and highlights the need for education and intervention strategies in this population.

10. What effect does NSAIDs have on kidney function?

A. Enhances renal perfusion

B. Promotes urinary retention

C. Can cause interstitial nephritis

D. Reduces glomerular filtration rate

The effect of NSAIDs, or non-steroidal anti-inflammatory drugs, on kidney function primarily involves their impact on renal blood flow and filtration processes. The correct response highlights that NSAIDs can cause interstitial nephritis, a condition characterized by inflammation of the kidney interstitium. NSAIDs inhibit the production of prostaglandins, which are compounds that play a crucial role in maintaining renal blood flow, particularly in situations where there is reduced perfusion, such as dehydration or congestive heart failure. When prostaglandin production is inhibited, this can lead to a decrease in renal perfusion and may result in acute interstitial nephritis, which is marked by inflammation and damage to the kidney tissue. This mechanism illustrates how NSAIDs, despite their efficacy in pain relief and inflammation, can adversely affect kidney function. Understanding this side effect is particularly important for patients who might be at risk, such as those with pre-existing renal conditions or other comorbidities. In contrast, other options do not accurately reflect the effects of NSAIDs. For example, they do not enhance renal perfusion or promote urinary retention directly. While NSAIDs may influence glomerular filtration rate, they typically do so by reducing it during renal impairment rather than as a

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cnn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE