CERTIFIED OSTOMY CARE NURSE (COCN) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cocn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What skin issues are common around stomas?

A. Skin cancer

B. Skin irritation or dermatitis

C. Acne

D. Eczema

2. Which medication is known to decrease peristaltic activity and reduce stool odor, but may cause general malaise with long-term use?

A. Loperamide

B. Bismuth subgallate

C. Diphenoxylate

D. Omeprazole

3. True or False: The clinician should secure the nephrostomy tube at a perpendicular angle to the body.

A. True

B. False

C. It depends on the patient's anatomy

D. None of the above

4. Which medications should ostomy patients be cautious with?

A. Antidepressants and pain relievers

B. Over-the-counter antihistamines

C. Laxatives and certain antibiotics

D. Vitamins and minerals

5. What is a stoma cap used for?

A. To block stoma output completely

B. To cover the stoma when no output is expected, aiding in discretion

C. To protect the stoma from water exposure

D. To assist in the healing of the stoma

SAMPLE

6. Following the second post-op visit after IPAA surgery, which instruction should be given to decrease stooling frequency?

A. Encourage the patient to delay defecation.

B. Encourage the patient to respond immediately to the urge to defecate.

C. Encourage Kegels prior to each defecation.

D. Encourage increased solid food volume at mealtime.

7. Which surgical procedure is a 75-year-old male likely to undergo after being admitted with an acute sigmoid obstruction due to perforated diverticulitis?

A. End-loop colostomy

B. End ileostomy

C. Colostomy with Hartmann's Pouch

D. Loop diverting ileostomy

8. After a radical cystectomy, what is one of the best ways for a patient with an orthotopic neobladder to prevent incontinence?

A. Performing pelvic floor muscle exercises 2-3 times daily

B. Clean intermittent catheterization

C. Irrigating the neobladder to reduce mucous

D. Drinking enough fluids to dilute the urine

9. Which of the following is considered an early complication following ostomy surgery?

A. Parastomal hernia

B. Stomal retraction

C. Stomal prolapse

D. All of the above

10. A patient with abdominal pain and cramps is marked for an ileostomy. The colonoscopy shows lesions that balloon out through the wall of the intestine. What condition does this indicate?

A. Ulcerative colitis

B. Diverticulitis

C. Crohn's disease

D. Diverticulosis

SAMPLE

Answers

SAMPLE

1. B
2. B
3. A
4. C
5. B
6. D
7. C
8. A
9. D
10. C

SAMPLE

Explanations

SAMPLE

1. What skin issues are common around stomas?

A. Skin cancer

B. Skin irritation or dermatitis

C. Acne

D. Eczema

Skin irritation or dermatitis is a common issue around stomas, primarily due to the presence of appliances and the interactions between the adhesive materials, skin, and bodily secretions. The stoma site is often subjected to moisture and trauma from the ostomy appliance changing procedures, which can lead to skin breakdown and irritation. This can manifest as redness, itching, and discomfort around the stoma. Factors contributing to skin irritation include improper fitting of the ostomy appliance, allergic reactions to adhesives, and exposure to digestive contents, which can cause chemical burns and inflammation. Managing skin integrity is crucial for individuals with stomas, as compromised skin can lead to additional complications. While options like skin cancer, acne, and eczema can occur in the general population, they are not specifically associated with the presence of a stoma in the same way that skin irritation and dermatitis are. Therefore, understanding the commonalities of skin issues related to ostomy care is essential for providing effective and compassionate patient care.

2. Which medication is known to decrease peristaltic activity and reduce stool odor, but may cause general malaise with long-term use?

A. Loperamide

B. Bismuth subgallate

C. Diphenoxylate

D. Omeprazole

Bismuth subgallate is known for its ability to decrease peristaltic activity in the intestines, which can lead to a reduction in stool frequency and urgency. Additionally, it possesses properties that help to reduce stool odor, making it beneficial in managing the symptoms associated with certain gastrointestinal issues, particularly in patients with ostomies. However, long-term use of bismuth subgallate may lead to general malaise. This refers to a feeling of discomfort, uneasiness, or lack of well-being that can arise from various sources, including the potential for buildup of the medication in the body, which may eventually cause some systemic effects. In the context of the other options, while loperamide and diphenoxylate also decrease peristalsis, they do not have the same properties regarding stool odor. Omeprazole, a proton pump inhibitor, is primarily used for reducing stomach acid and is not directly related to decreasing peristaltic activity or stool odor. Thus, bismuth subgallate emerges as the medication that aligns with all aspects of the question.

SAMPLE

3. True or False: The clinician should secure the nephrostomy tube at a perpendicular angle to the body.

A. True

B. False

C. It depends on the patient's anatomy

D. None of the above

The recommendation to secure the nephrostomy tube at a perpendicular angle to the body is based on ensuring proper drainage and minimizing the risk of complications. When a nephrostomy tube is positioned perpendicular to the body, it aligns more efficiently with the kidney's drainage system, allowing urine to flow unobstructed from the renal pelvis through the tube and into the external collection device. Additionally, ensuring that the tube is perpendicular can help reduce tension on the catheter, thus lowering the risk of dislodgement or kinking, which could lead to urinary retention, infection, or other complications. This positioning also aids in patient comfort and can facilitate easier care and maintenance of the tube. While anatomical variations might influence specific placement in individual cases, the general guideline is to maintain the tube at a perpendicular angle to maximize function and safety.

4. Which medications should ostomy patients be cautious with?

A. Antidepressants and pain relievers

B. Over-the-counter antihistamines

C. Laxatives and certain antibiotics

D. Vitamins and minerals

Ostomy patients should be cautious with laxatives and certain antibiotics due to their potential impact on bowel function and the overall health of the stoma. Laxatives can lead to increased bowel activity, which may result in diarrhea or dehydration, particularly in patients with a shortened bowel or altered bowel habits due to their ostomy. This can compromise the patient's ability to manage their output effectively, potentially leading to skin irritation or peristomal complications. Certain antibiotics, on the other hand, can upset the balance of gut flora, potentially resulting in antibiotic-associated diarrhea, which can be particularly problematic for those with an ostomy. This disruption could lead to changes in output consistency, increase the risk of dehydration, or even influence the stoma area health due to output that may irritate the surrounding skin. Given these considerations, the need for caution with these types of medications is particularly relevant for ostomy patients. It's essential for healthcare providers to educate patients on the appropriate use of these medications and monitor their effects closely.

SAMPLE

5. What is a stoma cap used for?

A. To block stoma output completely

B. To cover the stoma when no output is expected, aiding in discretion

C. To protect the stoma from water exposure

D. To assist in the healing of the stoma

A stoma cap is designed primarily to cover the stoma when no output is expected, providing a discreet solution for individuals with stomas. This product is particularly advantageous for patients who desire more confidence and comfort in social situations, allowing them to engage in daily activities without worrying about the visibility or sound associated with ostomy output. Stoma caps are typically small, lightweight, and designed to fit snugly against the skin around the stoma, ensuring that there is minimal disruption to the surrounding area and that the patient feels secure. This discreet coverage is essential for many ostomy patients, particularly when they are active or when they want to avoid drawing attention to their stoma. The other options, while relating to stoma care, do not accurately reflect the purpose of a stoma cap. For instance, completely blocking stoma output would not align with the function of a cap, as ostomates still need to manage output appropriately. Similarly, while stoma caps provide some level of protection from water, their primary purpose is discretion rather than protection from water exposure or assisting in healing, which would typically involve different types of products and care protocols.

6. Following the second post-op visit after IPAA surgery, which instruction should be given to decrease stooling frequency?

A. Encourage the patient to delay defecation.

B. Encourage the patient to respond immediately to the urge to defecate.

C. Encourage Kegels prior to each defecation.

D. Encourage increased solid food volume at mealtime.

Increasing solid food volume at mealtime is an effective instruction for decreasing stooling frequency after ileal pouch-anal anastomosis (IPAA) surgery. Solid foods generally take longer to digest, which can help form more solid stools and reduce the frequency of bowel movements. After this type of surgery, individuals often experience changes in bowel habits, and dietary adjustments play a crucial role in managing these symptoms. Patients who consume a diet richer in solid foods may experience improved stool consistency and form, leading to fewer urges to defecate. It's important to note that the choice of foods also matters; patients should focus on foods that are higher in soluble fiber, which can help absorb excess fluid and contribute to firmer stools. In contrast, encouraging the patient to delay defecation may lead to discomfort and potential rectal complications or increased urgency in the long run. Urging immediate response to the urge to defecate can lead to inconsistent bowel habits. Promoting Kegel exercises prior to each defecation is more about pelvic floor muscle tone and does not directly address stooling frequency or consistency and might not be as relevant to short-term stool management post-surgery. Therefore, increasing the volume of solid foods at meals is the most appropriate and effective instruction

SAMPLE

7. Which surgical procedure is a 75-year-old male likely to undergo after being admitted with an acute sigmoid obstruction due to perforated diverticulitis?

A. End-loop colostomy

B. End ileostomy

C. Colostomy with Hartmann's Pouch

D. Loop diverting ileostomy

The most appropriate surgical procedure for a 75-year-old male who is admitted with an acute sigmoid obstruction due to perforated diverticulitis is a colostomy with Hartmann's pouch. This approach is often utilized in cases of diverticulitis, especially when there is perforation and potential contamination of the peritoneal cavity. Hartmann's procedure involves the resection of the affected segment of the colon and the creation of a colostomy. The distal portion of the bowel is usually left in place but not connected to the rectum; instead, it is closed off. This is beneficial because it allows for immediate diversion of bowel contents, effectively managing the acute condition and preventing complications related to fecal contamination. Additionally, Hartmann's pouch provides a potential for future restoration of bowel continuity once the inflammation has resolved, which is a significant consideration in the management of diverticulitis. This option aligns with the need for urgent intervention in the case of perforation while also considering the possibility of reestablishing normal bowel function later. The other options may not be as appropriate given the specific circumstances of acute sigmoid obstruction and perforation.

8. After a radical cystectomy, what is one of the best ways for a patient with an orthotopic neobladder to prevent incontinence?

A. Performing pelvic floor muscle exercises 2-3 times daily

B. Clean intermittent catheterization

C. Irrigating the neobladder to reduce mucous

D. Drinking enough fluids to dilute the urine

Performing pelvic floor muscle exercises, often referred to as Kegel exercises, is an effective way for patients with an orthotopic neobladder to prevent incontinence after a radical cystectomy. These exercises strengthen the pelvic floor muscles, which play a crucial role in supporting the bladder and maintaining urinary control. By consistently engaging in pelvic floor muscle exercises 2-3 times daily, patients enhance their ability to contract these muscles when a strong urge to urinate arises, thereby reducing the risk of involuntary leakage. Other methods such as clean intermittent catheterization can help manage urinary retention issues but do not focus on preventing incontinence. Irrigating the neobladder primarily serves to clear mucus and potential blockages, which is important for maintaining the neobladder's function but does not directly address incontinence management. Drinking sufficient fluids is vital for overall health and can help dilute urine, but it does not specifically target the prevention of incontinence like pelvic floor muscle exercises do.

SAMPLE

9. Which of the following is considered an early complication following ostomy surgery?

A. Parastomal hernia

B. Stomal retraction

C. Stomal prolapse

D. All of the above

All of the options listed—parastomal hernia, stomal retraction, and stomal prolapse—can indeed be considered early complications following ostomy surgery. Parastomal hernia refers to the protrusion of intestine or abdominal contents through the muscle wall near the stoma site. This can occur soon after the surgery due to increased intra-abdominal pressure or weakness in the abdominal wall tissue. Stomal retraction involves the stoma being pulled back into the abdominal wall leading to a lower profile of the stoma. This condition can arise shortly after surgery if the stoma was not positioned or sutured adequately in relation to the skin. Stomal prolapse occurs when the stoma extends beyond its normal size and protrudes excessively. This can happen within weeks after the surgery, especially if the stoma is excessively mobilized or if there is increased abdominal pressure. Together, these conditions represent complications that can manifest relatively early in the post-operative period for patients who have undergone ostomy procedures, emphasizing the need for proper post-operative care and monitoring.

10. A patient with abdominal pain and cramps is marked for an ileostomy. The colonoscopy shows lesions that balloon out through the wall of the intestine. What condition does this indicate?

A. Ulcerative colitis

B. Diverticulitis

C. Crohn's disease

D. Diverticulosis

The presence of lesions that balloon out through the wall of the intestine is indicative of Crohn's disease. This condition is characterized by transmural inflammation, which can lead to the formation of fistulas, strictures, and abscesses, as well as the appearance of "cobblestoning" due to ulceration and scarring. The lesions described often result in complications that may necessitate surgical interventions, such as an ileostomy. While ulcerative colitis can also cause abdominal pain and has associated lesions, it typically affects only the mucosal layer of the colon and does not feature the same transmural involvement or significantly ballooning lesions. Diverticulitis, on the other hand, involves inflammation of diverticula, which are pouches in the colon, rather than lesions that cause wall protrusion throughout the intestine. Diverticulosis refers to the presence of diverticula without inflammation, which does not match the symptomatology or endoscopic findings described. This makes Crohn's disease the most fitting diagnosis in this context.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cocn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE