CERTIFIED PEDIATRIC EMERGENCY NURSE (CPEN) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cpen.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What constitutes the "A" in the ABCDE approach to pediatric emergencies?

A. Airway management

B. Access to veins

C. Assessment of pain

D. Administration of fluids

2. What are common causes of fever in pediatric patients?

A. Viral infections, dehydration, and diabetes

B. Infections, inflammatory diseases, and CNS disorders

C. Overheating, teething, and anxiety

D. Allergies, infections, and environmental factors

3. What does the presence of "moon facies" indicate in a pediatric assessment?

A. Possible dehydration.

B. Possible malnutrition.

C. Possible steroid use or Cushing's syndrome.

D. Possible genetic disorder.

4. What is the recommended compression to ventilation ratio for pediatric CPR?

A. 15:2 for single rescuer

B. 30:2 for single rescuer and 15:2 for two rescuers

C. 20:2 for single rescuer

D. 30:2 for two rescuers

5. What is a key indicator of airway obstruction in a pediatric patient?

A. Bradycardia

B. Tachypnea

C. Hypoxia

D. Stridor

SAMPLE

6. What should caregivers avoid to reduce the risk of SIDS?

A. Smokey environments

B. Exposing infants to sick individuals

C. Both A and B

D. Keeping infants in a separate room while they sleep

7. How do you calculate a child's blood pressure using the formula for hypotension?

A. MAP = (SBP + DBP) / 2

B. MAP = (2 x DBP + SBP) / 3

C. MAP = (DBP + 2 x SBP) / 3

D. MAP = DBP + SBP

8. During a neurological assessment, what does a level of alertness indicate?

A. Degree of pain tolerance

B. Response to external environments

C. Ability to follow commands

D. Stability of vital signs

9. How should you respond if a child presents with unstable cardiac rhythm?

A. Monitor continuously

B. Use electrical methods like defibrillation

C. Administer oxygen immediately

D. Consult a cardiologist

10. Which condition requires immediate administration of glucose?

A. Dehydration

B. Hypoglycemia

C. Sepsis

D. Cardiac arrest

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. D
6. C
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What constitutes the "A" in the ABCDE approach to pediatric emergencies?

A. Airway management

B. Access to veins

C. Assessment of pain

D. Administration of fluids

In the ABCDE approach to pediatric emergencies, the "A" stands for Airway management. This is a fundamental aspect of emergency care, as ensuring a patent airway is critical for effective ventilation and oxygenation. Without a clear airway, a child cannot breathe adequately, which can lead to hypoxia and potential respiratory failure or cardiac arrest. In pediatric patients, airway considerations are particularly important due to their anatomical differences and the fact that they can become obstructed more easily compared to adults. Proper assessment and management of the airway, which may involve positioning, suctioning, or even advanced airway intervention, are crucial to stabilize the patient and provide essential life support. Understanding airway management as the first step in the ABCDE approach emphasizes the priority of maintaining airway patency before moving on to breathing and circulation, ensuring a structured and efficient response to emergencies in children.

2. What are common causes of fever in pediatric patients?

A. Viral infections, dehydration, and diabetes

B. Infections, inflammatory diseases, and CNS disorders

C. Overheating, teething, and anxiety

D. Allergies, infections, and environmental factors

The correct choice highlights common causes of fever in pediatric patients, emphasizing the role of infections, inflammatory diseases, and central nervous system (CNS) disorders. Infections, such as viral and bacterial illnesses, are the most prevalent reasons for fever in children. When the body experiences an infection, the immune response generates pyrogens that increase the thermostat set point in the hypothalamus, resulting in fever as a means to combat the pathogen. Inflammatory diseases, including conditions such as juvenile idiopathic arthritis or systemic lupus erythematosus, can also lead to fever due to the underlying inflammatory processes influencing the body's temperature regulation. Additionally, central nervous system disorders, such as meningitis or encephalitis, can cause fever by directly affecting the brain's temperature control center or by causing inflammation, leading to an elevated temperature. This response effectively encompasses a range of significant medical conditions that can present with fever in children, making it a comprehensive choice for the causes of fever in pediatric patients.

SAMPLE

3. What does the presence of "moon facies" indicate in a pediatric assessment?

A. Possible dehydration.

B. Possible malnutrition.

C. Possible steroid use or Cushing's syndrome.

D. Possible genetic disorder.

The presence of "moon facies" is indicative of possible steroid use or Cushing's syndrome in a pediatric assessment. Moon facies refers to the characteristic rounding of the face due to fat redistribution, which is commonly associated with the use of corticosteroids or the endogenous overproduction of cortisol seen in Cushing's disease. In children, prolonged exposure to corticosteroids, whether through medication or other medical conditions affecting hormone production, can lead to a variety of physical changes, including the development of moon facies. These changes occur due to the accumulation of fat deposits in the facial region, leading to the typical full appearance of the cheeks and rounded shape of the face. Recognizing moon facies is crucial for pediatric healthcare providers, as it may prompt further investigation into the child's underlying medical condition and medication history, particularly regarding endocrine disorders or chronic steroid therapy. Understanding this manifestation helps in identifying and addressing health concerns effectively.

4. What is the recommended compression to ventilation ratio for pediatric CPR?

A. 15:2 for single rescuer

B. 30:2 for single rescuer and 15:2 for two rescuers

C. 20:2 for single rescuer

D. 30:2 for two rescuers

The recommended compression to ventilation ratio for pediatric cardiopulmonary resuscitation (CPR) is significant because it helps ensure effective circulation and oxygenation in a child who has experienced cardiac arrest. For a single rescuer, the compression to ventilation ratio is 30:2, which allows for optimal airflow and blood flow during the resuscitation process. This ratio is designed to maximize both CPR quality and the chances of return of spontaneous circulation (ROSC). In the case of two rescuers performing CPR on a child, the compression to ventilation ratio is adjusted to 15:2. This change acknowledges the increased efficiency and ability for ventilation that a second rescuer provides. The two rescuer team can switch off duties, allowing for effective compressions while maintaining airway control and ventilation, thus enhancing the overall effectiveness of resuscitation efforts for the pediatric patient. Applying these recommended ratios is crucial for pediatric resuscitation, as children have different airway and cardiovascular physiology compared to adults, necessitating tailored approaches that reflect these differences. By following the established guidelines, the chances of survival and recovery for pediatric patients are improved.

SAMPLE

5. What is a key indicator of airway obstruction in a pediatric patient?

A. Bradycardia

B. Tachypnea

C. Hypoxia

D. Stridor

Stridor is a key indicator of airway obstruction in a pediatric patient because it often signifies a partially obstructed airway, particularly in the upper airway. Stridor is characterized by a high-pitched, wheezing sound primarily heard during inhalation and is often caused by conditions such as croup, foreign body aspiration, or laryngeal edema. This sound indicates turbulence in airflow due to a narrowed airway, which is a critical sign that requires immediate assessment and potentially intervention. The presence of stridor suggests that the patient's airway may be compromised, making it an essential clinical finding in emergency settings. Immediate evaluation and management of the underlying cause of stridor are necessary to ensure adequate airflow and oxygenation. Other symptoms such as bradycardia, tachypnea, and hypoxia may also relate to respiratory distress or other complications in pediatric patients but are less specific indicators of airway obstruction compared to stridor. In an emergency context, identifying stridor can help clinicians prioritize treatment and initiate appropriate steps to relieve the obstruction swiftly.

6. What should caregivers avoid to reduce the risk of SIDS?

A. Smokey environments

B. Exposing infants to sick individuals

C. Both A and B

D. Keeping infants in a separate room while they sleep

To effectively reduce the risk of Sudden Infant Death Syndrome (SIDS), caregivers should actively avoid exposing infants to smokey environments and sick individuals. Smoking, particularly in close proximity to infants, has been consistently linked to a higher incidence of SIDS. The harmful substances present in cigarette smoke can affect an infant's respiratory health and potentially lead to unsafe sleeping conditions. Similarly, exposing infants to individuals who are ill may present a risk through various infectious diseases, which can compromise an infant's delicate immune system. Keeping an infant's environment as healthy and safe as possible plays a crucial role in their overall well-being. Choosing to keep infants in a separate room while they sleep, however, does not directly correlate with a reduction in the risk of SIDS. In fact, current recommendations often suggest that infants sleep in the same room as their caregivers, as this can enhance safety monitoring and reduce the likelihood of SIDS by ensuring that the infant is in a closer, more supervised environment. Therefore, avoiding both smokey environments and exposure to sick individuals is essential for caregivers aiming to minimize the risk of SIDS.

SAMPLE

7. How do you calculate a child's blood pressure using the formula for hypotension?

A. MAP = (SBP + DBP) / 2

B. MAP = (2 x DBP + SBP) / 3

C. MAP = (DBP + 2 x SBP) / 3

D. MAP = DBP + SBP

The correct formula to calculate the Mean Arterial Pressure (MAP) for predicting hypotension in children is represented by the equation where MAP is calculated as (2 x DBP + SBP) / 3. This formula is particularly relevant in pediatrics because it better reflects the average arterial pressure during one cardiac cycle given the longer diastolic phase in children relative to systolic pressure. The reasoning behind this method involves understanding how to represent blood pressure across the cardiac cycle. In children, diastolic blood pressure contributes more significantly to the overall blood pressure profile than systolic, thus the MAP formula gives double weight to the diastolic blood pressure. This method of calculation ensures a more accurate assessment of perfusion pressure in pediatric patients, which is essential when evaluating for hypotension. Utilizing the correct MAP formula is crucial during clinical assessments as it helps determine if the child is experiencing inadequate blood flow to their organs, which can lead to serious complications. In assessment protocols, it's important for healthcare professionals to recognize this formula as an effective tool, especially when differentiating between normal and concerning blood pressures in the pediatric population.

8. During a neurological assessment, what does a level of alertness indicate?

A. Degree of pain tolerance

B. Response to external environments

C. Ability to follow commands

D. Stability of vital signs

The level of alertness is a crucial component of a neurological assessment as it reflects a patient's response to their external environment. This means that a higher level of alertness indicates that the patient is aware of their surroundings and can interact with stimuli such as sounds, sights, and even gentle physical touches. When assessing alertness, healthcare providers look for the patient's ability to be responsive and engaged. This can involve checking whether a patient can maintain focus, respond to questions, or react to sensory inputs, which is vital for determining neurological function. Understanding a patient's level of alertness can help identify potential neurological deficits or changes in mental status, which may warrant further evaluation or intervention. The other options, while related to different aspects of neurological assessment and overall health, do not directly define what level of alertness signifies. For instance, pain tolerance and the ability to follow commands can be influenced by alertness but measure different attributes of neurological and physical function, while stability of vital signs relates to physiological parameters rather than cognitive engagement with the environment.

SAMPLE

9. How should you respond if a child presents with unstable cardiac rhythm?

A. Monitor continuously

B. Use electrical methods like defibrillation

C. Administer oxygen immediately

D. Consult a cardiologist

When a child presents with an unstable cardiac rhythm, the appropriate response is to use electrical methods such as defibrillation. Unstable cardiac rhythms can lead to serious complications, including decreased cardiac output and loss of consciousness, making it imperative to restore a normal heart rhythm promptly. Defibrillation is crucial in cases of life-threatening arrhythmias, such as ventricular fibrillation or pulseless ventricular tachycardia, as it delivers a therapeutic dose of electrical energy to the heart, allowing it to reset and ideally revert to a normal rhythm. This immediate intervention can be life-saving and is a critical step in managing arrythmias effectively. While continuous monitoring, administering oxygen, and consulting a cardiologist may play a role in the overall care for a child with cardiac issues, they do not address the immediate need to correct an unstable rhythm. Monitoring is essential but does not contribute to resolving the instability, oxygen may help with overall tissue oxygenation but not rhythm stabilization, and while consulting a cardiologist is important for ongoing care, it may not provide the urgent intervention needed in cases of unstable rhythms.

10. Which condition requires immediate administration of glucose?

A. Dehydration

B. Hypoglycemia

C. Sepsis

D. Cardiac arrest

The condition that necessitates the immediate administration of glucose is hypoglycemia. Hypoglycemia occurs when blood sugar levels drop below normal, leading to symptoms such as confusion, irritability, sweating, tremors, and in severe cases, loss of consciousness or seizures. These symptoms are due to the brain's reliance on glucose for energy; a deficit can quickly impair neurological function. In a pediatric emergency setting, recognizing hypoglycemia as a potentially life-threatening situation is crucial. Administering glucose can rapidly reverse these symptoms and restore normal brain function, making it a priority intervention. In contrast, while dehydration, sepsis, and cardiac arrest are serious conditions that require prompt assessment and treatment, they do not specifically require the immediate administration of glucose as part of their initial management. Dehydration typically involves the restoration of fluids and electrolytes, sepsis treatments may include antibiotics and fluids, and cardiac arrest necessitates advanced cardiac life support protocols.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cpen.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE