CERTIFIED REGISTERED NURSE ANESTHETIST (CRNA) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://crna.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Can tracheal intubation fully prevent aspiration?

A. Yes, it completely prevents aspiration

B. No, micro aspiration can still occur

C. Only in patients with strong gag reflexes

D. Only during emergency procedures

2. Why should patients discontinue certain medications before surgery?

A. To improve the effectiveness of the anesthetics

B. To ensure a higher tolerance to pain

C. To reduce the risk of adverse reactions and complications during anesthesia

D. To prevent postoperative infections

3. What is the most common type of tracheo-esophageal fistula?

A. Tracheoesophageal junction anomaly

B. Blind esophageal pouch with a fistulous connection to the trachea

C. Esophageal atresia with a distal tracheoesophageal fistula

D. Proximal tracheoesophageal fistula

4. What defines a full stomach in the context of tracheal intubation?

A. Presence of gastric contents or recent food intake

B. Increased esophageal pressure

C. Decreased lower esophageal pressure

D. Absence of gastrointestinal motility

5. Which are the only "fail safe" signs for the correct placement of an endotracheal tube?

A. Visual confirmation of tube placement

B. Persistence of appropriate levels of end-tidal carbon dioxide

C. Palpation of the cuff in the trachea

D. Presence of breath sounds bilaterally

SAMPLE

6. In patients receiving blood pressure medications, what should be the primary focus of the CRNA during anesthesia?

A. Maintaining hydration only

B. Controlling blood sugar levels

C. Monitoring and managing hemodynamic stability

D. Administering pain relief

7. Why is temperature monitoring critical during surgery?

A. To ensure optimal communication with staff

B. To monitor the progress of anesthesia

C. To prevent hypothermia and ensure optimal metabolic function

D. To measure blood loss

8. What medications are typically included in a balanced anesthesia technique?

A. Antibiotics, Hormones, Local anesthetics, and Opioids

B. Opioids, sedatives, neuromuscular blockers, and inhalational anesthetics

C. Only inhalational anesthetics and opioids

D. Only sedatives and neuromuscular blockers

9. What is the range for normal pulmonary artery pressure (PAP)?

A. 10-20/2-10 mmHg

B. 15-30/5-15 mmHg

C. 20-25/10-15 mmHg

D. 25-35/10-20 mmHg

10. What effect does epinephrine have on systolic blood pressure?

A. Decreased systolic BP

B. No effect on systolic BP

C. Increased systolic BP

D. Variable effects on systolic BP

SAMPLE

Answers

SAMPLE

1. B
2. C
3. B
4. A
5. B
6. C
7. C
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. Can tracheal intubation fully prevent aspiration?

A. Yes, it completely prevents aspiration

B. No, micro aspiration can still occur

C. Only in patients with strong gag reflexes

D. Only during emergency procedures

Tracheal intubation is a procedure used to secure the airway and facilitate ventilation. While it significantly reduces the risk of aspiration by creating a barrier between the trachea and the oral cavity, it does not completely prevent aspiration. This is because micro-aspiration can still occur, particularly if there are factors such as anatomical variations or dysfunction of the upper airway protective reflexes. Micro-aspiration refers to small amounts of fluid or secretions entering the trachea and lungs, which can happen despite the presence of an endotracheal tube. In situations where there is increased intra-abdominal pressure, such as during positive pressure ventilation, or in certain patient populations with impaired swallowing or protective airway reflexes, the risk for aspiration may still exist, leading to potential pulmonary complications. Understanding that tracheal intubation greatly decreases the likelihood of significant aspiration episodes but does not entirely eliminate the risk is essential for managing patients' airways effectively.

2. Why should patients discontinue certain medications before surgery?

A. To improve the effectiveness of the anesthetics

B. To ensure a higher tolerance to pain

C. To reduce the risk of adverse reactions and complications during anesthesia

D. To prevent postoperative infections

Discontinuing certain medications before surgery is essential primarily to reduce the risk of adverse reactions and complications during anesthesia. Some medications can interact negatively with anesthetic agents, leading to increased cardiac or respiratory complications, prolonged recovery times, or ineffective analgesia. For instance, anticoagulants can increase bleeding risk, while certain herbal supplements may affect blood pressure or lead to unpredictable responses to anesthesia. Ensuring patient safety is a primary concern during any surgical procedure, and understanding the implications of medication use is a crucial part of preoperative assessments. By identifying and managing these risks beforehand, healthcare providers can create a safer surgical environment for patients.

3. What is the most common type of tracheo-esophageal fistula?

A. Tracheoesophageal junction anomaly

B. Blind esophageal pouch with a fistulous connection to the trachea

C. Esophageal atresia with a distal tracheoesophageal fistula

D. Proximal tracheoesophageal fistula

The most prevalent type of tracheo-esophageal fistula is characterized as a blind esophageal pouch with a fistulous connection to the trachea. This specific configuration typically arises due to a developmental defect during the embryonic stage, where the esophagus does not form a complete passage, resulting in the upper part of the esophagus terminating in a pouch. The pouch then connects to the trachea, leading to the abnormal communication between the respiratory and digestive tracts. This condition is commonly associated with esophageal atresia, where the esophagus is obstructed. The pervasive nature of this type of fistula can lead to significant clinical complications, such as aspiration pneumonia, since swallowed materials can be diverted into the trachea. Understanding the anatomical and physiological implications of this anomaly is crucial for anesthetic management and surgical intervention, which are key components of a CRNA's responsibilities. While other types of tracheo-esophageal fistulas do exist, such as those involving various configurations of the esophagus and trachea, the blind pouch with a connection to the trachea stands out as the most frequently encountered form in clinical practice.

SAMPLE

4. What defines a full stomach in the context of tracheal intubation?

A. Presence of gastric contents or recent food intake

B. Increased esophageal pressure

C. Decreased lower esophageal pressure

D. Absence of gastrointestinal motility

A full stomach in the context of tracheal intubation refers to the presence of gastric contents or recent food intake. This is an important consideration in anesthesia practice because a full stomach increases the risk of aspiration during intubation. When a patient has not adequately fasted before undergoing anesthesia, there is a potential for stomach contents to regurgitate and enter the airway, leading to serious complications. Understanding the implications of a full stomach is critical for CRNAs, as it affects the management of airway and anesthetic techniques. Protocols may include preoperative fasting guidelines to mitigate this risk, highlighting the necessity for all healthcare providers involved in the perioperative care of patients to recognize signs of a full stomach. The other options may be relevant in other contexts related to gastrointestinal physiology or different aspects of patient health but do not define what constitutes a full stomach in the context of anesthesia and tracheal intubation. For instance, increases or decreases in esophageal pressure relate more to the function of the esophagus rather than the status of gastric contents. Similarly, the absence of gastrointestinal motility pertains to gut function but does not directly indicate fullness of the stomach.

5. Which are the only "fail safe" signs for the correct placement of an endotracheal tube?

A. Visual confirmation of tube placement

B. Persistence of appropriate levels of end-tidal carbon dioxide

C. Palpation of the cuff in the trachea

D. Presence of breath sounds bilaterally

The identification of the persistence of appropriate levels of end-tidal carbon dioxide is considered a "fail safe" sign for verifying correct endotracheal tube placement. When an endotracheal tube is positioned correctly within the trachea, it allows for the proper exchange of gases between the lungs and the environment. The presence of carbon dioxide in the exhaled breath indicates that the tube is effectively situated where it can access the alveoli for adequate ventilation. The measurement of end-tidal carbon dioxide is a reliable indicator, as it not only confirms the position of the tube within the trachea but also reflects the patient's respiratory status. This is particularly critical in assessing their ventilation, as any significant changes or absence of carbon dioxide can suggest displacement or misplacement of the tube, prompting necessary interventions. While visual confirmation of tube placement, palpation of the cuff in the trachea, and presence of breath sounds bilaterally are important checks in the intubation process, they do not provide as definitive a measure of proper placement as the continuous monitoring of end-tidal carbon dioxide. Each of these methods has limitations: visual confirmation may be ambiguous, palpation could lead to misinterpretation based on anatomical variations, and the presence of breath sounds

SAMPLE

6. In patients receiving blood pressure medications, what should be the primary focus of the CRNA during anesthesia?

A. Maintaining hydration only

B. Controlling blood sugar levels

C. Monitoring and managing hemodynamic stability

D. Administering pain relief

In patients receiving blood pressure medications, the primary focus of the CRNA during anesthesia should be on monitoring and managing hemodynamic stability. This is especially important because these patients may have compromised cardiovascular function due to their underlying conditions and the effects of their medications. Hemodynamic stability encompasses the assessment of blood pressure, heart rate, and overall cardiac output, which can be significantly affected by anesthetic agents and the surgical procedure. Carefully monitoring these parameters allows the CRNA to make timely interventions to prevent hypotension, hypertension, or other cardiovascular complications that might arise during anesthesia. Other factors, such as hydration, blood sugar levels, and pain relief, are also significant components of preoperative and intraoperative care but they are part of a broader focus. While maintaining hydration is crucial for overall patient health and can affect hemodynamic status, it is not the sole focus when managing patients on blood pressure medications. Similarly, while controlling blood sugar levels is essential for diabetic patients, it is not as immediate a concern in the context of anesthesia as maintaining hemodynamic stability. Administering pain relief is vital for postoperative comfort but again takes a secondary role compared to ensuring that the patient's cardiovascular system remains stable throughout the anesthesia process.

7. Why is temperature monitoring critical during surgery?

A. To ensure optimal communication with staff

B. To monitor the progress of anesthesia

C. To prevent hypothermia and ensure optimal metabolic function

D. To measure blood loss

Temperature monitoring during surgery is essential because it plays a crucial role in preventing hypothermia and ensuring optimal metabolic function. Maintaining normothermia is vital for several physiological reasons. The body's temperature affects enzymatic processes and metabolic rates; thus, a drop in body temperature can lead to slowed metabolism, impaired coagulation, and increased risk of perioperative complications. Hypothermia can also result from exposure to the cold environment of the operating room, the effects of anesthesia, and increased heat loss during surgery. By continuously monitoring the patient's temperature, the surgical team can implement warming measures like heated blankets or fluid warmers when necessary. This helps maintain the patient's core body temperature, which is crucial for stable hemodynamics and better overall surgical outcomes. Other aspects mentioned, such as communication with staff, monitoring the progress of anesthesia, or measuring blood loss, are important in their own right but are not the primary reasons for temperature monitoring during surgical procedures.

SAMPLE

8. What medications are typically included in a balanced anesthesia technique?

A. Antibiotics, Hormones, Local anesthetics, and Opioids

B. Opioids, sedatives, neuromuscular blockers, and inhalational anesthetics

C. Only inhalational anesthetics and opioids

D. Only sedatives and neuromuscular blockers

The practice of balanced anesthesia involves using a combination of medications to achieve optimal conditions for surgery while minimizing side effects. The correct choice includes opioids, sedatives, neuromuscular blockers, and inhalational anesthetics, which together cover different aspects of anesthesia management. Opioids provide effective analgesia and help to manage pain during and after the surgical procedure. They are a crucial component for ensuring patient comfort and reducing the perception of pain. Sedatives, on the other hand, help to maintain sedation and reduce anxiety, allowing for a more controlled and comfortable surgical experience for the patient. These medications also assist in achieving amnesia during the procedure. Neuromuscular blockers are essential for achieving muscle relaxation, facilitating intubation, and providing optimal surgical conditions by allowing the surgeon access to areas that may be difficult to operate on without muscle tone interference. Finally, inhalational anesthetics are typically used to maintain anesthesia throughout the procedure, providing a reliable means to sustain unconsciousness and analgesia during surgery. This combination of agents ensures comprehensive coverage of anesthesia needs, enhancing the safety and effectiveness of the surgical process while minimizing the risks associated with using a single-agent approach.

9. What is the range for normal pulmonary artery pressure (PAP)?

A. 10-20/2-10 mmHg

B. 15-30/5-15 mmHg

C. 20-25/10-15 mmHg

D. 25-35/10-20 mmHg

The normal range for pulmonary artery pressure (PAP) is widely accepted to be approximately 15-30 mmHg for systolic pressure and 5-15 mmHg for diastolic pressure. This range reflects the typical pressures seen in a healthy individual's pulmonary circulation, which is significantly lower than systemic arterial pressures. Understanding the normal PAP is crucial for anesthetists and healthcare providers in assessing cardiac function and diagnosing pulmonary hypertension. Elevated pressures can indicate underlying pathologies such as conditions leading to increased pulmonary vascular resistance, while lower pressures are often less concerning. When monitoring patients, particularly in anesthesia settings, it is important to know these values to make appropriate clinical decisions. The specified option provides a range that accurately encompasses these typical values observed in clinical practice, making it the best choice among those provided.

SAMPLE

10. What effect does epinephrine have on systolic blood pressure?

A. Decreased systolic BP

B. No effect on systolic BP

C. Increased systolic BP

D. Variable effects on systolic BP

Epinephrine, also known as adrenaline, is a powerful vasopressor and inotropic agent commonly used in emergency situations and anesthesia. Its primary mechanism of action involves stimulating both alpha and beta-adrenergic receptors. The stimulation of beta-1 adrenergic receptors in the heart increases heart rate and contractility, which typically leads to an increase in cardiac output. Additionally, the stimulation of beta-2 receptors causes vasodilation in certain vascular beds, while alpha-1 receptor activation can cause vasoconstriction in others. When considering its effects on systolic blood pressure, the increased cardiac output due to enhanced heart rate and contractility, combined with the increased vascular resistance from alpha-1 receptor stimulation, leads to an overall increase in systolic blood pressure. This is especially important in critical care settings where maintaining adequate perfusion pressure is crucial for organ function. Therefore, the correct answer highlights that epinephrine increases systolic blood pressure through its multifaceted actions on the cardiovascular system.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://crna.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE