CERTIFIED REGISTERED NURSE INFUSION (CRNI) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://crni.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What serious side effect can occur with rapid infusion of Ticarcillin?

A. Renal failure

B. Seizures

C. Anaphylaxis

D. Optic neuritis

2. Why is Clindamycin infusion potentially dangerous if administered rapidly?

A. It can cause seizures.

B. It can lead to cardiac arrest.

C. It causes respiratory failure.

D. It can result in suppressed immune function.

3. What is the significance of potassium replacement in high doses of Lasix?

A. To prevent kidney damage

B. To avoid hypokalemia

C. To enhance medication effects

D. To reduce fluid retention

4. Which common side effect is associated with excess administration of dextrose solutions?

A. Electrolyte imbalance

B. Hyperglycemia

C. Fluid overload

D. Nutritional deficiency

5. Which antineoplastic agent has a recommended total cumulative dose limit of 1400 mg/m²?

A. Carmustine

B. Gemcitabine

C. Paclitaxel

D. Docetaxel

SAMPLE

6. What is the primary cause of bacterial phlebitis?

A. Air contamination

B. Skin contamination

C. Improper catheter placement

D. Drug abnormalities

7. What size filter is required for administering Amphotericin B?

A. 0.45 micron

B. 1 micron

C. 5 micron

D. 10 micron

8. Which of the following medications is used to relax bronchial tubes?

A. IVIG

B. Theophylline

C. Caspofungin

D. Flagyl

9. Which third generation cephalosporin is indicated for use with pseudomonas?

A. Cefotaxime

B. Ceftriaxone

C. Ceftazidime

D. Cefepime

10. Which of the following is a critical safety measure when administering Amphotericin B?

A. Use a one micron filter

B. Administer with a saline bolus

C. Monitor vital signs every 15 minutes

D. Give as a fast intravenous push

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. B
5. A
6. B
7. B
8. B
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. What serious side effect can occur with rapid infusion of Ticarcillin?

A. Renal failure

B. Seizures

C. Anaphylaxis

D. Optic neuritis

Rapid infusion of Ticarcillin can lead to seizures due to its neurotoxic potential at high serum concentrations. When infused quickly, the medication can accumulate in the body and reach levels that may cause central nervous system stimulation, resulting in seizure activity. This response is particularly important to consider in patients who may already be at risk for seizures or those with compromised renal function, as they might have impaired drug clearance. While renal failure can happen with various medications, it is not a direct result of rapid infusion of Ticarcillin itself. Anaphylaxis is an allergic reaction that can occur but is not specifically associated with the rapid infusion of Ticarcillin. Optic neuritis is not a recognized side effect of Ticarcillin at any infusion rate. Understanding the specific side effects related to the infusion rate helps healthcare professionals manage medication administration safely and effectively.

2. Why is Clindamycin infusion potentially dangerous if administered rapidly?

A. It can cause seizures.

B. It can lead to cardiac arrest.

C. It causes respiratory failure.

D. It can result in suppressed immune function.

Clindamycin infusion can potentially be dangerous if administered rapidly primarily because it can lead to cardiac arrest. Rapid infusion of Clindamycin can cause significant cardiovascular effects, including hypotension and other adverse reactions that may compromise heart function. The risk of severe cardiac complications highlights the importance of adhering to recommended infusion rates when administering this medication. In this particular scenario, the concern centers around the physiological impact of a fast infusion rate, which can overwhelm the body's ability to tolerate the influx of medication, resulting in serious, life-threatening outcomes. Understanding the pharmacodynamics and proper administration protocols of Clindamycin is essential for healthcare providers to minimize such risks.

3. What is the significance of potassium replacement in high doses of Lasix?

A. To prevent kidney damage

B. To avoid hypokalemia

C. To enhance medication effects

D. To reduce fluid retention

Potassium replacement is particularly significant when high doses of Lasix (furosemide) are used due to the risk of hypokalemia, which is a condition characterized by low potassium levels in the blood. Lasix is a potent loop diuretic that promotes the excretion of sodium and water through the kidneys. However, this increased diuresis can also lead to the loss of electrolytes, including potassium. When patients are on high doses of Lasix, the risk of hypokalemia becomes greater due to the increased urinary excretion of potassium. Hypokalemia can lead to serious complications, such as cardiac arrhythmias, muscle weakness, and metabolic disturbances. Therefore, monitoring potassium levels and providing potassium replacement as necessary is crucial in patients receiving high doses of Lasix to maintain electrolyte balance and prevent these potentially harmful consequences. While other considerations, such as kidney protection, medication efficacy, and fluid retention, are important in the context of Lasix therapy, the primary concern when administering high doses is ensuring that potassium levels remain within an appropriate range to avoid the risks associated with its deficiency.

SAMPLE

4. Which common side effect is associated with excess administration of dextrose solutions?

A. Electrolyte imbalance

B. Hyperglycemia

C. Fluid overload

D. Nutritional deficiency

Hyperglycemia occurs as a common side effect when there is an excess administration of dextrose solutions. Dextrose is a form of glucose, and when administered in large amounts, it can lead to elevated blood sugar levels. This is particularly significant in patients who may have underlying conditions such as diabetes or impaired glucose metabolism, where their bodies cannot adequately process the increased glucose load. When dextrose is infused, it enters the bloodstream rapidly, resulting in an increase in blood glucose levels that can exceed the normal range. This spike can provoke a physiological response, leading to symptoms such as increased thirst, frequent urination, and fatigue. Monitoring blood glucose levels is essential when administering dextrose solutions to prevent hyperglycemia and to ensure patient safety. In comparison, while electrolyte imbalance, fluid overload, and nutritional deficiency can occur with various fluid and nutrient infusions, they are not exclusively tied to the excess administration of dextrose solutions in the same direct and significant way that hyperglycemia is.

5. Which antineoplastic agent has a recommended total cumulative dose limit of 1400 mg/m²?

A. Carmustine

B. Gemcitabine

C. Paclitaxel

D. Docetaxel

Carmustine is the antineoplastic agent that is associated with a recommended total cumulative dose limit of 1400 mg/m². This dose limit is significant because exceeding it can result in severe toxicities, particularly myelosuppression leading to a high risk of infections and bleeding complications. Carmustine is an alkylating agent that is often used in the treatment of certain types of cancers, including brain tumors and lymphomas, and its unique formulation allows for prolonged exposure in tissues. Understanding cumulative dose limits is crucial for patient safety and for preventing long-term complications from chemotherapy. Being aware of such dosing guidelines, especially for drugs like Carmustine, helps healthcare professionals manage therapy effectively and mitigate potential side effects.

6. What is the primary cause of bacterial phlebitis?

A. Air contamination

B. Skin contamination

C. Improper catheter placement

D. Drug abnormalities

Bacterial phlebitis is primarily caused by skin contamination. When a catheter is inserted into a vein, the potential exists for bacteria from the skin to enter the bloodstream and infect the site. The skin is a natural habitat for many bacteria, and even with careful antiseptic techniques, some bacteria may still be present. This can lead to localized inflammation and infection at the site of the catheter, resulting in phlebitis. Factors such as air contamination, improper catheter placement, and drug abnormalities can contribute to other complications or infections but are not the primary drivers of bacterial phlebitis. Skin contamination is a direct and common route for bacteria to gain access to the vascular system during the insertion and maintenance of intravenous devices. Therefore, ensuring proper skin antisepsis before catheter placement is crucial in preventing bacterial phlebitis.

SAMPLE

7. What size filter is required for administering Amphotericin B?

A. 0.45 micron

B. 1 micron

C. 5 micron

D. 10 micron

The requirement for a 1 micron filter for administering Amphotericin B is based on the need to prevent the infusion of any potential particulate matter or fungal contaminants that could arise from the formulation. Amphotericin B is known to aggregate, and using a filter of this size helps to trap any particles or aggregates that could cause adverse reactions or complications during administration. Filters that are smaller than 1 micron might be too restrictive, potentially causing issues with flow rates and leading to incomplete administration of the medication. Filters larger than 1 micron are not adequate to effectively trap the particulates associated with Amphotericin B, which could lead to potential complications for the patient. Thus, the 1 micron filter strikes an appropriate balance, ensuring safety while allowing for proper drug delivery.

8. Which of the following medications is used to relax bronchial tubes?

A. IVIG

B. Theophylline

C. Caspofungin

D. Flagyl

Theophylline is a medication that serves to relax the bronchial tubes, which can help to improve airflow in conditions such as asthma and chronic obstructive pulmonary disease (COPD). It acts as a bronchodilator by inhibiting certain enzymes and promoting the relaxation of smooth muscles within the respiratory tract, thereby widening the airways and enhancing breathing. This mechanism of action is crucial for managing symptoms and providing relief to patients experiencing bronchoconstriction. Theophylline can also have anti-inflammatory effects, adding further benefit to respiratory management. The other medications listed, while important in their respective categories, do not have bronchodilatory properties. IVIG (Intravenous Immunoglobulin) is primarily used for immunological conditions, Caspofungin is an antifungal medication, and Flagyl (Metronidazole) is an antibiotic used for treating various infections. None of these medications are indicated for relaxing bronchial tubes.

SAMPLE

9. Which third generation cephalosporin is indicated for use with pseudomonas?

A. Cefotaxime

B. Ceftriaxone

C. Ceftazidime

D. Cefepime

Ceftazidime is a third-generation cephalosporin that is specifically indicated for use against Pseudomonas aeruginosa, a common and opportunistic pathogen known for its resistance to many antibiotics. This particular cephalosporin is effective due to its ability to penetrate the outer membrane of Gram-negative bacteria, including Pseudomonas, and its unique structure that enhances its activity against this organism. Ceftazidime is often chosen for serious infections caused by Pseudomonas, particularly in immunocompromised patients or those with healthcare-associated infections. Its effectiveness in treating infections such as pneumonia, urinary tract infections, and bacteremia linked to this resistant pathogen makes it a critical option in the antimicrobial arsenal. The other third-generation cephalosporins listed do not have reliable activity against Pseudomonas. For instance, cefotaxime and ceftriaxone primarily target strains of Streptococcus pneumoniae and other Gram-positive organisms, while cefepime, although it is a fourth-generation cephalosporin recognized for its broader spectrum including Pseudomonas, is not part of the third generation being asked about in this context. Thus, cefotaxime, ceftriaxone, and cef

10. Which of the following is a critical safety measure when administering Amphotericin B?

A. Use a one micron filter

B. Administer with a saline bolus

C. Monitor vital signs every 15 minutes

D. Give as a fast intravenous push

Using a one micron filter is a critical safety measure when administering Amphotericin B due to the risk of potential particulate matter and fungal contamination in the medication. Amphotericin B is a lipid-based antifungal agent, and its formulation can lead to the presence of aggregates or particles that could cause adverse reactions if introduced into the bloodstream. The one micron filter acts as a safety barrier that traps these particles, ensuring the medication delivered is safe and effective. This filtration step is particularly important given the serious nature of infections treated with Amphotericin B, such as systemic fungal infections, where any contaminants could pose significant risks to patient safety. Proper filtration helps in administering the drug without introducing harmful particles that could lead to complications. While other measures, like hydration or monitoring vital signs, are important in the context of administering Amphotericin B, the specific use of a one micron filter directly addresses the prevention of particulates entering the bloodstream.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://crni.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE